Full wwPDB NMR Structure Validation Report (i)

Apr 27, 2016 – 03:30 AM BST

PDB ID	:	2MUM
Title	:	Solution structure of the PHD domain of Yeast YNG2
Authors	:	Taeb, S.; Kaustov, L.; Lemak, A.; Farhadi, S.; Sheng, Y.
Deposited on	:	2014-09-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 94%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	50	92% 6% •

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:223-A:271 (49)	0.46	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 6, 7, 8, 10, 11, 12, 13, 14, 15, 16, 18, 19
2	5, 17, 20
Single-model clusters	9

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 788 atoms, of which 373 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Chromatin modification-related protein YNG2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	50	786	266	373	64	75	8	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
			Total	Zn
2	A	2	2	2

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.2 Score per residue for model 2

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.3 Score per residue for model 3

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.4 Score per residue for model 4

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.5 Score per residue for model 5

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.7 Score per residue for model 7

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.8 Score per residue for model 8

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.9 Score per residue for model 9

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.10 Score per residue for model 10

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.11 Score per residue for model 11

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.12 Score per residue for model 12

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.13 Score per residue for model 13

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.14 Score per residue for model 14

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.15 Score per residue for model 15

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.16 Score per residue for model 16

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.17 Score per residue for model 17

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.18 Score per residue for model 18

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:



4.2.19 Score per residue for model 19

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

4.2.20 Score per residue for model 20

- Molecule 1: Chromatin modification-related protein YNG2

Chain A:

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
FMCGUI	refinement	
CYANA	structure solution	
CNSSOLVE	refinement	
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2mum_cs.str
Number of chemical shift lists	1
Total number of shifts	602
Number of shifts mapped to atoms	602
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	94%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	406	364	364	1±1
All	All	8160	7280	7280	28

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:231:SER:HB3	1:A:235:MET:HB3	0.55	1.78	2	2
1:A:238:CYS:HB3	1:A:246:GLU:HA	0.52	1.80	4	6
1:A:240:GLY:HA3	1:A:268:CYS:SG	0.50	2.47	17	3
1:A:224:TYR:HB3	1:A:245:TYR:HA	0.48	1.85	7	2
1:A:243:CYS:SG	1:A:246:GLU:HA	0.47	2.49	9	3
1:A:238:CYS:SG	1:A:264:TYR:HA	0.47	2.48	9	1
1:A:227:CYS:SG	1:A:229:ARG:HB2	0.45	2.51	10	1
1:A:249:HIS:CD2	1:A:252:CYS:HB2	0.44	2.48	17	3
1:A:265:CYS:SG	1:A:267:GLU:OE2	0.43	2.75	8	4
1:A:265:CYS:SG	1:A:267:GLU:OE1	0.42	2.78	10	1
1:A:265:CYS:SG	1:A:267:GLU:HG2	0.41	2.54	2	1
1:A:265:CYS:SG	1:A:267:GLU:HB2	0.41	2.55	15	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	48/50 (96%)	41±2 (86±3%)	6±2 (13±3%)	1±1 (2±1%)	17 61
All	All	960/1000 (96%)	821 (86%)	124 (13%)	15 (2%)	17 61

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	228	GLN	8
1	A	266	PRO	7

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	45/46 (98%)	42±1 (93±2%)	3±1 (7±2%)	22 68
All	All	900/920 (98%)	834 (93%)	66 (7%)	22 68

All 13 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	249	HIS	20
1	A	252	CYS	13
1	A	267	GLU	9
1	A	244	LYS	7
1	A	260	LYS	3
1	A	269	LYS	2
1	A	229	ARG	2
1	A	234	GLU	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	266	PRO	2
1	A	256	LYS	2
1	A	235	MET	2
1	A	223	LEU	1
1	A	257	GLU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 94% for the well-defined parts and 93% for the entire structure.

7.1 Chemical shift list 1

File name: 2mum_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	602
Number of shifts mapped to atoms	602
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	7

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	50	0.10 ± 0.27	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	47	0.05 ± 0.21	None needed (< 0.5 ppm)
$^{13}C'$	42	0.20 ± 0.32	None needed (< 0.5 ppm)
^{15}N	44	-0.25 ± 0.54	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 94%, i.e. 574 atoms were assigned a chemical shift out of a possible 613. 0 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	228/237 (96%)	93/94 (99%)	91/98 (93%)	44/45 (98%)
Sidechain	259/286 (91%)	166/173 (96%)	90/103 (87%)	3/10 (30%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	87/90 (97%)	45/47 (96%)	40/40 (100%)	2/3 (67%)
Overall	574/613 (94%)	304/314 (97%)	221/241 (92%)	49/58 (84%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 93%, i.e. 578 atoms were assigned a chemical shift out of a possible 622. 0 out of 5 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	230/242 (95%)	94/96 (98%)	92/100 (92%)	44/46 (96%)
Sidechain	261/290 (90%)	167/175 (95%)	91/105 (87%)	3/10 (30%)
Aromatic	87/90 (97%)	45/47 (96%)	40/40 (100%)	2/3 (67%)
Overall	578/622 (93%)	306/318 (96%)	223/245 (91%)	49/59 (83%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	234	GLU	HB3	0.41	3.10 – 0.90	-7.2
1	A	237	ALA	N	101.05	141.07 – 105.37	-6.2
1	A	271	GLU	N	99.31	138.24 – 103.14	-6.1
1	A	234	GLU	HB2	0.81	3.08 – 0.98	-5.8
1	A	237	ALA	HB1	0.11	2.61 – 0.11	-5.0
1	A	237	ALA	HB2	0.11	2.61 – 0.11	-5.0
1	A	237	ALA	HB3	0.11	2.61 – 0.11	-5.0

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:


RCI
0.3
0.2
0.1
0
222:THR
224:TYR
226:PHE
228:GLN
230:VAL
232:PHE
234:GLU
236:VAL
238:CYS
240:GLY
242:ASN
244:LYS
246:GLU
248:PHE
250:TYR
252:CYS
254:ASN
256:LYS
258:PRO
260:LYS
262:THR
264:TYR
266:PRO
268:CYS
270:ILE