



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:49 PM GMT

PDB ID : 4MVS
Title : Structural Basis for Ca²⁺ Selectivity of a Voltage-gated Calcium Channel
Authors : Tang, L.; Gamal El-Din, T.M.; Payandeh, J.; Martinez, G.Q.; Heard, T.M.; Scheuer, T.; Zheng, N.; Catterall, W.A.
Deposited on : 2013-09-24
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	 58% 31% 8%
1	B	237	 63% 28% 8%
1	C	237	 60% 27% 5% 8%
1	D	237	 62% 27% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PX4	A	1306	-	-	-	X
3	PX4	B	1304	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1798	1224	268	295	11			
1	B	219	Total	C	N	O	S	0	0	0
			1798	1224	268	295	11			
1	C	219	Total	C	N	O	S	0	0	0
			1798	1224	268	295	11			
1	D	219	Total	C	N	O	S	0	0	0
			1798	1224	268	295	11			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	983	MET	-	EXPRESSION TAG	UNP A8EVM5
A	984	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	985	TYR	-	EXPRESSION TAG	UNP A8EVM5
A	986	LYS	-	EXPRESSION TAG	UNP A8EVM5
A	987	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	988	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	989	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	990	ASP	-	EXPRESSION TAG	UNP A8EVM5
A	991	LYS	-	EXPRESSION TAG	UNP A8EVM5
A	992	GLY	-	EXPRESSION TAG	UNP A8EVM5
A	993	SER	-	EXPRESSION TAG	UNP A8EVM5
A	994	LEU	-	EXPRESSION TAG	UNP A8EVM5
A	995	VAL	-	EXPRESSION TAG	UNP A8EVM5
A	996	PRO	-	EXPRESSION TAG	UNP A8EVM5
A	997	ARG	-	EXPRESSION TAG	UNP A8EVM5
A	998	GLY	-	EXPRESSION TAG	UNP A8EVM5
A	999	SER	-	EXPRESSION TAG	UNP A8EVM5
A	1000	HIS	-	EXPRESSION TAG	UNP A8EVM5
A	1177	ASP	GLU	CONFLICT	UNP A8EVM5
A	1178	ASP	SER	CONFLICT	UNP A8EVM5
A	1181	ASP	MET	CONFLICT	UNP A8EVM5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1217	CYS	ILE	CONFLICT	UNP A8EVM5
B	983	MET	-	EXPRESSION TAG	UNP A8EVM5
B	984	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	985	TYR	-	EXPRESSION TAG	UNP A8EVM5
B	986	LYS	-	EXPRESSION TAG	UNP A8EVM5
B	987	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	988	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	989	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	990	ASP	-	EXPRESSION TAG	UNP A8EVM5
B	991	LYS	-	EXPRESSION TAG	UNP A8EVM5
B	992	GLY	-	EXPRESSION TAG	UNP A8EVM5
B	993	SER	-	EXPRESSION TAG	UNP A8EVM5
B	994	LEU	-	EXPRESSION TAG	UNP A8EVM5
B	995	VAL	-	EXPRESSION TAG	UNP A8EVM5
B	996	PRO	-	EXPRESSION TAG	UNP A8EVM5
B	997	ARG	-	EXPRESSION TAG	UNP A8EVM5
B	998	GLY	-	EXPRESSION TAG	UNP A8EVM5
B	999	SER	-	EXPRESSION TAG	UNP A8EVM5
B	1000	HIS	-	EXPRESSION TAG	UNP A8EVM5
B	1177	ASP	GLU	CONFLICT	UNP A8EVM5
B	1178	ASP	SER	CONFLICT	UNP A8EVM5
B	1181	ASP	MET	CONFLICT	UNP A8EVM5
B	1217	CYS	ILE	CONFLICT	UNP A8EVM5
C	983	MET	-	EXPRESSION TAG	UNP A8EVM5
C	984	ASP	-	EXPRESSION TAG	UNP A8EVM5
C	985	TYR	-	EXPRESSION TAG	UNP A8EVM5
C	986	LYS	-	EXPRESSION TAG	UNP A8EVM5
C	987	ASP	-	EXPRESSION TAG	UNP A8EVM5
C	988	ASP	-	EXPRESSION TAG	UNP A8EVM5
C	989	ASP	-	EXPRESSION TAG	UNP A8EVM5
C	990	ASP	-	EXPRESSION TAG	UNP A8EVM5
C	991	LYS	-	EXPRESSION TAG	UNP A8EVM5
C	992	GLY	-	EXPRESSION TAG	UNP A8EVM5
C	993	SER	-	EXPRESSION TAG	UNP A8EVM5
C	994	LEU	-	EXPRESSION TAG	UNP A8EVM5
C	995	VAL	-	EXPRESSION TAG	UNP A8EVM5
C	996	PRO	-	EXPRESSION TAG	UNP A8EVM5
C	997	ARG	-	EXPRESSION TAG	UNP A8EVM5
C	998	GLY	-	EXPRESSION TAG	UNP A8EVM5
C	999	SER	-	EXPRESSION TAG	UNP A8EVM5
C	1000	HIS	-	EXPRESSION TAG	UNP A8EVM5
C	1177	ASP	GLU	CONFLICT	UNP A8EVM5

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1178	ASP	SER	CONFLICT	UNP A8EVM5
C	1181	ASP	MET	CONFLICT	UNP A8EVM5
C	1217	CYS	ILE	CONFLICT	UNP A8EVM5
D	983	MET	-	EXPRESSION TAG	UNP A8EVM5
D	984	ASP	-	EXPRESSION TAG	UNP A8EVM5
D	985	TYR	-	EXPRESSION TAG	UNP A8EVM5
D	986	LYS	-	EXPRESSION TAG	UNP A8EVM5
D	987	ASP	-	EXPRESSION TAG	UNP A8EVM5
D	988	ASP	-	EXPRESSION TAG	UNP A8EVM5
D	989	ASP	-	EXPRESSION TAG	UNP A8EVM5
D	990	ASP	-	EXPRESSION TAG	UNP A8EVM5
D	991	LYS	-	EXPRESSION TAG	UNP A8EVM5
D	992	GLY	-	EXPRESSION TAG	UNP A8EVM5
D	993	SER	-	EXPRESSION TAG	UNP A8EVM5
D	994	LEU	-	EXPRESSION TAG	UNP A8EVM5
D	995	VAL	-	EXPRESSION TAG	UNP A8EVM5
D	996	PRO	-	EXPRESSION TAG	UNP A8EVM5
D	997	ARG	-	EXPRESSION TAG	UNP A8EVM5
D	998	GLY	-	EXPRESSION TAG	UNP A8EVM5
D	999	SER	-	EXPRESSION TAG	UNP A8EVM5
D	1000	HIS	-	EXPRESSION TAG	UNP A8EVM5
D	1177	ASP	GLU	CONFLICT	UNP A8EVM5
D	1178	ASP	SER	CONFLICT	UNP A8EVM5
D	1181	ASP	MET	CONFLICT	UNP A8EVM5
D	1217	CYS	ILE	CONFLICT	UNP A8EVM5

- Molecule 2 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cd 1 1	0	0
2	A	2	Total Cd 2 2	0	0
2	D	1	Total Cd 1 1	0	0
2	C	1	Total Cd 1 1	0	0

- Molecule 3 is 1,2-DIMYRISTOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PX4) (formula: C₃₆H₇₃NO₈P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		
3	A	1	Total	C	O	P	0	0
			21	13	7	1		
3	A	1	Total	C	O	P	0	0
			21	13	7	1		
3	A	1	Total	C			0	0
			6	6				
3	A	1	Total	C	O	P	0	0
			10	3	6	1		
3	B	1	Total	C	O	P	0	0
			10	3	6	1		
3	B	1	Total	C	O	P	0	0
			21	13	7	1		
3	B	1	Total	C			0	0
			6	6				
3	B	1	Total	C	O	P	0	0
			10	3	6	1		
3	C	1	Total	C	O	P	0	0
			10	3	6	1		
3	C	1	Total	C	O	P	0	0
			21	13	7	1		
3	C	1	Total	C			0	0
			6	6				
3	C	1	Total	C	O	P	0	0
			10	3	6	1		
3	D	1	Total	C	O	P	0	0
			10	3	6	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C			0	0
			6	6				
3	D	1	Total	C	O	P	0	0
			10	3	6	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			1	1		
4	B	1	Total	O	0	0
			1	1		
4	C	2	Total	O	0	0
			2	2		
4	D	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ion transport protein



- Molecule 1: Ion transport protein




- Molecule 1: Ion transport protein





● Molecule 1: Ion transport protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	178.59Å 178.65Å 131.13Å 90.00° 132.89° 90.00°	Depositor
Resolution (Å)	30.00 – 3.30 28.57 – 3.30	Depositor EDS
% Data completeness (in resolution range)	89.0 (30.00-3.30) 89.2 (28.57-3.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 3.31Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.233 , 0.269 0.236 , 0.273	Depositor DCC
R_{free} test set	2025 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	94.4	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 70.8	EDS
Estimated twinning fraction	0.428 for k,h,-1/2*h-1/2*k-l 0.418 for -k,-h,-1/2*h+1/2*k-l 0.428 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 40376 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7391	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CD, PX4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/1849	0.77	1/2517 (0.0%)
1	B	0.69	1/1849 (0.1%)	0.76	0/2517
1	C	0.66	0/1849	0.77	0/2517
1	D	0.70	0/1849	0.78	0/2517
All	All	0.68	1/7396 (0.0%)	0.77	1/10068 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1159	TRP	CZ3-CH2	-5.54	1.31	1.40

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1188	MET	CG-SD-CE	5.06	108.29	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1798	0	1861	54	0
1	B	1798	0	1861	46	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1798	0	1861	54	0
1	D	1798	0	1861	49	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	68	0	59	3	0
3	B	47	0	40	3	0
3	C	47	0	40	2	0
3	D	26	0	21	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
All	All	7391	0	7604	189	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1162:THR:HG22	1:C:1165:GLU:H	1.39	0.88
1:A:1133:VAL:HG11	1:A:1212:LEU:HD12	1.57	0.86
1:D:1162:THR:HG22	1:D:1165:GLU:H	1.48	0.79
1:D:1133:VAL:HG11	1:D:1212:LEU:HD12	1.65	0.78
1:A:1162:THR:HG22	1:A:1165:GLU:H	1.48	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/237 (92%)	200 (92%)	17 (8%)	0	100	100
1	B	217/237 (92%)	199 (92%)	18 (8%)	0	100	100
1	C	217/237 (92%)	199 (92%)	15 (7%)	3 (1%)	14	50
1	D	217/237 (92%)	193 (89%)	22 (10%)	2 (1%)	21	60
All	All	868/948 (92%)	791 (91%)	72 (8%)	5 (1%)	30	68

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	1093	SER
1	D	1091	THR
1	D	1093	SER
1	C	1115	GLN
1	C	1092	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/218 (93%)	188 (93%)	14 (7%)	19	57
1	B	202/218 (93%)	185 (92%)	17 (8%)	14	46
1	C	202/218 (93%)	188 (93%)	14 (7%)	19	57
1	D	202/218 (93%)	184 (91%)	18 (9%)	12	43
All	All	808/872 (93%)	745 (92%)	63 (8%)	16	50

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1206	THR
1	C	1070	SER
1	D	1162	THR
1	B	1209	MET
1	C	1024	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 5 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PX4	A	1303	-	9,9,45	0.62	0	10,12,53	1.45	2 (20%)
3	PX4	A	1304	-	20,20,45	1.49	2 (10%)	21,24,53	1.52	2 (9%)
3	PX4	A	1305	-	20,20,45	1.48	2 (10%)	21,24,53	1.49	2 (9%)
3	PX4	A	1306	-	5,5,45	0.55	0	4,4,53	0.42	0
3	PX4	A	1307	-	9,9,45	0.85	0	10,12,53	1.60	2 (20%)
3	PX4	B	1302	-	9,9,45	0.74	0	10,12,53	1.21	2 (20%)
3	PX4	B	1303	-	20,20,45	1.47	2 (10%)	21,24,53	1.52	2 (9%)
3	PX4	B	1304	-	5,5,45	0.39	0	4,4,53	0.37	0
3	PX4	B	1305	-	9,9,45	0.64	0	10,12,53	1.71	3 (30%)
3	PX4	C	1302	-	9,9,45	0.71	0	10,12,53	1.37	1 (10%)
3	PX4	C	1303	-	20,20,45	1.48	2 (10%)	21,24,53	1.51	2 (9%)
3	PX4	C	1304	-	5,5,45	0.42	0	4,4,53	0.40	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PX4	C	1305	-	9,9,45	0.63	0	10,12,53	1.18	1 (10%)
3	PX4	D	1302	-	9,9,45	0.54	0	10,12,53	1.24	1 (10%)
3	PX4	D	1303	-	5,5,45	0.49	0	4,4,53	0.26	0
3	PX4	D	1304	-	9,9,45	0.60	0	10,12,53	1.30	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PX4	A	1303	-	-	0/8/8/49	0/0/0/0
3	PX4	A	1304	-	-	0/22/22/49	0/0/0/0
3	PX4	A	1305	-	-	0/22/22/49	0/0/0/0
3	PX4	A	1306	-	-	0/3/3/49	0/0/0/0
3	PX4	A	1307	-	-	0/8/8/49	0/0/0/0
3	PX4	B	1302	-	-	0/8/8/49	0/0/0/0
3	PX4	B	1303	-	-	0/22/22/49	0/0/0/0
3	PX4	B	1304	-	-	0/3/3/49	0/0/0/0
3	PX4	B	1305	-	-	0/8/8/49	0/0/0/0
3	PX4	C	1302	-	-	0/8/8/49	0/0/0/0
3	PX4	C	1303	-	-	0/22/22/49	0/0/0/0
3	PX4	C	1304	-	-	0/3/3/49	0/0/0/0
3	PX4	C	1305	-	-	0/8/8/49	0/0/0/0
3	PX4	D	1302	-	-	0/8/8/49	0/0/0/0
3	PX4	D	1303	-	-	0/3/3/49	0/0/0/0
3	PX4	D	1304	-	-	0/8/8/49	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1305	PX4	O5-C9	2.80	1.41	1.33
3	B	1303	PX4	O5-C9	2.88	1.42	1.33
3	A	1304	PX4	O5-C9	2.92	1.42	1.33
3	C	1303	PX4	O5-C9	2.93	1.42	1.33
3	C	1303	PX4	O6-C9	5.40	1.38	1.22

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1303	PX4	O5-C9-O6	-5.47	109.37	123.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1304	PX4	O5-C9-O6	-5.47	109.38	123.49
3	C	1303	PX4	O5-C9-O6	-5.42	109.51	123.49
3	A	1305	PX4	O5-C9-O6	-5.34	109.72	123.49
3	C	1303	PX4	O6-C9-C10	-3.59	109.38	123.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1304	PX4	1	0
3	A	1305	PX4	2	0
3	B	1303	PX4	2	0
3	B	1305	PX4	1	0
3	C	1303	PX4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/237 (92%)	0.16	1 (0%) 91 90	41, 101, 152, 178	0
1	B	219/237 (92%)	0.19	3 (1%) 78 73	43, 101, 152, 172	0
1	C	219/237 (92%)	0.12	2 (0%) 85 82	38, 104, 155, 167	0
1	D	219/237 (92%)	0.13	1 (0%) 91 90	38, 101, 150, 176	0
All	All	876/948 (92%)	0.15	7 (0%) 87 84	38, 102, 153, 178	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1065	TYR	3.8
1	C	1065	TYR	3.0
1	A	1065	TYR	2.8
1	C	1001	MET	2.7
1	B	1069	ILE	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PX4	A	1306	6/46	0.91	0.38	3.18	54,59,63,64	0
3	PX4	B	1304	6/46	0.90	0.36	2.41	58,59,69,72	0
3	PX4	C	1304	6/46	0.89	0.32	1.98	57,65,74,74	0
2	CD	C	1301	1/1	0.99	0.23	1.69	112,112,112,112	0
2	CD	D	1301	1/1	0.99	0.22	1.28	107,107,107,107	0
2	CD	A	1301	1/1	0.99	0.30	1.22	52,52,52,52	0
3	PX4	D	1303	6/46	0.90	0.29	1.08	62,63,65,67	0
2	CD	B	1301	1/1	0.99	0.20	0.97	123,123,123,123	0
3	PX4	C	1303	21/46	0.88	0.27	0.66	66,87,102,106	0
3	PX4	A	1304	21/46	0.90	0.22	-0.44	69,83,100,108	0
3	PX4	A	1307	10/46	0.84	0.21	-0.51	70,94,110,113	0
3	PX4	B	1305	10/46	0.87	0.18	-0.64	75,92,111,140	0
3	PX4	B	1303	21/46	0.90	0.22	-0.67	66,82,103,114	0
3	PX4	C	1305	10/46	0.88	0.18	-0.74	79,110,133,141	0
3	PX4	A	1305	21/46	0.87	0.23	-0.75	71,85,106,109	0
3	PX4	D	1304	10/46	0.89	0.15	-1.18	90,98,112,133	0
2	CD	A	1302	1/1	0.97	0.18	-1.32	110,110,110,110	0
3	PX4	B	1302	10/46	0.83	0.17	-2.10	65,91,135,145	0
3	PX4	A	1303	10/46	0.87	0.17	-2.30	60,90,136,139	0
3	PX4	D	1302	10/46	0.81	0.16	-2.53	59,97,126,131	0
3	PX4	C	1302	10/46	0.75	0.18	-3.31	58,87,143,161	0

6.5 Other polymers [i](#)

There are no such residues in this entry.