Full wwPDB X-ray Structure Validation Report (i)

Jan 31, 2016 – 08:58 PM GMT

PDB ID	:	1MYT
Title	:	CRYSTAL STRUCTURE TO 1.74 ANGSTROMS RESOLUTION OF MET-MYOGLOBIN FROM YELLOWFIN TUNA (THUNNUS ALBACARES): AN EXAMPLE OF A MYOGLOBIN LACKING THE D HELIX
Authors	:	Birnbaum, G.I.; Evans, S.V.; Przybylska, M.; Rose, D.R.
Deposited on	:	1991-05-06
Resolution	:	1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2570 (1.76-1.72)
Ramachandran outliers	100387	2544 (1.76-1.72)
Sidechain outliers	100360	2544 (1.76-1.72)
RSRZ outliers	91569	2420 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HEM	A	154	-	-	-	X

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYOGLOBIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	146	1093	703	189	197	4	0	0	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	81	81	81	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MYOGLOBIN

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	44.50Å 72.32Å 52.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.74 6.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-1.74) 86.9 (6.00-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	7.65 (at 1.70Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.177 , (Not available) 0.178 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å^2)	18.1	Xtriage
Anisotropy	0.231	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.58 , 90.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	1 of 16202 reflections (0.006%)	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	1217	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.47% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.23	3/1113 (0.3%)	1.95	35/1500 (2.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	153	GLY	C-OXT	13.42	1.48	1.23
1	A	151	PHE	C-N	-9.53	1.12	1.34
1	A	153	GLY	CA-C	9.24	1.66	1.51

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	152	SER	C-N-CA	20.95	166.30	122.30
1	A	151	PHE	O-C-N	-15.45	97.98	122.70
1	A	152	SER	O-C-N	-13.21	100.73	123.20
1	A	46	PHE	CB-CG-CD2	-12.24	112.23	120.80
1	A	125	ALA	CB-CA-C	8.26	122.49	110.10
1	A	33	PHE	CB-CG-CD2	-8.24	115.03	120.80
1	A	29	LEU	CB-CG-CD1	8.19	124.92	111.00
1	A	46	PHE	CB-CG-CD1	7.58	126.10	120.80
1	A	20	ASP	O-C-N	7.37	134.50	122.70
1	A	74	GLU	CA-CB-CG	6.99	128.78	113.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	74	GLU	OE1-CD-OE2	6.96	131.66	123.30
1	A	149	LEU	CA-CB-CG	6.86	131.08	115.30
1	A	128	GLN	N-CA-CB	6.79	122.83	110.60
1	A	122	ASP	O-C-N	-6.75	111.91	122.70
1	A	111	LEU	CA-CB-CG	6.73	130.78	115.30
1	A	143	GLU	OE1-CD-OE2	6.62	131.25	123.30
1	A	146	TYR	CB-CG-CD1	6.53	124.92	121.00
1	A	74	GLU	CG-CD-OE2	-6.08	106.14	118.30
1	A	143	GLU	CG-CD-OE2	-5.82	106.67	118.30
1	A	109	GLU	OE1-CD-OE2	5.80	130.26	123.30
1	A	151	PHE	C-N-CA	5.80	136.21	121.70
1	A	11	LEU	O-C-N	5.72	131.85	122.70
1	A	22	THR	O-C-N	-5.68	113.61	122.70
1	A	152	SER	CA-C-N	5.64	127.48	116.20
1	A	121	LEU	O-C-N	5.61	131.67	122.70
1	A	122	ASP	CB-CG-OD1	5.54	123.28	118.30
1	A	146	TYR	O-C-N	5.54	131.56	122.70
1	A	148	GLU	CA-CB-CG	5.47	125.44	113.40
1	A	46	PHE	CZ-CE2-CD2	-5.45	113.56	120.10
1	A	106	LEU	O-C-N	5.43	131.39	122.70
1	A	23	THR	O-C-N	5.41	131.36	122.70
1	A	32	LEU	CB-CG-CD2	-5.14	102.25	111.00
1	A	75	LEU	CA-CB-CG	5.12	127.08	115.30
1	A	16	PRO	N-CA-CB	5.11	109.43	103.30
1	A	33	PHE	CB-CG-CD1	5.05	124.33	120.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	151	PHE	Mainchain
1	A	152	SER	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1093	0	1131	14	0
2	A	43	0	30	0	0
3	A	81	0	0	0	0
All	All	1217	0	1161	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:LYS:HE3	1:A:139:ILE:HD11	1.65	0.78
1:A:66:ALA:O	1:A:70:LYS:HG3	2.04	0.58
1:A:21:TYR:CD2	1:A:70:LYS:HG2	2.41	0.56
1:A:87:LYS:HE3	1:A:91:ASN:HD21	1.70	0.55
1:A:6:ASP:O	1:A:10:VAL:HG23	2.08	0.54
1:A:55:ILE:HG23	1:A:61:ILE:HD13	1.92	0.51
1:A:91:ASN:HD22	1:A:149:LEU:HD21	1.77	0.49
1:A:105:LYS:HE3	1:A:139:ILE:CD1	2.38	0.48
1:A:113:LYS:N	1:A:113:LYS:HD3	2.31	0.45
1:A:75:LEU:HD13	1:A:85:ILE:CG2	2.47	0.44
1:A:146:TYR:CD2	1:A:151:PHE:HD2	2.35	0.44
1:A:52:GLN:HA	1:A:55:ILE:HD12	2.02	0.42
1:A:24:MET:HE3	1:A:119:ALA:HB2	2.02	0.42
1:A:21:TYR:CE2	1:A:70:LYS:HG2	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	144/146 (99%)	139 (96%)	4 (3%)	1 (1%)	26 9

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	111/111 (100%)	100 (90%)	11 (10%)	10 1

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	29	LEU
1	A	31	ARG
1	A	71	LYS
1	A	75	LEU
1	A	109	GLU
1	A	111	LEU
1	A	113	LYS
1	A	139	ILE
1	A	148	GLU
1	A	149	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	GLN
1	A	91	ASN
1	A	133	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	HEM	A	154	1,3	30,50,50	2.65	11 (36%)	24,82,82	2.88	14 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	154	1,3	-	0/10/54/54	0/0/8/8

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	154	HEM	C2D-C3D	-7.02	1.33	1.54
2	A	154	HEM	C3B-C4B	-6.43	1.46	1.51
2	A	154	HEM	C3D-C4D	-5.01	1.45	1.51
2	A	154	HEM	C2C-C1C	-4.80	1.43	1.52
2	A	154	HEM	C2D-C1D	-2.29	1.44	1.51
2	A	154	HEM	FE-NC	2.29	2.04	1.95
2	A	154	HEM	CHD-C4C	2.45	1.42	1.36
2	A	154	HEM	CMA-C3A	2.49	1.56	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	154	HEM	C3C-CAC	2.60	1.56	1.51
2	A	154	HEM	C1C-NC	3.35	1.40	1.36
2	A	154	HEM	C4C-NC	3.51	1.40	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	154	HEM	C3C-CAC-CBC	-5.79	115.57	124.46
2	A	154	HEM	CAA-C2A-C1A	-3.50	123.21	127.01
2	A	154	HEM	CMA-C3A-C4A	-3.29	122.92	128.36
2	A	154	HEM	CBA-CAA-C2A	-2.02	108.91	112.53
2	A	154	HEM	C4B-CHC-C1C	2.08	129.30	125.82
2	A	154	HEM	C3B-CAB-CBB	2.09	127.66	124.46
2	A	154	HEM	CMA-C3A-C2A	2.19	129.81	125.24
2	A	154	HEM	CAA-CBA-CGA	2.26	116.88	112.75
2	A	154	HEM	CAD-C3D-C4D	2.53	121.40	112.47
2	A	154	HEM	CMD-C2D-C3D	3.09	128.01	114.35
2	A	154	HEM	C2D-C3D-C4D	3.54	107.50	101.50
2	A	154	HEM	CMB-C2B-C3B	4.57	127.93	116.53
2	A	154	HEM	CMC-C2C-C3C	4.92	128.81	116.53
2	A	154	HEM	CAD-C3D-C2D	6.08	130.68	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	146/146 (100%)	-0.21	4 (2%) 58 64	11, 22, 43, 66	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	GLY	10.6
1	A	152	SER	8.9
1	A	5	ALA	6.4
1	A	8	ASP	2.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
2	HEM	A	154	43/43	0.97	0.09	2.08	5,13,28,31	0

6.5 Other polymers (i)

There are no such residues in this entry.