



## wwPDB X-ray Structure Validation Summary Report

Feb 1, 2016 – 06:51 PM GMT

PDB ID : 4MYH  
Title : Structure of the Glutathione bound mitochondrial ABC transporter, Atm1  
Authors : Srinivasan, V.  
Deposited on : 2013-09-27  
Resolution : 3.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1084 (3.46-3.30)
Clashscore	102246	1158 (3.46-3.30)
Ramachandran outliers	100387	1139 (3.46-3.30)
Sidechain outliers	100360	1138 (3.46-3.30)
RSRZ outliers	91569	1089 (3.46-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	598	<div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 1%, green 63%, yellow 94%, orange 99%);"></div> <div style="margin-left: 10px;">%</div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>63%</span> <span>31%</span> <span>6%</span> </div>
1	B	598	<div style="display: flex; align-items: center;"> <div style="width: 10px; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 1%, green 65%, yellow 93%, orange 99%, red 100%);"></div> <div style="margin-left: 10px;">%</div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>65%</span> <span>28%</span> <span>6%</span> <span>•</span> </div>
1	C	598	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, green 73%, yellow 95%, orange 100%);"></div> </div> <div style="display: flex; justify-content: space-around; margin-top: 5px;"> <span>73%</span> <span>22%</span> <span>5%</span> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GSH	B	701	-	-	-	X

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 14048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Iron-sulfur clusters transporter ATM1, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	598	4676	3002	802	855	17	0	0	0
1	C	598	4676	3002	802	855	17	0	0	0
1	B	598	4676	3002	802	855	17	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	691	TRP	-	EXPRESSION TAG	UNP P40416
A	692	SER	-	EXPRESSION TAG	UNP P40416
A	693	HIS	-	EXPRESSION TAG	UNP P40416
A	694	PRO	-	EXPRESSION TAG	UNP P40416
A	695	GLN	-	EXPRESSION TAG	UNP P40416
C	691	TRP	-	EXPRESSION TAG	UNP P40416
C	692	SER	-	EXPRESSION TAG	UNP P40416
C	693	HIS	-	EXPRESSION TAG	UNP P40416
C	694	PRO	-	EXPRESSION TAG	UNP P40416
C	695	GLN	-	EXPRESSION TAG	UNP P40416
B	691	TRP	-	EXPRESSION TAG	UNP P40416
B	692	SER	-	EXPRESSION TAG	UNP P40416
B	693	HIS	-	EXPRESSION TAG	UNP P40416
B	694	PRO	-	EXPRESSION TAG	UNP P40416
B	695	GLN	-	EXPRESSION TAG	UNP P40416

- Molecule 2 is GLUTATHIONE (three-letter code: GSH) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	B	1	20	10	3	6	1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Iron-sulfur clusters transporter ATM1, mitochondrial




- Molecule 1: Iron-sulfur clusters transporter ATM1, mitochondrial





● Molecule 1: Iron-sulfur clusters transporter ATM1, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.69Å 157.69Å 523.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	58.99 – 3.38 58.99 – 3.38	Depositor EDS
% Data completeness (in resolution range)	99.7 (58.99-3.38) 99.6 (58.99-3.38)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 3.40Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.247 , 0.290 0.256 , 0.293	Depositor DCC
$R_{free}$ test set	2782 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	71.1	Xtrriage
Anisotropy	0.222	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 35.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Outliers	4 of 54830 reflections (0.007%)	Xtrriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	14048	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/4763	0.66	0/6451
1	B	0.56	0/4763	0.65	1/6451 (0.0%)
1	C	0.53	0/4763	0.64	0/6451
All	All	0.56	0/14289	0.65	1/19353 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	428	LEU	C-N-CD	-6.73	105.80	120.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	428	LEU	Peptide
1	B	428	LEU	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4676	0	4738	141	0
1	B	4676	0	4738	145	0
1	C	4676	0	4738	74	0
2	B	20	0	15	3	0
All	All	14048	0	14229	326	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 326 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:261:ILE:HD11	1:B:357:MET:HG3	1.59	0.84
1:A:531:LYS:NZ	1:A:535:ILE:O	2.12	0.82
1:A:443:PHE:HZ	1:A:486:TYR:HD1	1.31	0.78
1:A:472:GLY:HA3	1:B:694:PRO:HG3	1.64	0.77
1:A:280:ARG:NH1	1:A:338:SER:OG	2.18	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	596/598 (100%)	541 (91%)	34 (6%)	21 (4%)	4	32
1	B	596/598 (100%)	540 (91%)	38 (6%)	18 (3%)	5	36

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	596/598 (100%)	542 (91%)	40 (7%)	14 (2%)	8	42
All	All	1788/1794 (100%)	1623 (91%)	112 (6%)	53 (3%)	5	36

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	312	ASN
1	A	392	LEU
1	A	395	VAL
1	A	621	SER
1	C	312	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	501/514 (98%)	446 (89%)	55 (11%)	8	30
1	B	501/514 (98%)	441 (88%)	60 (12%)	6	26
1	C	501/514 (98%)	442 (88%)	59 (12%)	6	26
All	All	1503/1542 (98%)	1329 (88%)	174 (12%)	7	28

5 of 174 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	391	PHE
1	C	578	LYS
1	B	623	SER
1	C	399	LEU
1	C	456	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	647	ASN
1	B	695	GLN
1	B	384	GLN
1	A	380	GLN
1	B	379	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GSH	B	701	-	13,19,19	2.46	3 (23%)	15,24,24	1.94	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GSH	B	701	-	-	0/18/24/24	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	GSH	CB2-SG2	-2.31	1.76	1.81
2	B	701	GSH	C2-N3	5.77	1.45	1.33
2	B	701	GSH	CD1-N2	5.94	1.46	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	GSH	CA2-CB2-SG2	-5.14	107.83	114.16
2	B	701	GSH	CB1-CG1-CD1	-2.59	107.12	113.27
2	B	701	GSH	C2-CA2-N2	2.07	117.09	111.26
2	B	701	GSH	CB2-CA2-C2	2.52	115.34	109.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	701	GSH	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	598/598 (100%)	0.09	8 (1%) 79 81	12, 41, 59, 82	0
1	B	598/598 (100%)	0.01	7 (1%) 81 82	9, 37, 53, 70	0
1	C	598/598 (100%)	-0.02	2 (0%) 94 95	8, 28, 46, 63	0
All	All	1794/1794 (100%)	0.03	17 (0%) 85 88	8, 35, 55, 82	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	ASN	5.6
1	B	282	HIS	3.8
1	A	401	GLN	3.4
1	C	392	LEU	2.9
1	B	692	SER	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GSH	B	701	20/20	0.81	0.42	2.11	61,66,71,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.