



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:52 PM GMT

PDB ID : 4MZ7
Title : Structural insight into dGTP-dependent activation of tetrameric SAMHD1 deoxynucleoside triphosphate triphosphohydrolase
Authors : Zhu, C.; Gao, W.; Zhao, K.; Qin, X.; Zhang, Y.; Peng, X.; Zhang, L.; Dong, Y.; Zhang, W.; Li, P.; Wei, W.; Gong, Y.; Yu, X.F.
Deposited on : 2013-09-29
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	539	<div> <div>7%</div> <div>82% 7% 11%</div> </div>
1	B	539	<div> <div>3%</div> <div>86% 5% 9%</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxynucleoside triphosphate triphosphohydrolase SAMHD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	481	Total	C	N	O	S	0	2	0
			3955	2534	688	714	19			
1	B	493	Total	C	N	O	S	0	2	0
			4050	2596	704	731	19			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	88	MET	-	EXPRESSION TAG	UNP Q9Y3Z3
A	89	GLY	-	EXPRESSION TAG	UNP Q9Y3Z3
A	90	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
A	91	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
A	92	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
A	93	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
A	94	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
A	95	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
A	96	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
A	97	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
A	98	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
A	99	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
A	100	GLY	-	EXPRESSION TAG	UNP Q9Y3Z3
A	101	GLU	-	EXPRESSION TAG	UNP Q9Y3Z3
A	102	ASN	-	EXPRESSION TAG	UNP Q9Y3Z3
A	103	LEU	-	EXPRESSION TAG	UNP Q9Y3Z3
A	104	TYR	-	EXPRESSION TAG	UNP Q9Y3Z3
A	105	PHE	-	EXPRESSION TAG	UNP Q9Y3Z3
A	106	GLN	-	EXPRESSION TAG	UNP Q9Y3Z3
A	107	GLY	-	EXPRESSION TAG	UNP Q9Y3Z3
A	108	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
A	266	TYR	CYS	ENGINEERED MUTATION	UNP Q9Y3Z3
B	88	MET	-	EXPRESSION TAG	UNP Q9Y3Z3
B	89	GLY	-	EXPRESSION TAG	UNP Q9Y3Z3
B	90	SER	-	EXPRESSION TAG	UNP Q9Y3Z3

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	91	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
B	92	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
B	93	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
B	94	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
B	95	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
B	96	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
B	97	HIS	-	EXPRESSION TAG	UNP Q9Y3Z3
B	98	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
B	99	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
B	100	GLY	-	EXPRESSION TAG	UNP Q9Y3Z3
B	101	GLU	-	EXPRESSION TAG	UNP Q9Y3Z3
B	102	ASN	-	EXPRESSION TAG	UNP Q9Y3Z3
B	103	LEU	-	EXPRESSION TAG	UNP Q9Y3Z3
B	104	TYR	-	EXPRESSION TAG	UNP Q9Y3Z3
B	105	PHE	-	EXPRESSION TAG	UNP Q9Y3Z3
B	106	GLN	-	EXPRESSION TAG	UNP Q9Y3Z3
B	107	GLY	-	EXPRESSION TAG	UNP Q9Y3Z3
B	108	SER	-	EXPRESSION TAG	UNP Q9Y3Z3
B	266	TYR	CYS	ENGINEERED MUTATION	UNP Q9Y3Z3

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (three-letter code: DGT) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
3	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
3	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
3	B	1	Total 31	C 10	N 5	O 13	P 3	0	0
3	B	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: $C_{10}H_{16}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			30	10	5	12	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	347	Total	O	0	0
			347	347		
6	B	472	Total	O	0	0
			472	472		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxynucleoside triphosphate triphosphohydrolase SAMHD1



- Molecule 1: Deoxynucleoside triphosphate triphosphohydrolase SAMHD1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	147.40 Å 103.61 Å 92.13 Å 90.00° 121.98° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 48.27 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.2 (50.00-1.80) 97.2 (48.27-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 1.79 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.194 , 0.226 0.194 , 0.226	Depositor DCC
R_{free} test set	5284 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 105546 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9013	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, DTP, MG, DGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	1/4054 (0.0%)	0.56	0/5474
1	B	0.52	1/4155 (0.0%)	0.58	0/5611
All	All	0.53	2/8209 (0.0%)	0.57	0/11085

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	231	TRP	CD2-CE2	5.13	1.47	1.41
1	B	231	TRP	CD2-CE2	5.07	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3955	0	3928	21	0
1	B	4050	0	4020	24	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	93	0	36	2	0
3	B	62	0	24	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
5	B	30	0	12	0	0
6	A	347	0	0	6	0
6	B	472	0	0	6	0
All	All	9013	0	8020	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:355:GLU:HG3	6:A:1062:HOH:O	1.82	0.80
1:A:591:ILE:O	1:A:594:GLN:HG2	1.83	0.78
1:B:243:HIS:HE1	6:B:1370:HOH:O	1.66	0.77
1:B:179:VAL:HG22	1:B:300:ILE:HD13	1.70	0.72
1:A:158:PRO:HG3	6:A:1059:HOH:O	1.94	0.67
1:B:397:ILE:HG21	1:B:426:ILE:HD11	1.77	0.64
1:A:326:GLN:NE2	6:A:1093:HOH:O	2.32	0.61
1:A:485:PRO:HG2	1:A:489:LEU:HD11	1.83	0.59
1:B:397:ILE:HB	6:B:1054:HOH:O	2.03	0.58
1:B:175:ALA:HB1	1:B:199:VAL:HG12	1.86	0.57
1:A:115:MET:HE1	1:A:127:GLU:HB3	1.87	0.57
1:A:395:ASP:OD1	1:A:408:ARG:NH2	2.39	0.56
1:B:179:VAL:CG2	1:B:300:ILE:HD13	2.35	0.56
1:A:494:LYS:HG3	1:A:496:GLU:HG2	1.90	0.53
3:A:703:DGT:N2	6:A:859:HOH:O	2.34	0.52
1:A:397:ILE:HG21	1:A:426:ILE:HD11	1.90	0.52
1:A:143:ARG:HD2	1:A:420:THR:HA	1.91	0.51
1:A:446:LYS:HE2	6:A:1147:HOH:O	2.09	0.51
1:A:439:LYS:O	1:A:443:GLU:HG2	2.10	0.51
1:B:297:LEU:HA	1:B:300:ILE:HD12	1.93	0.49
3:A:703:DGT:HN2A	1:B:119:ASN:ND2	2.11	0.49
1:B:509:MET:HE1	1:B:517:HIS:CD2	2.48	0.49
1:A:175:ALA:HB1	1:A:199:VAL:HG12	1.95	0.48
1:B:302:SER:O	6:B:905:HOH:O	2.20	0.48
1:A:302:SER:O	6:A:805:HOH:O	2.21	0.46
1:B:478:LYS:HD3	6:B:980:HOH:O	2.16	0.45
1:B:595:LYS:HE3	1:B:597:GLU:HG2	1.97	0.45
1:A:381:ILE:HA	1:A:381:ILE:HD12	1.84	0.45
1:A:390:PHE:CZ	1:A:426:ILE:HG23	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:ALA:O	1:B:520:PHE:HA	2.17	0.45
1:B:577:ASN:HD21	1:B:597:GLU:HG3	1.81	0.45
1:B:143:ARG:HD2	1:B:420:THR:HA	1.98	0.44
1:B:179:VAL:HG22	1:B:300:ILE:CD1	2.45	0.44
1:A:351:ALA:O	1:A:520:PHE:HA	2.17	0.44
1:B:215[A]:HIS:CD2	6:B:1252:HOH:O	2.70	0.44
1:B:179:VAL:HG21	1:B:199:VAL:HG11	2.00	0.43
1:B:580:LYS:HE3	1:B:592:THR:HG21	2.00	0.43
1:B:381:ILE:HD12	1:B:381:ILE:HA	1.91	0.43
1:B:390:PHE:CZ	1:B:426:ILE:HG23	2.54	0.42
1:A:331:TYR:CD1	1:A:331:TYR:C	2.92	0.42
1:B:509:MET:HB2	1:B:509:MET:HE2	1.79	0.42
1:A:291:PRO:HG2	1:A:293:ASN:OD1	2.20	0.41
1:B:179:VAL:CG2	1:B:199:VAL:HG11	2.50	0.41
1:B:417:GLU:OE2	6:B:1272:HOH:O	2.22	0.40
1:A:334:PHE:CE2	1:A:359:LEU:HD21	2.56	0.40
1:A:509:MET:HE2	1:A:509:MET:HB2	1.80	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/539 (89%)	471 (98%)	8 (2%)	0	100	100
1	B	491/539 (91%)	483 (98%)	8 (2%)	0	100	100
All	All	970/1078 (90%)	954 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	429/481 (89%)	423 (99%)	6 (1%)	74	65
1	B	440/481 (92%)	436 (99%)	4 (1%)	84	80
All	All	869/962 (90%)	859 (99%)	10 (1%)	78	71

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	MET
1	A	315	TYR
1	A	331	TYR
1	A	425	ASN
1	A	528	ARG
1	A	543	GLU
1	B	112	VAL
1	B	315	TYR
1	B	528	ARG
1	B	577	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	326	GLN
1	A	328	ASN
1	B	119	ASN
1	B	243	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	DGT	A	702	-	25,33,33	1.05	2 (8%)	35,52,52	1.91	6 (17%)
3	DGT	A	703	4	25,33,33	1.09	2 (8%)	35,52,52	1.76	6 (17%)
3	DGT	A	704	4	25,33,33	1.12	2 (8%)	35,52,52	1.80	6 (17%)
3	DGT	B	801	4	25,33,33	1.15	2 (8%)	35,52,52	1.83	6 (17%)
3	DGT	B	804	-	25,33,33	1.03	2 (8%)	35,52,52	1.78	7 (20%)
5	DTP	B	805	4	24,32,32	0.96	0	32,50,50	2.22	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DGT	A	702	-	-	0/18/34/34	0/3/3/3
3	DGT	A	703	4	-	0/18/34/34	0/3/3/3
3	DGT	A	704	4	-	0/18/34/34	0/3/3/3
3	DGT	B	801	4	-	0/18/34/34	0/3/3/3
3	DGT	B	804	-	-	0/18/34/34	0/3/3/3
5	DTP	B	805	4	-	0/18/34/34	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	804	DGT	C5-C4	2.22	1.45	1.40
3	A	703	DGT	C5-C4	2.26	1.45	1.40
3	A	704	DGT	C5-C4	2.36	1.45	1.40
3	B	801	DGT	C5-C4	2.47	1.46	1.40
3	B	801	DGT	C6-C5	2.73	1.46	1.41
3	A	702	DGT	C5-C4	2.87	1.47	1.40
3	A	702	DGT	C6-C5	2.93	1.47	1.41
3	B	804	DGT	C6-C5	2.98	1.47	1.41
3	A	704	DGT	C6-C5	2.98	1.47	1.41
3	A	703	DGT	C6-C5	3.09	1.47	1.41

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	805	DTP	N3-C2-N1	-9.72	121.45	128.89
3	B	801	DGT	C5-C6-N1	-4.92	116.87	123.59
3	B	804	DGT	PA-O3A-PB	-4.82	119.20	132.73
3	A	704	DGT	C5-C6-N1	-4.79	117.05	123.59
3	A	702	DGT	PA-O3A-PB	-4.74	119.42	132.73
3	A	703	DGT	C5-C6-N1	-4.45	117.50	123.59
5	B	805	DTP	C1'-N9-C4	-4.38	119.74	127.16
3	A	704	DGT	N3-C2-N1	-4.15	121.12	127.44
3	A	702	DGT	N3-C2-N1	-4.11	121.18	127.44
3	B	801	DGT	N3-C2-N1	-3.91	121.49	127.44
3	A	702	DGT	C5-C6-N1	-3.85	118.32	123.59
3	B	804	DGT	N3-C2-N1	-3.81	121.63	127.44
3	A	703	DGT	N3-C2-N1	-3.70	121.81	127.44
3	B	804	DGT	C5-C6-N1	-3.61	118.65	123.59
3	A	703	DGT	C1'-N9-C4	-3.60	121.06	127.16
3	B	801	DGT	PA-O3A-PB	-3.36	123.30	132.73
3	A	702	DGT	C6-C5-C4	-3.18	117.10	120.90
3	A	703	DGT	C6-C5-C4	-2.90	117.43	120.90
3	A	702	DGT	PB-O3B-PG	-2.82	123.20	132.67
3	A	704	DGT	PA-O3A-PB	-2.75	125.01	132.73
3	B	804	DGT	C6-C5-C4	-2.69	117.69	120.90
3	A	703	DGT	C4-C5-N7	-2.61	107.08	109.48
3	B	801	DGT	C6-C5-C4	-2.54	117.86	120.90
3	A	704	DGT	C6-C5-C4	-2.53	117.88	120.90
5	B	805	DTP	C4-C5-N7	-2.52	107.16	109.48
3	B	804	DGT	C1'-N9-C4	-2.50	122.92	127.16
3	B	804	DGT	C4-C5-N7	-2.38	107.29	109.48
5	B	805	DTP	PB-O3B-PG	-2.37	124.71	132.67
3	B	801	DGT	C4-C5-N7	-2.20	107.46	109.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	704	DGT	C4-C5-N7	-2.16	107.49	109.48
5	B	805	DTP	O2B-PB-O3B	2.16	114.89	105.09
5	B	805	DTP	N6-C6-N1	2.39	124.33	119.20
5	B	805	DTP	C2-N1-C6	2.44	123.12	118.77
3	B	804	DGT	C6-N1-C2	4.17	121.73	115.94
3	A	703	DGT	C6-N1-C2	4.89	122.73	115.94
3	A	702	DGT	C6-N1-C2	4.92	122.76	115.94
3	B	801	DGT	C6-N1-C2	5.42	123.46	115.94
3	A	704	DGT	C6-N1-C2	5.58	123.68	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	703	DGT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	481/539 (89%)	0.39	38 (7%)	15 12	6, 22, 52, 77	0
1	B	493/539 (91%)	0.07	17 (3%)	49 43	4, 17, 40, 81	0
All	All	974/1078 (90%)	0.23	55 (5%)	28 22	4, 19, 46, 81	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	111	HIS	7.9
1	A	488	LEU	7.5
1	A	491	VAL	6.2
1	A	489	LEU	5.7
1	A	593	PRO	5.1
1	A	277	GLU	5.0
1	A	594	GLN	5.0
1	A	596	LYS	5.0
1	A	284	LEU	5.0
1	A	590	LEU	4.9
1	B	284	LEU	4.7
1	A	487	VAL	4.6
1	A	599	ASN	4.3
1	A	403	GLY	4.3
1	A	486	LYS	4.1
1	A	490	ASP	4.1
1	B	593	PRO	3.9
1	A	591	ILE	3.9
1	A	465	GLN	3.8
1	B	277	GLU	3.8
1	B	110	ILE	3.7
1	A	285	TRP	3.7
1	B	403	GLY	3.6
1	A	577	ASN	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	535	ASN	3.5
1	A	464	GLY	3.5
1	A	592	THR	3.3
1	B	596	LYS	3.3
1	A	463	THR	3.2
1	A	597	GLU	3.2
1	A	559	ARG	3.1
1	A	598	TRP	3.1
1	A	471	GLU	3.1
1	A	345	ASN	3.0
1	A	466	ILE	3.0
1	A	230	LYS	2.9
1	B	278	SER	2.9
1	B	112	VAL	2.9
1	A	344	ASP	2.8
1	B	328	ASN	2.7
1	B	466	ILE	2.6
1	B	113	ASP	2.6
1	B	230	LYS	2.6
1	A	114	THR	2.6
1	B	599	ASN	2.5
1	A	563	TYR	2.4
1	A	113	ASP	2.4
1	B	102	ASN	2.4
1	B	464	GLY	2.4
1	A	328	ASN	2.3
1	A	574	ALA	2.3
1	B	325	ILE	2.2
1	A	470	ARG	2.1
1	A	326	GLN	2.1
1	A	584	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	DGT	A	702	31/31	0.83	0.15	1.47	10,17,43,47	0
3	DGT	B	804	31/31	0.86	0.15	1.39	7,13,38,39	0
5	DTP	B	805	30/30	0.98	0.12	-0.52	6,8,11,12	0
3	DGT	B	801	31/31	0.97	0.09	-0.54	9,11,20,22	0
2	ZN	A	701	1/1	0.99	0.10	-0.56	14,14,14,14	0
3	DGT	A	704	31/31	0.98	0.09	-0.82	8,9,15,16	0
2	ZN	B	803	1/1	0.99	0.08	-0.88	11,11,11,11	0
3	DGT	A	703	31/31	0.98	0.10	-1.09	8,11,15,17	0
4	MG	B	802	1/1	0.95	0.12	-	19,19,19,19	0
4	MG	A	705	1/1	0.95	0.14	-	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.