



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:52 PM GMT

PDB ID : 4N1G
Title : Crystal Structure of Ca(2+)- discharged F88Y obelin mutant from Obelia longissima at 1.50 Angstrom resolution
Authors : Natashin, P.V.; Markova, S.V.; Lee, J.; Vysotski, E.S.; Liu, Z.J.
Deposited on : 2013-10-04
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 88%, yellow 88%, yellow 99%, green 99%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 88% 11% .. </div> </div>
1	B	195	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 7%, orange 7%, orange 87%, yellow 87%, yellow 99%, green 99%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 7% 87% 12% .. </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Obelin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	3	0
			1575	997	264	305	9			
1	B	194	Total	C	N	O	S	0	3	0
			1578	996	267	306	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	SER	ENGINEERED MUTATION	UNP Q27709
A	88	TYR	PHE	ENGINEERED MUTATION	UNP Q27709
B	2	ALA	SER	ENGINEERED MUTATION	UNP Q27709
B	88	TYR	PHE	ENGINEERED MUTATION	UNP Q27709

- Molecule 2 is N-[3-BENZYL-5-(4-HYDROXYPHENYL)PYRAZIN-2-YL]-2-(4-HYDROXY PHENYL)ACETAMIDE (three-letter code: CEI) (formula: C₂₅H₂₁N₃O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			31	25	3	3		
2	B	1	Total	C	N	O	0	0
			31	25	3	3		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Ca	0	0
			3	3		
3	A	3	Total	Ca	0	0
			3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	308	Total	O	0	0
			308	308		
4	B	258	Total	O	0	0
			258	258		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


● Molecule 1: Obelin

Chain A: 



● Molecule 1: Obelin

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.92Å 85.13Å 90.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.54 – 1.50 49.54 – 1.50	Depositor EDS
% Data completeness (in resolution range)	77.8 (49.54-1.50) 74.7 (49.54-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.169 , 0.209 0.172 , 0.207	Depositor DCC
R_{free} test set	2001 reflections (3.66%)	DCC
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.332	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 58839 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3787	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CEI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	2/1620 (0.1%)	0.88	2/2187 (0.1%)
1	B	0.94	2/1620 (0.1%)	0.89	1/2187 (0.0%)
All	All	0.97	4/3240 (0.1%)	0.88	3/4374 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	75	CYS	CB-SG	-5.95	1.72	1.81
1	B	75	CYS	CB-SG	-5.79	1.72	1.81
1	B	59	GLU	CD-OE2	5.44	1.31	1.25
1	A	185	GLU	CD-OE1	5.23	1.31	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	48	ASP	CB-CG-OD1	5.27	123.04	118.30
1	B	21	ARG	CG-CD-NE	-5.21	100.86	111.80
1	A	165	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1575	0	1499	14	0
1	B	1578	0	1497	20	0
2	A	31	0	20	1	0
2	B	31	0	21	5	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	308	0	0	4	0
4	B	258	0	0	7	0
All	All	3787	0	3037	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:LYS:HE3	1:B:83:ILE:HD11	1.58	0.85
1:B:28:PHE:HE2	2:B:201:CEI:H29	1.43	0.83
1:B:28:PHE:CE2	2:B:201:CEI:H29	2.20	0.76
1:A:81:LYS:HE3	1:A:83:ILE:HD11	1.78	0.65
1:A:12:ASP:OD1	1:A:12:ASP:N	2.32	0.62
1:A:11:THR:HG21	1:A:181:THR:HA	1.82	0.61
1:B:21:ARG:HG2	1:B:195:PRO:O	2.00	0.61
1:B:115:GLY:HA3	1:B:172[B]:THR:HG22	1.83	0.61
1:B:173[A]:ARG:NH2	1:B:183:ASP:OD2	2.34	0.60
1:B:103:TRP:HZ3	4:B:503:HOH:O	1.85	0.59
1:B:93:LYS:HE3	4:B:555:HOH:O	2.03	0.57
1:B:8:LYS:NZ	4:B:548:HOH:O	2.30	0.57
1:A:62:LYS:NZ	4:A:598:HOH:O	2.39	0.55
1:B:111:ILE:HG12	4:B:504:HOH:O	2.05	0.55
1:B:142:SER:HB2	2:B:201:CEI:C30	2.39	0.52
1:B:172[A]:THR:HG23	4:B:504:HOH:O	2.10	0.50
1:A:11:THR:O	4:A:476:HOH:O	2.20	0.49
1:A:119:PHE:CE2	1:A:130[B]:ILE:HG12	2.47	0.49
1:A:62:LYS:HE3	1:B:103:TRP:CZ2	2.48	0.49
1:B:112:ARG:NH2	1:B:169:ASP:OD1	2.43	0.48
1:B:45:LYS:HE3	1:B:45:LYS:HB3	1.75	0.47
1:A:105:ARG:NH2	4:A:442:HOH:O	2.48	0.46
1:A:102:LYS:NZ	4:A:441:HOH:O	2.48	0.45
1:B:11:THR:HB	4:B:441:HOH:O	2.16	0.44
1:A:119:PHE:HD2	1:A:171[A]:MET:SD	2.41	0.44
1:A:140:LYS:HE2	1:A:148:GLN:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:PHE:O	1:A:18:TRP:CD1	2.72	0.43
1:B:17:ARG:HG2	1:B:17:ARG:HH21	1.83	0.42
1:A:179:TRP:CD2	2:A:201:CEI:H24	2.54	0.42
1:A:75:CYS:SG	1:A:95:LEU:HD22	2.60	0.42
1:B:103:TRP:HH2	4:B:387:HOH:O	2.02	0.42
1:B:179:TRP:CE2	2:B:201:CEI:H20	2.56	0.41
1:B:179:TRP:CD2	2:B:201:CEI:H20	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/195 (100%)	192 (98%)	3 (2%)	0	100	100
1	B	195/195 (100%)	192 (98%)	3 (2%)	0	100	100
All	All	390/390 (100%)	384 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/166 (101%)	165 (98%)	3 (2%)	66	35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	168/166 (101%)	165 (98%)	3 (2%)	66	35
All	All	336/332 (101%)	330 (98%)	6 (2%)	66	35

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	THR
1	A	53	LYS
1	A	116	ASP
1	B	10	LYS
1	B	89	LEU
1	B	136	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CEI	A	201	-	34,34,34	1.68	6 (17%)	41,46,46	1.75	10 (24%)
2	CEI	B	201	-	34,34,34	1.60	7 (20%)	41,46,46	2.15	13 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CEI	A	201	-	-	0/16/16/16	0/4/4/4
2	CEI	B	201	-	-	0/16/16/16	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	CEI	C19-C6	-5.23	1.40	1.48
2	B	201	CEI	C19-C6	-4.47	1.42	1.48
2	B	201	CEI	C26-C8	-2.52	1.49	1.51
2	A	201	CEI	C9-C8	-2.20	1.38	1.42
2	B	201	CEI	O25-C22	2.18	1.42	1.37
2	B	201	CEI	C6-N7	2.26	1.38	1.34
2	A	201	CEI	C21-C20	2.28	1.42	1.38
2	B	201	CEI	C13-C12	2.40	1.43	1.38
2	B	201	CEI	C5-N4	2.54	1.39	1.34
2	A	201	CEI	C9-N4	2.57	1.39	1.35
2	B	201	CEI	C12-C11	2.62	1.44	1.38
2	A	201	CEI	C2-N1	2.72	1.42	1.35
2	A	201	CEI	C9-N1	2.78	1.45	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	CEI	C27-C26-C8	-4.63	103.25	112.58
2	B	201	CEI	C12-C13-C14	-4.22	114.99	119.87
2	A	201	CEI	C6-C5-N4	-3.89	116.82	122.43
2	B	201	CEI	C6-C5-N4	-3.58	117.25	122.43
2	A	201	CEI	C11-C10-C2	-3.17	102.49	112.84
2	B	201	CEI	C15-C16-C11	-3.10	116.80	121.04
2	A	201	CEI	C9-N1-C2	-3.06	123.00	128.12
2	B	201	CEI	C9-N1-C2	-3.00	123.11	128.12
2	A	201	CEI	O33-C2-N1	-2.93	118.48	123.72
2	B	201	CEI	C11-C10-C2	-2.74	103.91	112.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	201	CEI	O33-C2-N1	-2.52	119.22	123.72
2	A	201	CEI	C16-C11-C12	-2.48	114.16	118.13
2	A	201	CEI	C27-C26-C8	-2.41	107.73	112.58
2	B	201	CEI	C24-C23-C22	2.17	122.38	119.87
2	A	201	CEI	C5-N4-C9	2.55	121.34	116.30
2	B	201	CEI	C13-C12-C11	3.01	125.16	121.04
2	A	201	CEI	C15-C16-C11	3.06	125.23	121.04
2	B	201	CEI	C5-N4-C9	3.09	122.41	116.30
2	B	201	CEI	C32-C27-C28	3.23	123.31	118.13
2	A	201	CEI	C10-C11-C12	3.30	125.81	120.86
2	A	201	CEI	C6-N7-C8	3.84	121.14	118.11
2	B	201	CEI	C16-C15-C14	4.65	125.24	119.87
2	B	201	CEI	C6-N7-C8	5.02	122.07	118.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	CEI	1	0
2	B	201	CEI	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/195 (99%)	-0.25	2 (1%) 84 86	9, 16, 32, 47	0
1	B	194/195 (99%)	-0.02	14 (7%) 18 18	10, 19, 45, 63	0
All	All	388/390 (99%)	-0.13	16 (4%) 41 43	9, 17, 42, 63	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	ILE	4.7
1	B	137	ALA	4.6
1	B	192	ASN	4.3
1	B	193	GLY	3.9
1	B	194	VAL	3.6
1	B	195	PRO	3.3
1	B	138	TYR	3.2
1	B	147	SER	2.9
1	A	11	THR	2.9
1	B	146	PRO	2.8
1	B	141	ILE	2.5
1	B	142	SER	2.5
1	B	191	GLY	2.4
1	A	14	ASP	2.2
1	B	11	THR	2.1
1	B	145	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CA	A	204	1/1	1.00	0.09	1.33	12,12,12,12	0
3	CA	B	202	1/1	1.00	0.06	0.07	14,14,14,14	0
3	CA	A	202	1/1	1.00	0.07	-0.03	16,16,16,16	0
2	CEI	B	201	31/31	0.97	0.08	-0.07	10,13,42,46	0
3	CA	A	203	1/1	1.00	0.07	-0.23	16,16,16,16	0
2	CEI	A	201	31/31	0.97	0.07	-0.35	9,12,20,22	0
3	CA	B	203	1/1	0.99	0.06	-0.58	19,19,19,19	0
3	CA	B	204	1/1	0.99	0.05	-2.12	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.