



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 06:54 PM GMT

PDB ID : 4N4T

Title : Co-crystal structure of tankyrase 1 with compound 3 [(4S)-3-{4-[6-amino-5-(pyrimidin-2-yl)pyridin-3-yl]phenyl}-5,5-dimethyl-4-phenyl-1,3-oxazolidin-2-one]

Authors : Huang, X.

Deposited on : 2013-10-08

Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.7 (RC4), CSD as536be (2015)

Xtriage (Phenix) : 1.9-1692

EDS : rb-20026688

Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)

Refmac : 5.8.0135

CCP4 : 6.5.0

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	2GV	A	1402	-	-	-	X
3	2GV	B	1402	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tankyrase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C 1694	N 1066	O 309	S 308	11	0	0
1	B	199	Total	C 1598	N 1006	O 292	S 290	10	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1315	HIS	-	EXPRESSION TAG	UNP Q6PFX9
A	1316	HIS	-	EXPRESSION TAG	UNP Q6PFX9
A	1317	HIS	-	EXPRESSION TAG	UNP Q6PFX9
A	1318	HIS	-	EXPRESSION TAG	UNP Q6PFX9
A	1319	HIS	-	EXPRESSION TAG	UNP Q6PFX9
A	1320	HIS	-	EXPRESSION TAG	UNP Q6PFX9
B	1315	HIS	-	EXPRESSION TAG	UNP Q6PFX9
B	1316	HIS	-	EXPRESSION TAG	UNP Q6PFX9
B	1317	HIS	-	EXPRESSION TAG	UNP Q6PFX9
B	1318	HIS	-	EXPRESSION TAG	UNP Q6PFX9
B	1319	HIS	-	EXPRESSION TAG	UNP Q6PFX9
B	1320	HIS	-	EXPRESSION TAG	UNP Q6PFX9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is (4S)-3-{4-[6-AMINO-5-(PYRIMIDIN-2-YL)PYRIDIN-3-YL]PHENYL}-5,5-DIMETHYL-4-PHENYL-1,3-OXAZOLIDIN-2-ONE (three-letter code: 2GV) (formula: C₂₆H₂₃N₅O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 33 26 5 2	0	0
3	B	1	Total C N O 33 26 5 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	74	Total O 74 74	0	0
4	B	67	Total O 67 67	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tankyrase-1



- Molecule 1: Tankyrase-1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.94Å 77.88Å 147.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.31 41.60 – 2.31	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.31) 99.2 (41.60-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.40 (at 2.32Å)	Xtriage
Refinement program	CNS	Depositor
R , R_{free}	0.251 , 0.275 0.260 , 0.280	Depositor DCC
R_{free} test set	1111 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.342	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.51$, $< L^2 > = 0.35$	Xtriage
Outliers	0 of 21689 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3501	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.03 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.5043e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 2GV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.45	0 / 1738	0.70	1 / 2340 (0.0%)
1	B	0.45	1 / 1637 (0.1%)	0.68	0 / 2200
All	All	0.45	1 / 3375 (0.0%)	0.69	1 / 4540 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1148	ILE	CB-CG2	5.45	1.69	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1281	ARG	NE-CZ-NH1	-5.21	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1694	0	1608	45	0
1	B	1598	0	1504	57	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	33	0	23	3	0
3	B	33	0	23	2	0
4	A	74	0	0	3	0
4	B	67	0	0	6	0
All	All	3501	0	3158	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1122:GLU:HG3	1:B:1147:VAL:HG21	1.53	0.89
1:A:1204:ILE:HD13	1:A:1209:GLY:HA2	1.56	0.87
1:B:1104:GLN:O	1:B:1104:GLN:HG3	1.79	0.82
1:B:1144:ARG:HD2	1:B:1311:MET:SD	2.23	0.78
1:A:1281:ARG:NH1	1:B:1259:SER:O	2.17	0.77
1:B:1152:LYS:HD3	4:B:1550:HOH:O	1.87	0.73
1:B:1156:LYS:O	1:B:1160:GLU:HG3	1.89	0.72
1:B:1199:GLU:HG2	1:B:1295:TYR:O	1.90	0.72
1:A:1262:GLN:NE2	1:A:1264:SER:O	2.23	0.71
1:A:1204:ILE:HD11	1:A:1268:MET:O	1.88	0.71
1:B:1220:LYS:NZ	1:B:1220:LYS:HB2	2.08	0.69
1:A:1199:GLU:HG2	1:A:1295:TYR:O	1.92	0.69
1:A:1260:PHE:HE2	1:A:1262:GLN:HG2	1.58	0.69
1:A:1146:ASN:ND2	1:A:1148:ILE:HD11	2.07	0.69
1:B:1204:ILE:HD11	1:B:1268:MET:O	1.93	0.69
1:B:1190:ASN:O	1:B:1194:HIS:HD2	1.77	0.67
1:B:1114:GLU:N	1:B:1114:GLU:OE1	2.23	0.67
1:B:1131:GLU:CD	1:B:1131:GLU:H	1.97	0.66
1:A:1171:GLU:HG3	4:A:1552:HOH:O	1.97	0.65
1:B:1148:ILE:HD11	1:B:1307:THR:HG22	1.78	0.65
1:B:1314:GLU:O	1:B:1315:HIS:HB2	1.96	0.65
1:B:1204:ILE:HD13	1:B:1209:GLY:HA2	1.81	0.62
1:A:1132:HIS:C	1:A:1134:ASP:H	2.01	0.62
1:B:1217:ASN:OD1	1:B:1289:TYR:CB	2.47	0.62
1:B:1176:HIS:HB3	4:B:1520:HOH:O	1.98	0.62
1:B:1178:ASN:ND2	1:B:1256:LEU:HB2	2.15	0.62
1:A:1117:GLU:OE2	1:A:1152:LYS:HE2	2.00	0.61
1:B:1232:THR:HG22	4:B:1508:HOH:O	2.00	0.60
1:A:1260:PHE:CE2	1:A:1262:GLN:HG2	2.36	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1234:CYS:SG	1:B:1237:HIS:HB2	2.43	0.58
1:B:1235:PRO:O	1:B:1238:LYS:NZ	2.35	0.58
1:B:1199:GLU:H	1:B:1199:GLU:CD	2.06	0.57
1:B:1167:LYS:O	1:B:1171:GLU:HG3	2.03	0.57
1:A:1199:GLU:HG3	1:A:1297:GLY:H	1.70	0.57
1:B:1216:GLU:HG3	1:B:1292:TYR:HE2	1.70	0.56
1:B:1263:PHE:CD2	1:B:1263:PHE:N	2.74	0.56
1:A:1204:ILE:HD13	1:A:1209:GLY:CA	2.30	0.56
1:A:1184:HIS:HE1	3:A:1402:2GV:H14	1.71	0.55
1:A:1252:CYS:HB3	1:A:1301:TYR:O	2.06	0.55
3:A:1402:2GV:O2	3:A:1402:2GV:H14	2.07	0.54
1:B:1314:GLU:O	1:B:1315:HIS:CB	2.55	0.54
1:A:1128:THR:HB	1:A:1217:ASN:HA	1.90	0.54
1:A:1266:MET:O	1:A:1267:LYS:CB	2.56	0.53
1:B:1220:LYS:HZ2	1:B:1220:LYS:HB2	1.74	0.52
1:A:1189:ILE:HD12	1:A:1250:LEU:HG	1.92	0.52
1:B:1148:ILE:HD11	1:B:1307:THR:CG2	2.40	0.51
1:B:1279:ILE:HG12	1:B:1292:TYR:CD1	2.45	0.51
1:B:1248:GLN:HA	1:B:1306:ILE:O	2.11	0.51
1:B:1258:LYS:HD3	1:B:1274:GLY:O	2.10	0.51
1:B:1220:LYS:HB2	1:B:1220:LYS:HZ3	1.75	0.50
1:A:1281:ARG:HD3	4:A:1524:HOH:O	2.12	0.49
1:A:1178:ASN:HA	1:B:1176:HIS:CE1	2.47	0.49
1:B:1267:LYS:N	4:B:1518:HOH:O	2.45	0.49
1:A:1184:HIS:CE1	3:A:1402:2GV:H14	2.48	0.49
1:B:1220:LYS:NZ	1:B:1220:LYS:CB	2.74	0.48
1:A:1213:TYR:OH	1:A:1265:THR:HB	2.13	0.48
1:A:1253:ARG:HB2	1:A:1303:GLU:OE1	2.13	0.48
1:A:1178:ASN:C	1:A:1178:ASN:HD22	2.16	0.47
1:A:1146:ASN:HD21	1:A:1148:ILE:HD11	1.76	0.47
1:B:1289:TYR:O	1:B:1290:ALA:HB3	2.14	0.47
1:B:1216:GLU:OE1	1:B:1290:ALA:HB3	2.14	0.47
1:B:1128:THR:HB	1:B:1217:ASN:HA	1.95	0.47
1:B:1186:SER:O	1:B:1189:ILE:HG12	2.14	0.47
1:B:1117:GLU:CD	1:B:1152:LYS:HZ1	2.19	0.47
1:A:1129:ILE:HD11	1:A:1145:TYR:CE2	2.50	0.46
1:B:1117:GLU:OE2	1:B:1152:LYS:NZ	2.49	0.46
1:A:1207:MET:CB	4:A:1566:HOH:O	2.62	0.46
1:A:1199:GLU:HG2	1:A:1296:ARG:HA	1.97	0.46
1:A:1134:ASP:HB3	1:A:1136:GLY:H	1.81	0.45
1:B:1117:GLU:CD	1:B:1152:LYS:NZ	2.69	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1222:ASN:HA	1:B:1225:VAL:HG23	1.99	0.45
1:A:1153:VAL:HB	1:A:1302:PRO:O	2.16	0.45
1:A:1113:PRO:HA	1:A:1118:TYR:CD2	2.52	0.45
1:B:1253:ARG:HG2	1:B:1253:ARG:HH11	1.81	0.45
1:A:1124:GLU:O	1:A:1128:THR:HG23	2.17	0.44
1:A:1260:PHE:CE2	1:A:1262:GLN:CG	3.01	0.44
1:A:1167:LYS:HG2	1:A:1171:GLU:OE2	2.17	0.44
1:A:1198:ASP:OD1	1:A:1200:ARG:HB2	2.17	0.44
1:B:1209:GLY:HA3	4:B:1564:HOH:O	2.17	0.44
3:B:1402:2GV:H14	3:B:1402:2GV:O2	2.17	0.44
1:B:1144:ARG:HD3	1:B:1145:TYR:O	2.17	0.44
1:A:1134:ASP:C	1:A:1136:GLY:H	2.21	0.43
1:A:1165:ARG:HB2	1:A:1298:GLU:HG3	2.00	0.43
1:B:1173:ASN:ND2	1:B:1176:HIS:O	2.31	0.43
1:A:1158:LEU:HD21	1:A:1196:GLY:HA3	2.00	0.43
1:B:1140:GLY:HA2	1:B:1241:SER:HB2	2.01	0.43
1:A:1262:GLN:OE1	1:A:1268:MET:SD	2.77	0.43
1:A:1188:PHE:O	1:A:1192:ILE:HG13	2.19	0.43
1:B:1149:ARG:HB3	1:B:1307:THR:HB	2.02	0.42
1:B:1253:ARG:HG2	1:B:1253:ARG:NH1	2.34	0.42
1:B:1197:PHE:HB3	1:B:1212:ILE:HD13	2.01	0.42
1:B:1258:LYS:HE2	4:B:1559:HOH:O	2.18	0.42
1:B:1117:GLU:OE1	1:B:1152:LYS:NZ	2.53	0.42
1:B:1158:LEU:HD21	1:B:1196:GLY:CA	2.50	0.42
1:A:1129:ILE:HD11	1:A:1145:TYR:CD2	2.55	0.42
1:A:1178:ASN:ND2	1:A:1178:ASN:C	2.73	0.42
1:A:1173:ASN:HB3	1:A:1176:HIS:O	2.21	0.41
1:A:1148:ILE:HG23	1:A:1148:ILE:O	2.19	0.41
3:B:1402:2GV:N4	3:B:1402:2GV:N5	2.63	0.41
1:B:1204:ILE:HD13	1:B:1209:GLY:CA	2.51	0.41
1:B:1298:GLU:H	1:B:1298:GLU:CD	2.24	0.40
1:A:1132:HIS:C	1:A:1134:ASP:N	2.72	0.40
1:B:1312:LYS:HB2	1:B:1312:LYS:HE2	1.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	210/217 (97%)	204 (97%)	6 (3%)	0	100 100
1	B	191/217 (88%)	185 (97%)	5 (3%)	1 (0%)	34 40
All	All	401/434 (92%)	389 (97%)	11 (3%)	1 (0%)	52 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1290	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	176/185 (95%)	164 (93%)	12 (7%)	20 25
1	B	166/185 (90%)	155 (93%)	11 (7%)	21 26
All	All	342/370 (92%)	319 (93%)	23 (7%)	20 26

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1104	GLN
1	A	1123	GLU
1	A	1144	ARG
1	A	1148	ILE
1	A	1161	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1178	ASN
1	A	1199	GLU
1	A	1216	GLU
1	A	1252	CYS
1	A	1253	ARG
1	A	1283	SER
1	A	1303	GLU
1	B	1104	GLN
1	B	1110	ASP
1	B	1131	GLU
1	B	1144	ARG
1	B	1152	LYS
1	B	1199	GLU
1	B	1208	PHE
1	B	1220	LYS
1	B	1249	MET
1	B	1263	PHE
1	B	1312	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1146	ASN
1	A	1166	GLN
1	A	1173	ASN
1	A	1178	ASN
1	A	1262	GLN
1	B	1104	GLN
1	B	1143	ASN
1	B	1176	HIS
1	B	1194	HIS
1	B	1262	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	2GV	A	1402	-	37,37,37	3.52	23 (62%)	51,54,54	3.59	28 (54%)
3	2GV	B	1402	-	37,37,37	3.53	23 (62%)	51,54,54	3.63	22 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2GV	A	1402	-	-	0/16/35/35	0/5/5/5
3	2GV	B	1402	-	-	0/16/35/35	0/5/5/5

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1402	2GV	C8-C7	-7.54	1.47	1.55
3	A	1402	2GV	C8-C7	-6.70	1.48	1.55
3	A	1402	2GV	C19-N2	-6.39	1.31	1.43
3	B	1402	2GV	C19-N2	-5.78	1.32	1.43
3	A	1402	2GV	C17-C7	-2.56	1.47	1.51
3	B	1402	2GV	C17-C7	-2.20	1.48	1.51
3	B	1402	2GV	C18-C22	2.10	1.42	1.38
3	A	1402	2GV	C18-C22	2.16	1.42	1.38
3	A	1402	2GV	C11-N2	2.37	1.39	1.36
3	A	1402	2GV	C15-C16	2.48	1.44	1.38
3	A	1402	2GV	C24-C25	2.53	1.46	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1402	2GV	C3-C2	2.60	1.45	1.37
3	B	1402	2GV	C15-C16	2.69	1.44	1.38
3	B	1402	2GV	C3-C4	2.70	1.45	1.37
3	B	1402	2GV	C7-N2	2.72	1.53	1.48
3	A	1402	2GV	C13-C12	2.81	1.44	1.38
3	B	1402	2GV	C15-C14	2.84	1.45	1.38
3	A	1402	2GV	C3-C2	2.86	1.46	1.37
3	B	1402	2GV	C13-C14	2.86	1.45	1.38
3	A	1402	2GV	C3-C4	2.92	1.46	1.37
3	B	1402	2GV	C13-C12	3.01	1.45	1.38
3	A	1402	2GV	C20-C19	3.14	1.45	1.39
3	A	1402	2GV	C18-C19	3.22	1.45	1.39
3	B	1402	2GV	C18-C19	3.23	1.45	1.39
3	B	1402	2GV	C21-C5	3.24	1.46	1.39
3	A	1402	2GV	O1-C11	3.37	1.41	1.35
3	A	1402	2GV	C15-C14	3.47	1.46	1.38
3	A	1402	2GV	C13-C14	3.50	1.47	1.38
3	A	1402	2GV	C21-C5	3.57	1.47	1.39
3	B	1402	2GV	C20-C19	3.73	1.46	1.39
3	B	1402	2GV	C23-C1	3.83	1.46	1.39
3	A	1402	2GV	C12-C17	3.99	1.45	1.39
3	A	1402	2GV	C22-C5	3.99	1.47	1.39
3	B	1402	2GV	C22-C5	4.07	1.48	1.39
3	B	1402	2GV	O1-C8	4.23	1.55	1.48
3	B	1402	2GV	C26-C1	4.39	1.47	1.39
3	A	1402	2GV	C26-C1	4.66	1.47	1.39
3	B	1402	2GV	C12-C17	4.97	1.47	1.39
3	B	1402	2GV	C16-C17	5.38	1.47	1.39
3	A	1402	2GV	C16-C17	5.76	1.48	1.39
3	A	1402	2GV	C6-N5	5.80	1.42	1.34
3	B	1402	2GV	C6-N1	6.53	1.43	1.34
3	B	1402	2GV	O1-C11	6.63	1.46	1.35
3	B	1402	2GV	C6-N5	6.76	1.43	1.34
3	A	1402	2GV	O1-C8	7.30	1.59	1.48
3	A	1402	2GV	C6-N1	7.52	1.44	1.34

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1402	2GV	O1-C8-C7	-10.02	94.14	102.73
3	B	1402	2GV	O1-C11-N2	-7.45	103.17	108.71
3	A	1402	2GV	O1-C8-C7	-7.32	96.45	102.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1402	2GV	C4-N5-C6	7.07	122.69	115.93
3	A	1402	2GV	O2-C11-N2	8.10	139.59	128.71
3	B	1402	2GV	C24-C6-N1	8.11	124.03	116.85
3	A	1402	2GV	C24-C6-N1	8.74	124.59	116.85
3	B	1402	2GV	C8-C7-N2	8.78	108.48	100.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1402	2GV	3	0
3	B	1402	2GV	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/217 (97%)	0.19	9 (4%) 40 48	17, 26, 37, 41	0
1	B	199/217 (91%)	0.29	12 (6%) 25 33	18, 29, 39, 44	0
All	All	411/434 (94%)	0.24	21 (5%) 32 41	17, 27, 38, 44	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1315	HIS	6.2
1	B	1104	GLN	5.8
1	A	1206	GLY	5.4
1	B	1264	SER	5.2
1	A	1104	GLN	4.0
1	A	1315	HIS	3.9
1	A	1207	MET	3.9
1	B	1263	PHE	3.7
1	B	1204	ILE	3.5
1	A	1284	VAL	3.2
1	B	1289	TYR	3.0
1	B	1208	PHE	2.7
1	B	1131	GLU	2.6
1	A	1131	GLU	2.6
1	B	1133	ARG	2.5
1	A	1167	LYS	2.4
1	B	1171	GLU	2.4
1	B	1134	ASP	2.3
1	A	1156	LYS	2.2
1	A	1116	LYS	2.1
1	B	1314	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	2GV	A	1402	33/33	0.86	0.27	4.22	39,49,63,67	0
3	2GV	B	1402	33/33	0.85	0.26	2.47	44,53,67,71	0
2	ZN	A	1401	1/1	0.99	0.12	1.47	24,24,24,24	0
2	ZN	B	1401	1/1	0.98	0.10	-0.92	30,30,30,30	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.