wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 03:59 AM BST

PDB ID	:	2N6A
Title	:	NMR structure of a human calmodulin/connexin-36 peptide hybrid
Authors	:	Donaldson, L.
Deposited on	:	2015-08-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 69%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	173	70% 22% •• 6%

2 Ensemble composition and analysis (i)

This entry contains 18 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:8-A:168 (161)	0.50	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 3, 5, 7, 10, 11, 12, 18
2	4, 14, 15, 17
3	1, 8
Single-model clusters	6; 9; 13; 16

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2235 atoms, of which 975 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called human calmodulin/connexin-36 peptide hybrid.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	162	2231	772	975	211	264	9	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	HIS	-	EXPRESSION TAG	UNP P62158
A	0	HIS	-	EXPRESSION TAG	UNP P62158
A	1	HIS	-	EXPRESSION TAG	UNP P62158
A	2	HIS	-	EXPRESSION TAG	UNP P62158
A	3	HIS	-	EXPRESSION TAG	UNP P62158
A	4	HIS	-	EXPRESSION TAG	UNP P62158
A	147	GLY	-	LINKER	UNP P62158
A	148	ALA	-	LINKER	UNP P62158
A	149	SER	-	LINKER	UNP P62158
A	150	THR	-	LINKER	UNP P62158
A	151	ALA	-	LINKER	UNP P62158
A	152	ALA	-	LINKER	UNP P62158
A	153	GLY	-	LINKER	UNP P62158
A	154	SER	-	LINKER	UNP P62158

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	
			Total	Ca
2	A	4	4	4

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: human calmodulin/connexin-36 peptide hybrid

Chain A: 70% 22% • • 6%

HIS HIS HIS HIS HIS HIS THR GLU GLU Q8 A15 L18 F19 D22 T29 K30 E31 L32 V35 M36 R37 S38 L39 L48 I52 D58 I63 D64 F65 P66 L69 I85 R86 E87 A88 V91 F92 I100 E104 L105 R106 H107 V108 L112

L116 D131 A148 A151 W156 R157 K158 I159 K160 L161 A162 V163 A166 K169 ARG LYS

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

- Molecule 1: human calmodulin/connexin-36 peptide hybrid

Chain A: 66% 24% • • 6%

HIS HIS HIS HIS HIS HIS THR GLU GLU Q8 A15 F16 F19 D22 I27 E31 L32 V35 M36 R37 S38 L39 L48 I52 D56 F65 P66 E67 F68 L69 K75 M76 K77 D78 I85 R86 E87 A88 V91 F92 D95 A102 A103 E104

V108 L112 L116 T117 V121 M124 E127 D131 G132 D133 M145 A148 A151 A152 I159 K160 L161 A162 V163 A166 K169 ARG LYS

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 400 calculated structures, 18 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2n6a_cs.str
Number of chemical shift lists	1
Total number of shifts	1575
Number of shifts mapped to atoms	1575
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	69%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1247	966	1188	24±3
All	All	22518	17388	21384	434

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

5 of 124 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:15:ALA:HB2	1:A:161:LEU:HD22	0.98	1.35	2	12
1:A:15:ALA:HB2	1:A:161:LEU:HD23	0.98	1.31	3	3
1:A:100:ILE:HG21	1:A:105:LEU:HD13	0.79	1.51	6	11
1:A:18:LEU:HD11	1:A:162:ALA:HB2	0.75	1.57	11	6
1:A:148:ALA:HB1	1:A:151:ALA:HB3	0.75	1.59	15	11

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	160/173 (92%)	146±3 (91±2%)	13±3 (8±2%)	1±0 (1±0%)	31 76
All	All	2880/3114 (92%)	2623 (91%)	238 (8%)	19 (1%)	31 76

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	85	ILE	17
1	A	114	GLU	1
1	A	78	ASP	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	131/143 (92%)	113±4 (87±3%)	18±4 (13±3%)	9 50
All	All	2358/2574 (92%)	2041 (87%)	317 (13%)	9 50

5 of 71 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	32	LEU	18
1	A	112	LEU	16
1	A	58	ASP	15
1	A	19	PHE	14
1	A	131	ASP	14

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 69% for the well-defined parts and 69% for the entire structure.

7.1 Chemical shift list 1

File name: 2n6a_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1575
Number of shifts mapped to atoms	1575
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	158	-0.33 ± 0.12	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	144	0.16 ± 0.09	None needed (< 0.5 ppm)
$^{13}C'$	155	-0.49 ± 0.06	None needed (< 0.5 ppm)
^{15}N	151	0.36 ± 0.29	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 69%, i.e. 1318 atoms were assigned a chemical shift out of a possible 1899. 0 out of 17 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	726/801 (91%)	282/320 (88%)	299/322 (93%)	145/159 (91%)
Sidechain	524/992 (53%)	294/574 (51%)	230/373 (62%)	0/45 (0%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	68/106 (64%)	34/58 (59%)	33/47 (70%)	1/1 (100%)
Overall	1318/1899 (69%)	610/952 (64%)	562/742 (76%)	146/205 (71%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	93	ASP	HB2	1.32	4.07 – 1.37	-5.2

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: