



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:55 PM GMT

PDB ID : 4N84
Title : Crystal structure of 14-3-3zeta in complex with a 12-carbon-linker cyclic peptide derived from ExoS
Authors : Bier, D.; Glas, A.; Hahne, G.; Grossmann, T.; Ottmann, C.
Deposited on : 2013-10-17
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	 4% 70% 24% . .
1	B	230	 3% 75% 20% . .
2	C	10	 40% 20% 40%
2	E	10	 20% 40% 40%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MKD	C	422	-	-	X	-
2	MKD	C	425	-	-	X	-
2	MKD	E	422	-	-	X	-
2	MKD	E	425	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3760 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein zeta/delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1793	1123	304	356	10			
1	B	226	Total	C	N	O	S	0	0	0
			1771	1109	297	355	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	EXPRESSION TAG	UNP P63104
A	?	-	LEU	DELETION	UNP P63104
B	0	GLY	-	EXPRESSION TAG	UNP P63104
B	?	-	LEU	DELETION	UNP P63104

- Molecule 2 is a protein called Exoenzyme S.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	0	0	0
			77	52	10	15			
2	E	10	Total	C	N	O	0	0	0
			77	52	10	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	422	MKD	LEU	ENGINEERED MUTATION	UNP Q93SQ3
C	425	MKD	ALA	ENGINEERED MUTATION	UNP Q93SQ3
E	422	MKD	LEU	ENGINEERED MUTATION	UNP Q93SQ3
E	425	MKD	ALA	ENGINEERED MUTATION	UNP Q93SQ3


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total 18	O 18	0	0
3	B	14	Total 14	O 14	0	0
3	C	5	Total 5	O 5	0	0
3	E	5	Total 5	O 5	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3 protein zeta/delta




- Molecule 1: 14-3-3 protein zeta/delta



- Molecule 2: Exoenzyme S



- Molecule 2: Exoenzyme S



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.24Å 105.18Å 113.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.74 – 2.50 47.74 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.74-2.50) 100.0 (47.74-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.86 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.202 , 0.249 0.218 , 0.256	Depositor DCC
R_{free} test set	1764 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	52.9	Xtriage
Anisotropy	0.453	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	6 of 35305 reflections (0.017%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3760	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MKD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	2/1816 (0.1%)	1.05	4/2442 (0.2%)
1	B	0.95	1/1793 (0.1%)	0.98	3/2415 (0.1%)
2	C	1.14	0/52	0.94	0/66
2	E	0.86	0/52	1.01	0/66
All	All	0.97	3/3713 (0.1%)	1.01	7/4989 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	4
2	E	0	4
All	All	0	8

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	102	GLU	CD-OE2	5.84	1.32	1.25
1	A	59	TRP	CD2-CE2	5.43	1.47	1.41
1	B	59	TRP	CD2-CE2	5.00	1.47	1.41

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	20	ASP	CB-CG-OD2	5.98	123.68	118.30
1	B	104	PHE	N-CA-C	5.80	126.65	111.00
1	A	75	LYS	CB-CA-C	5.73	121.85	110.40
1	B	124	ASP	CB-CG-OD1	5.71	123.44	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	ASP	CB-CA-C	-5.68	99.03	110.40
1	B	69	THR	N-CA-CB	-5.57	99.72	110.30
1	A	91	ARG	NE-CZ-NH1	-5.10	117.75	120.30

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	421	GLY	Mainchain
2	C	422	MKD	Mainchain
2	C	424	ASP	Mainchain
2	C	425	MKD	Mainchain
2	E	421	GLY	Mainchain
2	E	422	MKD	Mainchain
2	E	424	ASP	Mainchain
2	E	425	MKD	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1793	0	1774	39	0
1	B	1771	0	1713	42	0
2	C	77	0	85	21	0
2	E	77	0	85	16	0
3	A	18	0	0	1	0
3	B	14	0	0	1	0
3	C	5	0	0	0	0
3	E	5	0	0	1	0
All	All	3760	0	3657	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:422:MKD:H17	2:C:425:MKD:CH	1.17	1.55
2:E:422:MKD:H17	2:E:425:MKD:CH	1.04	1.51
2:C:422:MKD:CH	2:C:425:MKD:H17	1.00	1.47
2:C:422:MKD:CF	2:C:425:MKD:CH	2.14	1.25
2:E:422:MKD:CH	2:E:425:MKD:H17	1.67	1.11
2:E:422:MKD:H17	2:E:425:MKD:H18	1.18	1.10
2:E:422:MKD:H17	2:E:425:MKD:CF	1.83	1.08
2:C:422:MKD:CF	2:C:425:MKD:H19	1.83	1.04
1:B:48:TYR:O	1:B:52:VAL:HG23	1.56	1.02
2:E:422:MKD:CH	2:E:425:MKD:H18	1.77	0.96
2:E:422:MKD:CF	2:E:425:MKD:H18	1.98	0.93
2:C:422:MKD:CF	2:C:425:MKD:H17	1.88	0.93
2:C:422:MKD:H16	2:C:425:MKD:H19	1.53	0.87
1:A:116:VAL:HG21	1:A:160:MET:CE	2.06	0.86
2:E:422:MKD:H17	2:E:425:MKD:H17	0.86	0.84
1:B:49:LYS:HE2	3:E:505:HOH:O	1.77	0.83
1:B:52:VAL:HG22	1:B:93:ILE:HG21	1.59	0.82
1:B:1:MET:HB3	1:B:5:GLU:OE1	1.80	0.82
1:A:69:THR:O	1:A:76:GLN:HG3	1.78	0.82
1:A:164:HIS:HD2	1:A:166:ILE:H	1.26	0.82
2:C:422:MKD:H17	2:C:425:MKD:CF	2.11	0.81
2:C:422:MKD:H15	2:C:425:MKD:CH	2.08	0.80
1:B:196:PHE:CE2	1:B:200:ILE:HD11	2.17	0.80
2:C:422:MKD:CH	2:C:425:MKD:CF	2.60	0.80
1:B:168:LEU:HB3	1:B:217:ILE:HG21	1.63	0.79
1:B:163:THR:OG1	1:B:208:GLU:HG2	1.85	0.76
1:B:52:VAL:HG22	1:B:93:ILE:CG2	2.16	0.76
2:C:422:MKD:H15	2:C:425:MKD:H15	1.69	0.74
1:A:190:SER:O	1:A:194:THR:HG23	1.87	0.74
2:C:422:MKD:H15	2:C:425:MKD:CF	2.18	0.73
1:A:156:SER:HA	1:A:160:MET:HE3	1.70	0.73
1:A:116:VAL:HG21	1:A:160:MET:HE2	1.70	0.73
1:B:65:ILE:O	1:B:69:THR:HG23	1.89	0.72
1:A:116:VAL:HG11	1:A:160:MET:HE1	1.73	0.69
1:B:196:PHE:CE2	1:B:200:ILE:CD1	2.76	0.69
1:A:72:ALA:HB3	1:A:75:LYS:HB2	1.76	0.68
1:B:211:TYR:CB	1:B:214:SER:H	2.07	0.68
1:A:72:ALA:CB	1:A:75:LYS:HB2	2.27	0.65
1:A:18:ARG:NH2	1:B:89:GLU:OE2	2.27	0.64
1:A:183:ASN:O	1:A:185:PRO:HD3	1.98	0.64
1:B:227:LEU:O	1:B:227:LEU:HD22	1.97	0.64
2:E:422:MKD:CH	2:E:425:MKD:CF	2.58	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:421:GLY:O	2:C:422:MKD:H2	1.99	0.62
1:B:182:LEU:O	1:B:183:ASN:HB2	1.97	0.62
1:B:52:VAL:HG21	1:B:125:TYR:HE2	1.63	0.62
2:E:422:MKD:CH	2:E:425:MKD:H16	2.28	0.61
1:A:83:ARG:O	1:A:87:GLU:HG3	2.01	0.61
1:A:127:ARG:HD2	3:A:304:HOH:O	2.00	0.60
1:A:211:TYR:O	1:A:215:THR:HB	2.04	0.57
1:B:196:PHE:O	1:B:200:ILE:HD12	2.05	0.56
1:A:203:LEU:O	1:A:203:LEU:HD23	2.05	0.56
1:B:132:VAL:CG1	1:B:132:VAL:O	2.54	0.55
1:A:45:SER:HB3	2:C:425:MKD:H3	1.89	0.54
1:A:22:MET:HG2	1:A:47:ALA:HB2	1.90	0.54
2:C:422:MKD:H15	2:C:425:MKD:H19	1.76	0.54
1:B:65:ILE:O	1:B:69:THR:CG2	2.56	0.54
2:E:429:ALA:O	2:E:430:SER:C	2.45	0.54
1:A:116:VAL:CG2	1:A:160:MET:HE2	2.38	0.53
2:E:421:GLY:O	2:E:424:ASP:HB3	2.07	0.53
1:A:164:HIS:CD2	1:A:166:ILE:H	2.16	0.53
2:C:421:GLY:O	2:C:422:MKD:C	2.57	0.53
2:C:421:GLY:O	2:C:423:LEU:N	2.42	0.53
1:B:202:GLU:OE2	1:B:208:GLU:OE2	2.27	0.52
1:A:84:GLU:O	1:A:88:THR:HG23	2.11	0.51
1:B:183:ASN:O	1:B:185:PRO:HD3	2.11	0.51
1:B:59:TRP:O	1:B:63:SER:HB2	2.11	0.50
1:B:190:SER:O	1:B:194:THR:HG23	2.11	0.50
2:E:422:MKD:H16	2:E:425:MKD:H18	1.89	0.50
1:B:196:PHE:CZ	1:B:200:ILE:HD11	2.46	0.50
2:E:422:MKD:H15	2:E:425:MKD:H18	1.90	0.50
2:C:422:MKD:CF	2:C:425:MKD:CF	2.75	0.50
1:B:211:TYR:CB	1:B:214:SER:HB2	2.42	0.50
1:A:138:LYS:O	1:A:138:LYS:HD3	2.11	0.49
1:B:127:ARG:HD2	3:B:312:HOH:O	2.12	0.49
1:A:183:ASN:C	1:A:185:PRO:HD3	2.33	0.49
1:B:48:TYR:O	1:B:52:VAL:CG2	2.45	0.49
1:B:84:GLU:O	1:B:88:THR:HG23	2.13	0.48
1:B:52:VAL:HG21	1:B:125:TYR:CE2	2.46	0.48
1:A:227:LEU:O	1:A:230:SER:HB2	2.13	0.48
1:B:163:THR:HG1	1:B:208:GLU:HG2	1.75	0.48
2:E:426:LEU:HD23	2:E:426:LEU:HA	1.76	0.48
2:C:422:MKD:H16	2:C:425:MKD:CH	2.18	0.47
1:A:78:MET:HE1	1:B:8:GLN:HB3	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:422:MKD:H18	2:E:425:MKD:H16	1.95	0.47
1:A:133:ALA:HB3	1:A:138:LYS:HG2	1.96	0.47
2:C:422:MKD:CF	2:C:425:MKD:H15	2.42	0.47
1:A:164:HIS:CD2	1:A:165:PRO:HD2	2.50	0.47
1:B:202:GLU:HG3	1:B:208:GLU:OE2	2.15	0.46
1:A:6:LEU:HD13	1:A:28:SER:HB2	1.97	0.46
1:B:59:TRP:CE2	1:B:132:VAL:HG22	2.50	0.45
2:C:424:ASP:O	2:C:425:MKD:H2	2.16	0.45
1:B:132:VAL:O	1:B:132:VAL:HG13	2.17	0.45
1:A:72:ALA:HB3	1:A:75:LYS:CB	2.46	0.44
1:A:116:VAL:HG21	1:A:160:MET:HE1	1.93	0.44
1:B:196:PHE:CD2	1:B:200:ILE:CD1	3.02	0.43
1:B:202:GLU:HA	1:B:202:GLU:OE1	2.17	0.43
1:B:209:GLU:HA	1:B:210:SER:HA	1.74	0.43
1:B:213:ASP:HB3	2:E:423:LEU:HD21	2.00	0.43
1:A:225:LEU:HD23	1:A:225:LEU:HA	1.80	0.43
1:B:196:PHE:CE2	1:B:200:ILE:HD13	2.52	0.43
1:B:200:ILE:O	1:B:203:LEU:HG	2.19	0.42
1:B:6:LEU:HD13	1:B:28:SER:HB2	2.01	0.42
1:A:78:MET:HB3	1:A:78:MET:HE3	1.69	0.42
1:B:141:ILE:O	1:B:144:GLN:HB2	2.19	0.42
1:A:161:GLN:HA	1:A:161:GLN:OE1	2.19	0.42
1:B:202:GLU:CG	1:B:208:GLU:OE2	2.68	0.42
1:A:138:LYS:O	1:A:142:VAL:HG23	2.20	0.42
1:A:12:LEU:HD23	1:A:12:LEU:HA	1.86	0.42
1:A:186:GLU:CD	1:A:186:GLU:H	2.23	0.41
1:A:143:ASP:O	1:A:147:GLN:HB2	2.19	0.41
1:A:132:VAL:O	1:A:132:VAL:HG13	2.21	0.41
2:C:422:MKD:H19	2:C:422:MKD:H12	1.84	0.41
1:A:140:GLY:O	1:A:144:GLN:HG3	2.21	0.41
1:A:91:ARG:HG2	1:A:129:LEU:CD2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/230 (97%)	219 (98%)	4 (2%)	0	100	100
1	B	220/230 (96%)	217 (99%)	3 (1%)	0	100	100
2	C	6/10 (60%)	5 (83%)	1 (17%)	0	100	100
2	E	6/10 (60%)	6 (100%)	0	0	100	100
All	All	455/480 (95%)	447 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/199 (96%)	166 (87%)	25 (13%)	5	9
1	B	185/199 (93%)	168 (91%)	17 (9%)	11	21
2	C	6/6 (100%)	5 (83%)	1 (17%)	3	5
2	E	6/6 (100%)	6 (100%)	0	100	100
All	All	388/410 (95%)	345 (89%)	43 (11%)	8	14

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	9	LYS
1	A	49	LYS
1	A	64	SER
1	A	75	LYS
1	A	76	GLN
1	A	88	THR
1	A	99	SER
1	A	100	LEU
1	A	113	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	114	SER
1	A	132	VAL
1	A	136	ASP
1	A	138	LYS
1	A	147	GLN
1	A	154	GLU
1	A	168	LEU
1	A	187	LYS
1	A	193	LYS
1	A	194	THR
1	A	202	GLU
1	A	215	THR
1	A	216	LEU
1	A	222	ARG
1	A	229	THR
1	B	1	MET
1	B	60	ARG
1	B	64	SER
1	B	70	GLU
1	B	76	GLN
1	B	88	THR
1	B	99	SER
1	B	100	LEU
1	B	132	VAL
1	B	137	ASP
1	B	145	SER
1	B	158	LYS
1	B	168	LEU
1	B	184	SER
1	B	200	ILE
1	B	208	GLU
1	B	229	THR
2	C	430	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	A	147	GLN
1	A	164	HIS
1	B	50	ASN
1	B	146	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	183	ASN
1	B	224	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MKD	C	422	2	7,10,11	0.94	0	6,12,14	0.96	1 (16%)
2	MKD	C	425	2	7,10,11	0.79	0	6,12,14	1.19	1 (16%)
2	MKD	E	422	2	7,10,11	0.93	0	6,12,14	1.29	1 (16%)
2	MKD	E	425	2	7,10,11	1.43	2 (28%)	6,12,14	1.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MKD	C	422	2	-	0/7/10/13	0/0/0/0
2	MKD	C	425	2	-	0/7/10/13	0/0/0/0
2	MKD	E	422	2	-	0/7/10/13	0/0/0/0
2	MKD	E	425	2	-	0/7/10/13	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	425	MKD	O-C	2.22	1.27	1.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	425	MKD	CB-CA	3.02	1.58	1.55

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	425	MKD	CE-CD-CG	-2.62	101.03	114.53
2	C	422	MKD	CB1-CA-CB	2.18	114.78	110.92
2	E	422	MKD	CH-CF-CE	2.47	132.88	113.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	422	MKD	18	0
2	C	425	MKD	17	0
2	E	422	MKD	12	0
2	E	425	MKD	12	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/230 (98%)	0.15	10 (4%) 38 43	34, 49, 92, 120	0
1	B	226/230 (98%)	0.10	8 (3%) 48 53	30, 48, 91, 122	0
2	C	8/10 (80%)	0.06	0 100 100	39, 55, 66, 86	0
2	E	8/10 (80%)	-0.05	0 100 100	42, 58, 73, 88	0
All	All	469/480 (97%)	0.12	18 (3%) 44 49	30, 49, 92, 122	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	GLU	6.9
1	B	0	GLY	6.2
1	B	134	ALA	5.7
1	B	1	MET	5.2
1	A	203	LEU	4.4
1	A	71	GLY	3.8
1	B	203	LEU	3.6
1	A	75	LYS	3.5
1	B	208	GLU	3.1
1	A	204	ASP	2.8
1	B	52	VAL	2.7
1	A	136	ASP	2.6
1	A	72	ALA	2.4
1	A	134	ALA	2.2
1	B	230	SER	2.1
1	A	76	GLN	2.1
1	B	210	SER	2.1
1	A	137	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MKD	C	425	11/12	0.96	0.14	-	48,51,54,57	0
2	MKD	E	422	11/12	0.90	0.18	-	47,62,78,85	0
2	MKD	E	425	11/12	0.97	0.14	-	43,47,54,59	0
2	MKD	C	422	11/12	0.96	0.15	-	56,66,71,80	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.