



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:50 AM GMT

PDB ID : 3N98
Title : Crystal structure of TK1436, a GH57 branching enzyme from hyperthermophilic archaeon *Thermococcus kodakaraensis*, in complex with glucose and additives
Authors : Santos, C.R.; Tonoli, C.C.C.; Trindade, D.M.; Betzel, C.; Takata, H.; Kuriki, T.; Kanai, T.; Imanaka, T.; Arni, R.K.; Murakami, M.T.
Deposited on : 2010-05-28
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6965 (1.90-1.86)
Clashscore	102246	7778 (1.90-1.86)
Ramachandran outliers	100387	7691 (1.90-1.86)
Sidechain outliers	100360	7692 (1.90-1.86)
RSRZ outliers	91569	6979 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	<div> <div>4%</div> <div>80%</div> <div>16%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BGC	A	564	-	-	X	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	566	-	-	-	X
4	DIO	A	569	-	-	-	X
4	DIO	A	570	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 5128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-amylase, GH57 family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	550	4694	3033	796	853	12	0	25	0

- Molecule 2 is glucose (three-letter code: BGC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	12	6	6	0	0
2	A	1	12	6	6	0	0
2	A	1	12	6	6	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 1,4-DIETHYLENE DIOXIDE (three-letter code: DIO) (formula: C₄H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	4	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	348	Total	O	0	0
			348	348		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha-amylase, GH57 family



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.67Å 79.02Å 134.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.08 – 1.87 33.52 – 1.87	Depositor EDS
% Data completeness (in resolution range)	95.4 (68.08-1.87) 95.4 (33.52-1.87)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.87Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.159 , 0.196 0.157 , 0.195	Depositor DCC
R_{free} test set	3024 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 70.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 59878 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5128	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, PG4, BGC, DIO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.43	23/4945 (0.5%)	1.16	23/6689 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	364	GLU	CB-CG	6.19	1.64	1.52
1	A	158	ARG	CG-CD	6.11	1.67	1.51
1	A	310	GLU	CB-CG	6.09	1.63	1.52
1	A	274	TYR	CD1-CE1	6.02	1.48	1.39
1	A	333	GLU	CG-CD	5.90	1.60	1.51
1	A	326	ARG	CG-CD	5.55	1.65	1.51
1	A	439	VAL	CB-CG2	5.41	1.64	1.52
1	A	526	VAL	CB-CG2	5.40	1.64	1.52
1	A	363	PHE	CE1-CZ	5.34	1.47	1.37
1	A	407	TRP	CE3-CZ3	5.31	1.47	1.38
1	A	27	GLU	CB-CG	5.29	1.62	1.52
1	A	172	PHE	CE1-CZ	5.27	1.47	1.37
1	A	31	GLU	CB-CG	5.23	1.62	1.52
1	A	502	TYR	CE1-CZ	5.22	1.45	1.38
1	A	256	VAL	CB-CG2	5.21	1.63	1.52
1	A	332	VAL	CB-CG2	5.21	1.63	1.52
1	A	143	ALA	CA-CB	5.20	1.63	1.52
1	A	298	ARG	CB-CG	5.18	1.66	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	168	TYR	CG-CD1	5.14	1.45	1.39
1	A	138	VAL	CB-CG2	5.13	1.63	1.52
1	A	35[A]	GLU	CD-OE2	-5.10	1.20	1.25
1	A	35[B]	GLU	CD-OE2	-5.10	1.20	1.25
1	A	389	PHE	CE1-CZ	5.01	1.46	1.37

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	490	ARG	NE-CZ-NH1	13.36	126.98	120.30
1	A	490	ARG	NE-CZ-NH2	-10.42	115.09	120.30
1	A	128	ARG	NE-CZ-NH2	-9.53	115.53	120.30
1	A	247[A]	ARG	NE-CZ-NH2	-8.42	116.09	120.30
1	A	247[B]	ARG	NE-CZ-NH2	-8.42	116.09	120.30
1	A	371	ARG	NE-CZ-NH2	-7.86	116.37	120.30
1	A	198	ARG	NE-CZ-NH2	-7.69	116.46	120.30
1	A	509	ASP	CB-CG-OD2	6.68	124.31	118.30
1	A	264	GLU	OE1-CD-OE2	-6.51	115.49	123.30
1	A	198	ARG	NE-CZ-NH1	6.37	123.48	120.30
1	A	298	ARG	NE-CZ-NH1	6.26	123.43	120.30
1	A	66	GLU	OE1-CD-OE2	-6.16	115.91	123.30
1	A	337	ARG	NE-CZ-NH1	6.13	123.36	120.30
1	A	495	LEU	CB-CG-CD2	6.02	121.23	111.00
1	A	225	ASP	CB-CG-OD1	5.96	123.67	118.30
1	A	283[A]	ARG	NE-CZ-NH1	-5.59	117.50	120.30
1	A	283[B]	ARG	NE-CZ-NH1	-5.59	117.50	120.30
1	A	298	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	A	128	ARG	NE-CZ-NH1	5.53	123.07	120.30
1	A	215	ARG	NE-CZ-NH1	5.20	122.90	120.30
1	A	412	ASP	CB-CG-OD1	5.17	122.96	118.30
1	A	509	ASP	CB-CG-OD1	-5.16	113.65	118.30
1	A	48	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	305	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4694	0	4551	73	0
2	A	36	0	36	11	0
3	A	18	0	24	2	0
4	A	12	0	16	3	0
5	A	7	0	10	0	0
6	A	13	0	18	3	0
7	A	348	0	0	25	0
All	All	5128	0	4655	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270[B]:TRP:HZ2	7:A:911:HOH:O	1.21	1.20
1:A:169[B]:GLU:OE2	1:A:175:LYS:HG2	1.46	1.13
1:A:270[B]:TRP:HZ3	7:A:617:HOH:O	1.34	1.08
1:A:483[A]:ARG:NH1	1:A:549:PRO:O	1.90	1.03
1:A:270[B]:TRP:CZ2	7:A:911:HOH:O	2.02	0.99
6:A:572:PG4:H61	7:A:880:HOH:O	1.63	0.97
1:A:247[B]:ARG:NH1	1:A:398:HIS:CE1	2.33	0.95
1:A:169[B]:GLU:OE2	1:A:175:LYS:CG	2.16	0.93
1:A:270[B]:TRP:CZ3	7:A:617:HOH:O	2.14	0.92
1:A:247[B]:ARG:HH11	1:A:398:HIS:CE1	1.86	0.91
1:A:476:GLN:HE22	2:A:564:BGC:C6	1.84	0.89
3:A:567:GOL:O3	7:A:911:HOH:O	1.79	0.86
1:A:129[B]:GLU:HG2	1:A:133:GLN:NE2	1.95	0.80
1:A:476:GLN:HE22	2:A:564:BGC:H6C1	1.51	0.75
1:A:42[A]:MET:CE	7:A:862:HOH:O	2.36	0.74
1:A:169[B]:GLU:OE2	1:A:175:LYS:CD	2.35	0.74
1:A:224:ILE:HD12	1:A:260:ALA:HB2	1.70	0.74
2:A:565:BGC:O6	7:A:868:HOH:O	2.05	0.73
1:A:39:PRO:HA	1:A:42[A]:MET:CE	2.19	0.73
1:A:283[B]:ARG:HD2	1:A:295:GLN:HB3	1.71	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:572:PG4:H42	7:A:880:HOH:O	1.91	0.70
1:A:243[B]:LYS:HE2	7:A:913:HOH:O	1.92	0.69
1:A:77:PHE:CZ	1:A:81:MET:HG2	2.29	0.67
6:A:572:PG4:H21	7:A:693:HOH:O	1.93	0.67
1:A:39:PRO:HA	1:A:42[A]:MET:HE2	1.79	0.65
1:A:42[A]:MET:HE1	7:A:862:HOH:O	1.95	0.64
2:A:563:BGC:H3	7:A:893:HOH:O	1.97	0.63
1:A:476:GLN:NE2	2:A:564:BGC:C6	2.60	0.62
1:A:79[A]:GLU:HG2	1:A:83:ARG:NH2	2.15	0.61
1:A:129[B]:GLU:CG	1:A:133:GLN:NE2	2.64	0.61
1:A:205:ILE:O	1:A:209[A]:LEU:HD13	2.01	0.60
1:A:39:PRO:HA	1:A:42[A]:MET:HE3	1.83	0.59
1:A:28:TRP:CE3	2:A:564:BGC:O3	2.56	0.59
1:A:247[A]:ARG:NH2	7:A:890:HOH:O	2.35	0.59
1:A:100[A]:LYS:HD2	7:A:580:HOH:O	2.04	0.57
1:A:38[A]:ILE:HD11	1:A:116:TRP:HE3	1.71	0.56
1:A:270[B]:TRP:HH2	1:A:352:PRO:O	1.89	0.55
1:A:42[A]:MET:HE3	7:A:862:HOH:O	2.01	0.55
1:A:62:PRO:HG2	1:A:164:GLY:HA2	1.88	0.54
1:A:476:GLN:NE2	2:A:564:BGC:H6C1	2.19	0.54
1:A:35[B]:GLU:OE2	1:A:360:HIS:HD2	1.91	0.54
1:A:169[B]:GLU:OE2	1:A:175:LYS:CE	2.56	0.53
1:A:338:GLU:OE1	7:A:729:HOH:O	2.19	0.53
1:A:169[B]:GLU:OE2	1:A:175:LYS:NZ	2.42	0.53
1:A:318:MET:HG2	4:A:570:DIO:H22	1.91	0.52
1:A:129[B]:GLU:HG2	1:A:133:GLN:HE22	1.74	0.52
1:A:283[B]:ARG:HD2	1:A:295:GLN:CB	2.39	0.51
1:A:370:GLY:O	1:A:374[A]:GLU:HG3	2.11	0.50
1:A:490:ARG:HD3	1:A:494[C]:ARG:HD2	1.94	0.49
1:A:75:LYS:HE3	1:A:78:GLU:OE1	2.13	0.49
1:A:356[B]:GLU:OE1	7:A:865:HOH:O	2.20	0.49
1:A:11:THR:HG21	4:A:569:DIO:H2'2	1.96	0.48
2:A:565:BGC:H5	7:A:918:HOH:O	2.13	0.48
1:A:270[B]:TRP:CH2	1:A:352:PRO:O	2.67	0.47
1:A:247[B]:ARG:NH2	1:A:343:PHE:CD1	2.82	0.47
1:A:105[B]:MET:HE3	7:A:579:HOH:O	2.14	0.47
1:A:476:GLN:NE2	2:A:564:BGC:H6C2	2.27	0.47
1:A:432:GLU:OE2	1:A:464:GLU:OE2	2.32	0.47
1:A:483[A]:ARG:HG2	1:A:487:ILE:CD1	2.45	0.46
1:A:326:ARG:NE	7:A:916:HOH:O	2.47	0.46
1:A:14:PRO:HD2	2:A:564:BGC:H2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:312:TYR:OH	4:A:570:DIO:H2'2	2.15	0.45
1:A:483[B]:ARG:HG3	7:A:876:HOH:O	2.15	0.45
1:A:247[B]:ARG:HH12	1:A:398:HIS:CE1	2.28	0.44
1:A:270[B]:TRP:CH2	1:A:353:TYR:CD2	3.06	0.43
1:A:476:GLN:HE22	2:A:564:BGC:H6C2	1.71	0.43
1:A:490:ARG:HD2	1:A:491:ASP:OD1	2.18	0.43
1:A:39:PRO:CA	1:A:42[A]:MET:HE2	2.48	0.42
1:A:247[B]:ARG:NH1	1:A:398:HIS:HE1	2.10	0.42
1:A:129[B]:GLU:CG	1:A:133:GLN:HE22	2.31	0.42
1:A:342:LYS:HG2	1:A:343:PHE:CE2	2.54	0.42
1:A:407:TRP:CE2	3:A:567:GOL:H12	2.55	0.41
1:A:174:MET:CE	7:A:902:HOH:O	2.67	0.41
1:A:53:LYS:NZ	7:A:639:HOH:O	2.29	0.41
1:A:100[B]:LYS:HB2	1:A:100[B]:LYS:HE3	1.13	0.41
1:A:5:LEU:O	1:A:383:THR:HA	2.22	0.40
1:A:270[B]:TRP:CZ2	1:A:353:TYR:HD2	2.39	0.40
1:A:447:GLU:HG3	7:A:751:HOH:O	2.22	0.40
1:A:183:GLU:HG3	1:A:416:TRP:CZ2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	573/562 (102%)	560 (98%)	10 (2%)	3 (0%)	34 20

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	VAL
1	A	523	PHE
1	A	521	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	502/489 (103%)	488 (97%)	14 (3%)	51 39

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	LYS
1	A	127	LEU
1	A	175	LYS
1	A	240[A]	ASP
1	A	240[B]	ASP
1	A	302	LYS
1	A	303	GLU
1	A	396	GLU
1	A	407	TRP
1	A	422[A]	GLU
1	A	422[B]	GLU
1	A	494[B]	ARG
1	A	494[C]	ARG
1	A	495	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	268	GLN
1	A	360	HIS
1	A	378	GLN
1	A	398	HIS
1	A	476	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	A	563	-	12,12,12	0.89	0	17,17,17	1.79	6 (35%)
2	BGC	A	564	-	12,12,12	0.68	0	17,17,17	1.17	2 (11%)
2	BGC	A	565	-	12,12,12	0.56	0	17,17,17	1.16	2 (11%)
3	GOL	A	566	-	5,5,5	0.36	0	5,5,5	0.87	0
3	GOL	A	567	-	5,5,5	0.44	0	5,5,5	1.39	0
3	GOL	A	568	-	5,5,5	0.94	0	5,5,5	0.83	0
4	DIO	A	569	-	6,6,6	0.63	0	6,6,6	1.26	0
4	DIO	A	570	-	6,6,6	0.78	0	6,6,6	1.89	2 (33%)
5	PEG	A	571	-	6,6,6	0.29	0	5,5,5	0.88	0
6	PG4	A	572	-	12,12,12	0.56	0	11,11,11	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	563	-	-	0/2/22/22	0/1/1/1
2	BGC	A	564	-	-	0/2/22/22	0/1/1/1
2	BGC	A	565	-	-	0/2/22/22	0/1/1/1
3	GOL	A	566	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	567	-	-	0/4/4/4	0/0/0/0
3	GOL	A	568	-	-	0/4/4/4	0/0/0/0
4	DIO	A	569	-	-	0/0/6/6	0/1/1/1
4	DIO	A	570	-	-	0/0/6/6	0/1/1/1
5	PEG	A	571	-	-	0/4/4/4	0/0/0/0
6	PG4	A	572	-	-	0/10/10/10	0/0/0/0

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	563	BGC	O4-C4-C3	-2.68	104.30	110.34
2	A	565	BGC	C3-C4-C5	-2.34	106.11	110.20
2	A	564	BGC	O3-C3-C2	-2.18	105.44	110.34
2	A	565	BGC	C1-C2-C3	-2.02	107.42	110.43
2	A	563	BGC	O5-C5-C6	2.06	111.55	106.36
2	A	563	BGC	C3-C4-C5	2.15	113.95	110.20
2	A	563	BGC	O5-C5-C4	2.44	114.26	109.68
2	A	564	BGC	O5-C1-C2	2.74	114.17	109.80
4	A	570	DIO	C2'-O1'-C1'	2.92	119.73	109.89
4	A	570	DIO	C2-O1-C1	3.27	120.89	109.89
2	A	563	BGC	O5-C1-C2	3.61	115.56	109.80
2	A	563	BGC	C1-O5-C5	3.92	120.72	113.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	563	BGC	1	0
2	A	564	BGC	8	0
2	A	565	BGC	2	0
3	A	567	GOL	2	0
4	A	569	DIO	1	0
4	A	570	DIO	2	0
6	A	572	PG4	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	550/562 (97%)	-0.00	22 (4%)	42 44	13, 22, 38, 63	23 (4%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	304	VAL	12.7
1	A	305	GLY	6.6
1	A	22	TRP	6.0
1	A	270[A]	TRP	4.9
1	A	550	PRO	3.9
1	A	549	PRO	3.6
1	A	197	GLY	3.3
1	A	303	GLU	3.0
1	A	24	PHE	3.0
1	A	306	LEU	2.9
1	A	444	GLY	2.7
1	A	242	GLU	2.7
1	A	240[A]	ASP	2.5
1	A	462	ILE	2.3
1	A	394	SER	2.3
1	A	344	GLY	2.2
1	A	419	GLU	2.2
1	A	535[A]	ASN	2.1
1	A	23	PRO	2.1
1	A	64	LEU	2.1
1	A	148	LEU	2.1
1	A	422[A]	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	DIO	A	569	6/6	0.95	0.27	5.66	31,34,39,43	0
3	GOL	A	566	6/6	0.90	0.17	5.33	44,49,50,52	0
4	DIO	A	570	6/6	0.90	0.15	4.67	41,43,45,50	0
2	BGC	A	564	12/12	0.69	0.25	3.03	79,82,83,84	0
6	PG4	A	572	13/13	0.89	0.11	1.87	45,49,54,59	0
5	PEG	A	571	7/7	0.96	0.09	0.83	40,42,46,47	0
2	BGC	A	563	12/12	0.69	0.23	0.76	65,70,76,77	0
3	GOL	A	568	6/6	0.93	0.11	0.50	32,37,41,42	0
3	GOL	A	567	6/6	0.93	0.18	0.47	25,46,50,55	0
2	BGC	A	565	12/12	0.77	0.23	-	82,85,86,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.