



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 06:58 PM GMT

PDB ID : 4ND5
Title : Crystal structure of the lactate dehydrogenase from cryptosporidium parvum
Authors : Chattopadhyay, D.; Cook, W.J.
Deposited on : 2013-10-25
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 9099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactate dehydrogenase, adjacent gene encodes predicted malate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C 2240	N 1423	O 376	S 427	14	0	0
1	B	301	Total	C 2232	N 1419	O 375	S 424	14	0	0
1	C	300	Total	C 2224	N 1414	O 374	S 423	13	0	0
1	D	298	Total	C 2209	N 1403	O 372	S 421	13	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	ALA	VAL	SEE REMARK 999	UNP Q5CYZ2
B	202	ALA	VAL	SEE REMARK 999	UNP Q5CYZ2
C	202	ALA	VAL	SEE REMARK 999	UNP Q5CYZ2
D	202	ALA	VAL	SEE REMARK 999	UNP Q5CYZ2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	55	Total O 55 55	0	0
2	B	89	Total O 89 89	0	0
2	C	34	Total O 34 34	0	0
2	D	16	Total O 16 16	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lactate dehydrogenase, adjacent gene encodes predicted malate dehydrogenase




- Molecule 1: Lactate dehydrogenase, adjacent gene encodes predicted malate dehydrogenase



- Molecule 1: Lactate dehydrogenase, adjacent gene encodes predicted malate dehydrogenase



- Molecule 1: Lactate dehydrogenase, adjacent gene encodes predicted malate dehydrogenase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	136.49 Å 136.49 Å 170.22 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.88 – 2.10 19.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	91.3 (19.88-2.10) 91.4 (19.88-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.89 (at 2.09 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R , R_{free}	0.230 , 0.255 0.224 , 0.248	Depositor DCC
R_{free} test set	5057 reflections (5.52%)	DCC
Wilson B-factor (Å ²)	38.5	Xtriage
Anisotropy	0.500	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.3	EDS
Estimated twinning fraction	0.218 for -h,-k,l	Xtriage
L-test for twinning ²	$< L > = 0.42$, $< L^2 > = 0.24$	Xtriage
Outliers	0 of 100011 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9099	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.27	0/2275	0.47	0/3081
1	B	0.28	0/2267	0.48	0/3070
1	C	0.27	0/2259	0.45	0/3060
1	D	0.26	0/2244	0.43	0/3039
All	All	0.27	0/9045	0.46	0/12250

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	0	2277	12	0
1	B	2232	0	2273	6	0
1	C	2224	0	2264	4	0
1	D	2209	0	2244	4	0
2	A	55	0	0	0	0
2	B	89	0	0	0	0
2	C	34	0	0	0	0
2	D	16	0	0	0	0
All	All	9099	0	9058	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:ARG:NH2	1:B:75:THR:O	2.41	0.54
1:B:275:CYS:SG	1:B:278:HIS:HB2	2.49	0.52
1:C:39:ILE:HG23	1:D:42:LYS:HE3	1.92	0.51
1:B:142:LEU:HD22	1:B:164:ALA:HB2	1.93	0.51
1:C:42:LYS:HE3	1:D:39:ILE:HG23	1.92	0.50
1:A:142:LEU:HD22	1:A:164:ALA:HB2	1.96	0.48
1:A:137:CYS:HB2	1:A:146:VAL:HG22	1.97	0.46
1:A:275:CYS:SG	1:A:278:HIS:HB2	2.55	0.46
1:D:175:PHE:CE1	1:D:231:HIS:CE1	3.06	0.43
1:D:142:LEU:HD22	1:D:164:ALA:HB2	2.00	0.43
1:A:276:SER:O	1:A:282:LYS:HA	2.20	0.42
1:A:176:ILE:HD13	1:A:206:VAL:HG13	2.00	0.42
1:A:278:HIS:O	1:A:281:VAL:HG22	2.20	0.42
1:A:172:PHE:CD2	1:A:189:ALA:HB1	2.55	0.42
1:A:42:LYS:HE3	1:B:39:ILE:HG23	2.02	0.41
1:C:85:TYR:O	1:C:88:ILE:HG12	2.19	0.41
1:A:163:MET:SD	1:A:163:MET:C	2.99	0.41
1:B:142:LEU:CD2	1:B:164:ALA:HB2	2.51	0.41
1:A:162:GLY:O	1:A:271:CYS:HB2	2.21	0.41
1:A:216:ILE:HD11	1:A:225:ILE:CD1	2.51	0.41
1:C:142:LEU:HD22	1:C:164:ALA:HB2	2.02	0.41
1:A:39:ILE:HG23	1:B:42:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	298/321 (93%)	285 (96%)	13 (4%)	0	100 100
1	B	297/321 (92%)	286 (96%)	10 (3%)	1 (0%)	46 45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	296/321 (92%)	287 (97%)	8 (3%)	1 (0%)	46 45
1	D	294/321 (92%)	280 (95%)	13 (4%)	1 (0%)	46 45
All	All	1185/1284 (92%)	1138 (96%)	44 (4%)	3 (0%)	46 45

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	283	GLY
1	D	164	ALA
1	B	283	GLY

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	245/258 (95%)	244 (100%)	1 (0%)	93 96
1	B	244/258 (95%)	244 (100%)	0	100 100
1	C	243/258 (94%)	241 (99%)	2 (1%)	86 91
1	D	241/258 (93%)	240 (100%)	1 (0%)	93 96
All	All	973/1032 (94%)	969 (100%)	4 (0%)	93 96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	THR
1	C	195	HIS
1	C	197	ASP
1	D	197	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/321 (94%)	0.38	12 (3%) 42 51	30, 47, 77, 107	0
1	B	301/321 (93%)	0.30	13 (4%) 39 48	28, 40, 64, 98	0
1	C	300/321 (93%)	0.33	12 (4%) 42 51	37, 52, 72, 84	0
1	D	298/321 (92%)	0.79	37 (12%) 5 7	38, 64, 97, 113	0
All	All	1201/1284 (93%)	0.45	74 (6%) 24 32	28, 50, 86, 113	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	196	GLY	6.9
1	B	98	ALA	6.2
1	B	112	LEU	5.8
1	D	114	PHE	5.7
1	C	112	LEU	5.0
1	D	320	ASN	4.9
1	D	195	HIS	4.6
1	B	113	LEU	4.5
1	B	115	GLY	4.3
1	D	118	ARG	4.3
1	C	113	LEU	4.2
1	A	322	VAL	4.2
1	C	118	ARG	4.2
1	A	17	MET	4.1
1	D	280	GLY	4.1
1	D	54	ILE	4.0
1	D	113	LEU	3.9
1	C	197	ASP	3.9
1	A	329	LEU	3.8
1	B	114	PHE	3.7
1	D	304	GLU	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	277	ASN	3.6
1	A	277	ASN	3.6
1	B	327	LYS	3.4
1	C	328	VAL	3.2
1	C	54	ILE	3.1
1	B	195	HIS	2.9
1	A	327	LYS	2.9
1	B	118	ARG	2.8
1	D	322	VAL	2.7
1	C	56	GLU	2.7
1	D	197	ASP	2.7
1	D	112	LEU	2.7
1	D	313	LYS	2.7
1	D	234	ILE	2.6
1	A	323	ASN	2.6
1	B	56	GLU	2.6
1	A	330	ASP	2.6
1	A	282	LYS	2.5
1	D	318	SER	2.5
1	D	279	TYR	2.5
1	D	180	PHE	2.5
1	D	274	PHE	2.5
1	D	276	SER	2.5
1	A	314	LEU	2.5
1	D	151	LYS	2.5
1	C	195	HIS	2.4
1	D	307	LEU	2.4
1	D	321	GLU	2.4
1	D	327	LYS	2.4
1	B	119	ILE	2.4
1	A	113	LEU	2.4
1	B	197	ASP	2.3
1	D	121	ASP	2.3
1	D	306	ASP	2.3
1	D	236	TRP	2.2
1	D	314	LEU	2.2
1	B	328	VAL	2.2
1	D	285	ILE	2.2
1	C	116	ASN	2.2
1	D	281	VAL	2.2
1	C	314	LEU	2.2
1	C	329	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	139	THR	2.1
1	B	282	LYS	2.1
1	D	192	ILE	2.1
1	A	309	PRO	2.1
1	A	234	ILE	2.1
1	D	140	ASN	2.1
1	D	144	VAL	2.1
1	D	289	VAL	2.1
1	C	309	PRO	2.0
1	D	120	LEU	2.0
1	D	305	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.