



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:52 AM GMT

PDB ID : 3NED
Title : mRouge
Authors : Mayo, S.L.; Chica, R.A.; Moore, M.M.
Deposited on : 2010-06-08
Resolution : 0.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 0.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1319 (1.10-0.82)
Clashscore	102246	1031 (1.08-0.84)
Ramachandran outliers	100387	1331 (1.10-0.82)
Sidechain outliers	100360	1330 (1.10-0.82)
RSRZ outliers	91569	1324 (1.10-0.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	EYG	A	67[C]	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAmCherry1 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	11	45	0
			2140	1367	346	411	16			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	EXPRESSION TAG	UNP D1MPT3
A	-11	GLY	-	EXPRESSION TAG	UNP D1MPT3
A	-10	HIS	-	EXPRESSION TAG	UNP D1MPT3
A	-9	HIS	-	EXPRESSION TAG	UNP D1MPT3
A	-8	HIS	-	EXPRESSION TAG	UNP D1MPT3
A	-7	HIS	-	EXPRESSION TAG	UNP D1MPT3
A	-6	HIS	-	EXPRESSION TAG	UNP D1MPT3
A	-5	HIS	-	EXPRESSION TAG	UNP D1MPT3
A	-4	GLY	-	EXPRESSION TAG	UNP D1MPT3
A	16	THR	VAL	ENGINEERED MUTATION	UNP D1MPT3
A	26	GLU	VAL	ENGINEERED MUTATION	UNP D1MPT3
A	57	ALA	THR	ENGINEERED MUTATION	UNP D1MPT3
A	67	NRQ	MET	CHROMOPHORE	UNP D1MPT3
A	67	NRQ	TYR	CHROMOPHORE	UNP D1MPT3
A	67	NRQ	GLY	CHROMOPHORE	UNP D1MPT3
A	70	LYS	ASN	ENGINEERED MUTATION	UNP D1MPT3
A	83	LEU	PHE	ENGINEERED MUTATION	UNP D1MPT3
A	98	ASN	LYS	ENGINEERED MUTATION	UNP D1MPT3
A	146	CYS	LEU	ENGINEERED MUTATION	UNP D1MPT3
A	161	MET	VAL	ENGINEERED MUTATION	UNP D1MPT3
A	163	MET	PRO	ENGINEERED MUTATION	UNP D1MPT3
A	165	LEU	VAL	ENGINEERED MUTATION	UNP D1MPT3
A	195	THR	VAL	ENGINEERED MUTATION	UNP D1MPT3
A	197	THR	ARG	ENGINEERED MUTATION	UNP D1MPT3
A	217	ASN	ALA	ENGINEERED MUTATION	UNP D1MPT3

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Na	0	0
			3	3		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	414	Total	O	0	46
			470	470		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PAmCherry1 protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.81Å 42.76Å 61.20Å 90.00° 112.35° 90.00°	Depositor
Resolution (Å)	56.60 – 0.95 56.61 – 0.95	Depositor EDS
% Data completeness (in resolution range)	91.9 (56.60-0.95) 92.0 (56.61-0.95)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 0.95Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.133 , 0.157 0.135 , 0.159	Depositor DCC
R_{free} test set	6732 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	5.8	Xtriage
Anisotropy	1.569	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.0	EDS
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 134996 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2617	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NRQ, CH6, EYG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.39	21/2253 (0.9%)	1.19	8/3021 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65[A]	PHE	CA-C	13.57	1.88	1.52
1	A	65[B]	PHE	CA-C	13.57	1.88	1.52
1	A	65[C]	PHE	CA-C	13.57	1.88	1.52
1	A	146[A]	CYS	CB-SG	8.09	1.96	1.82
1	A	146[B]	CYS	CB-SG	8.09	1.96	1.82
1	A	231	LYS	CE-NZ	7.91	1.68	1.49
1	A	89[A]	GLU	CD-OE1	6.44	1.32	1.25
1	A	89[B]	GLU	CD-OE1	6.44	1.32	1.25
1	A	147[A]	SER	CB-OG	-6.20	1.34	1.42
1	A	147[C]	SER	CB-OG	-6.20	1.34	1.42
1	A	4	ASN	CG-ND2	5.78	1.47	1.32
1	A	121[A]	LYS	CD-CE	5.53	1.65	1.51
1	A	121[B]	LYS	CD-CE	5.53	1.65	1.51
1	A	125[A]	ARG	CG-CD	5.30	1.65	1.51
1	A	125[B]	ARG	CG-CD	5.30	1.65	1.51
1	A	10	GLU	CD-OE1	-5.11	1.20	1.25
1	A	28	GLU	CD-OE1	-5.07	1.20	1.25
1	A	89[A]	GLU	CG-CD	5.04	1.59	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	89[B]	GLU	CG-CD	5.04	1.59	1.51
1	A	128[A]	ASN	C-O	5.03	1.32	1.23
1	A	128[B]	ASN	C-O	5.03	1.32	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121[A]	LYS	CD-CE-NZ	10.55	135.97	111.70
1	A	121[B]	LYS	CD-CE-NZ	10.55	135.97	111.70
1	A	220	ARG	NE-CZ-NH1	8.31	124.46	120.30
1	A	36	ARG	NE-CZ-NH1	8.30	124.45	120.30
1	A	92	LYS	CD-CE-NZ	-7.57	94.29	111.70
1	A	174	ASP	CB-CG-OD2	-6.32	112.61	118.30
1	A	139	LYS	CB-CG-CD	5.15	124.98	111.60
1	A	36	ARG	NE-CZ-NH2	-5.00	117.80	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	67[C]	EYG	CA3

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2140	0	2063	83	0
2	A	3	0	0	0	0
3	A	4	0	3	0	0
4	A	470	0	0	42	0
All	All	2617	0	2066	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:LYS:HD2	4:A:425:HOH:O	1.31	1.29
1:A:114[A]:GLN:HG2	4:A:365:HOH:O	1.17	1.27
1:A:128[B]:ASN:ND2	4:A:261:HOH:O	1.69	1.22
1:A:161:MET:SD	4:A:420[A]:HOH:O	1.99	1.18
1:A:141[A]:MET:SD	1:A:168:LYS:HG2	1.85	1.17
1:A:139:LYS:HE2	4:A:522[B]:HOH:O	1.47	1.12
1:A:26:GLU:HG3	4:A:601[B]:HOH:O	1.52	1.09
1:A:70[B]:LYS:HE2	1:A:181[B]:TYR:OH	1.52	1.08
1:A:161:MET:CB	4:A:420[A]:HOH:O	2.03	1.05
1:A:141[B]:MET:HE3	1:A:168:LYS:HG2	1.43	0.98
1:A:67[A]:NRQ:HE2	1:A:146[A]:CYS:SG	2.09	0.92
1:A:180[B]:THR:OG1	4:A:370:HOH:O	1.87	0.89
1:A:67[A]:NRQ:OH	1:A:146[A]:CYS:SG	2.32	0.87
1:A:26:GLU:CG	4:A:601[B]:HOH:O	2.13	0.86
1:A:151[B]:TYR:OH	4:A:593:HOH:O	1.95	0.85
1:A:218[B]:GLU:OE2	1:A:220:ARG:CZ	2.25	0.85
1:A:17:HIS:CE1	1:A:19[B]:GLU:HG3	2.13	0.84
1:A:139:LYS:CE	4:A:522[B]:HOH:O	2.12	0.83
1:A:141[B]:MET:CE	1:A:168:LYS:HG2	2.08	0.82
1:A:198:LYS:HG3	4:A:423[B]:HOH:O	1.80	0.82
1:A:114[A]:GLN:CG	4:A:365:HOH:O	1.90	0.81
1:A:164[B]:ARG:NH1	4:A:421:HOH:O	2.06	0.80
1:A:5:MET:HE3	4:A:383:HOH:O	1.82	0.80
1:A:168:LYS:CD	4:A:425:HOH:O	2.06	0.78
1:A:67[A]:NRQ:CE2	1:A:146[A]:CYS:SG	2.74	0.75
1:A:141[A]:MET:SD	1:A:168:LYS:CG	2.71	0.75
1:A:114[A]:GLN:CD	4:A:365:HOH:O	2.19	0.74
1:A:132:ASP:OD1	4:A:402:HOH:O	2.06	0.74
1:A:70[B]:LYS:HE2	1:A:181[B]:TYR:HH	1.51	0.73
1:A:17:HIS:HE1	1:A:19[B]:GLU:HG3	1.51	0.72
1:A:121[A]:LYS:HE2	4:A:505:HOH:O	1.92	0.70
1:A:54:LEU:HD12	4:A:468:HOH:O	1.91	0.70
1:A:161:MET:HB3	4:A:420[A]:HOH:O	1.74	0.69
1:A:70[B]:LYS:HE2	1:A:181[B]:TYR:CZ	2.28	0.69
1:A:70[B]:LYS:HB3	1:A:83:LEU:CD1	2.24	0.68
1:A:141[B]:MET:SD	4:A:390[B]:HOH:O	2.50	0.68
1:A:164[B]:ARG:NE	4:A:421:HOH:O	2.22	0.68
1:A:70[B]:LYS:HB2	4:A:591:HOH:O	1.95	0.66
1:A:45[B]:LYS:NZ	4:A:577:HOH:O	2.28	0.66
1:A:164[B]:ARG:CZ	4:A:421:HOH:O	2.42	0.65
1:A:153:GLU:OE2	4:A:398[B]:HOH:O	2.14	0.65
1:A:161:MET:HB2	4:A:420[A]:HOH:O	1.80	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141[A]:MET:CG	1:A:168:LYS:HG2	2.25	0.64
1:A:70[B]:LYS:HG3	1:A:181[B]:TYR:CE1	2.32	0.64
1:A:67[A]:NRQ:CZ	1:A:146[A]:CYS:SG	2.86	0.63
1:A:228:GLU:HG3	4:A:609:HOH:O	1.99	0.62
1:A:26:GLU:CD	4:A:601[B]:HOH:O	2.33	0.62
1:A:70[B]:LYS:HB3	1:A:83:LEU:HD12	1.82	0.61
1:A:140:THR:HB	1:A:165[B]:LEU:HD22	1.83	0.60
1:A:141[B]:MET:HE3	1:A:168:LYS:CG	2.27	0.59
1:A:70[B]:LYS:CB	4:A:591:HOH:O	2.52	0.58
1:A:70[B]:LYS:CG	4:A:591:HOH:O	2.54	0.56
1:A:108[B]:THR:CG2	4:A:272[B]:HOH:O	2.56	0.54
1:A:70[B]:LYS:HG3	1:A:181[B]:TYR:CZ	2.44	0.53
1:A:67[A]:NRQ:HD2	4:A:497[A]:HOH:O	2.09	0.52
1:A:141[A]:MET:CG	1:A:168:LYS:CG	2.87	0.52
1:A:218[B]:GLU:OE2	1:A:220:ARG:NH2	2.43	0.52
1:A:112[A]:SER:HB2	4:A:580:HOH:O	2.10	0.51
1:A:108[B]:THR:HG22	4:A:272[B]:HOH:O	2.11	0.51
1:A:95:ARG:HB3	1:A:107[B]:VAL:HG13	1.93	0.51
1:A:70[A]:LYS:HE3	1:A:148:GLU:OE1	2.11	0.50
1:A:59:ASP:HB3	1:A:165[B]:LEU:HD21	1.93	0.50
1:A:70[B]:LYS:HB3	1:A:83:LEU:HD13	1.92	0.49
1:A:141[A]:MET:HG2	1:A:168:LYS:HG3	1.95	0.47
1:A:182[A]:LYS:HE3	4:A:320:HOH:O	2.13	0.47
1:A:141[A]:MET:HG2	1:A:168:LYS:CG	2.45	0.47
1:A:121[A]:LYS:NZ	4:A:386:HOH:O	2.29	0.47
1:A:65[A]:PHE:C	1:A:67[A]:NRQ:HG12	2.36	0.47
1:A:17:HIS:HE1	1:A:19[B]:GLU:CG	2.25	0.47
1:A:125[B]:ARG:HD2	4:A:504:HOH:O	2.16	0.45
1:A:141[B]:MET:HB3	1:A:141[B]:MET:HE3	1.74	0.45
1:A:143:TRP:CZ3	1:A:165[B]:LEU:HG	2.53	0.44
1:A:95:ARG:NH2	4:A:415:HOH:O	2.51	0.43
1:A:166:LYS:NZ	4:A:422[C]:HOH:O	2.48	0.43
1:A:1:GLU:HG2	1:A:2:GLU:N	2.35	0.42
1:A:65[A]:PHE:C	1:A:67[A]:NRQ:CG1	2.87	0.42
1:A:67[A]:NRQ:N1	1:A:67[A]:NRQ:CA3	2.82	0.42
1:A:121[A]:LYS:CD	4:A:268:HOH:O	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/244 (110%)	261 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/205 (117%)	238 (100%)	1 (0%)	93	72

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	231	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CH6	A	67[B]	1	24,24,25	1.62	3 (12%)	28,32,34	2.89	10 (35%)
1	EYG	A	67[C]	1	19,23,26	3.16	5 (26%)	22,33,37	2.97	6 (27%)
1	NRQ	A	67[A]	1	23,24,25	1.47	3 (13%)	23,32,34	2.73	9 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CH6	A	67[B]	1	-	0/11/31/32	0/2/2/2
1	EYG	A	67[C]	1	1/1/5/8	0/6/34/37	0/1/2/2
1	NRQ	A	67[A]	1	-	0/9/31/32	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	67[C]	EYG	CA3-N1	-11.37	1.24	1.46
1	A	67[C]	EYG	CAR-C2	-4.25	1.44	1.49
1	A	67[B]	CH6	CA2-C2	-3.88	1.44	1.48
1	A	67[A]	NRQ	CA2-C2	-3.88	1.44	1.48
1	A	67[C]	EYG	CB-CA3	-3.73	1.50	1.53
1	A	67[A]	NRQ	C1-N2	2.19	1.37	1.33
1	A	67[C]	EYG	O13-C13	2.63	1.42	1.38
1	A	67[C]	EYG	CD1-CG2	3.14	1.45	1.39
1	A	67[B]	CH6	CD1-CG2	3.14	1.45	1.39
1	A	67[A]	NRQ	CD1-CG2	3.14	1.45	1.39
1	A	67[B]	CH6	C1-N2	3.92	1.38	1.32

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67[B]	CH6	CE-SD-CG1	-6.10	79.56	100.37
1	A	67[C]	EYG	O2-C2-N3	-3.40	122.27	125.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67[A]	NRQ	C2-CA2-N2	-3.38	106.22	108.91
1	A	67[A]	NRQ	CB1-CG1-SD	-3.29	104.81	112.88
1	A	67[A]	NRQ	N3-C1-N2	-3.27	109.55	113.26
1	A	67[B]	CH6	O2-C2-CA2	-2.22	129.75	130.95
1	A	67[A]	NRQ	O2-C2-CA2	-2.22	129.75	130.95
1	A	67[C]	EYG	CD2-CG2-CB2	-2.12	113.98	121.23
1	A	67[B]	CH6	CD2-CG2-CB2	-2.12	113.98	121.23
1	A	67[A]	NRQ	CD2-CG2-CB2	-2.12	113.98	121.23
1	A	67[B]	CH6	CA1-C1-N2	2.09	127.90	123.38
1	A	67[B]	CH6	CG2-CB2-CA2	2.13	132.98	130.22
1	A	67[A]	NRQ	CG2-CB2-CA2	2.13	132.98	130.22
1	A	67[A]	NRQ	CB2-CA2-N2	2.25	132.70	128.67
1	A	67[B]	CH6	CA3-N3-C1	2.49	130.24	127.36
1	A	67[B]	CH6	CB1-CG1-SD	2.73	124.69	113.17
1	A	67[C]	EYG	C-CAG-N3	3.49	120.91	112.22
1	A	67[B]	CH6	C3-CA3-N3	3.61	120.91	113.00
1	A	67[A]	NRQ	C3-CA3-N3	3.61	120.91	113.00
1	A	67[C]	EYG	CAR-C2-N3	4.80	107.95	105.56
1	A	67[C]	EYG	CB-CA3-N1	5.11	122.91	108.81
1	A	67[B]	CH6	CB1-CA1-N1	6.99	124.46	109.81
1	A	67[B]	CH6	CA2-C2-N3	9.08	107.95	103.40
1	A	67[A]	NRQ	CA2-C2-N3	9.08	107.95	103.40
1	A	67[C]	EYG	C2-CAR-N23	9.69	113.06	105.13

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	67[C]	EYG	CA3

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	67[A]	NRQ	8	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	233	-	1,3,3	0.08	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	233	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	228/244 (93%)	-0.02	8 (3%) 48 36	5, 8, 16, 31	3 (1%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ASP	5.8
1	A	230	TYR	4.8
1	A	2	GLU	4.6
1	A	229	LEU	4.0
1	A	227	ASP	3.2
1	A	231	LYS	2.8
1	A	4	ASN	2.5
1	A	226[A]	MET	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	EYG	A	67[C]	22/25	0.97	0.17	-	6,14,28,34	22
1	CH6	A	67[B]	23/24	0.97	0.16	-	6,9,28,34	23
1	NRQ	A	67[A]	23/24	0.98	0.15	-	6,10,28,34	23

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ACT	A	233	4/4	0.98	0.06	1.51	8,8,8,9	0
2	NA	A	232	1/1	0.99	0.09	1.14	8,8,8,8	0
2	NA	A	235	1/1	1.00	0.05	-1.97	8,8,8,8	0
2	NA	A	234	1/1	1.00	0.03	-5.11	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.