



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:56 AM GMT

PDB ID : 3NLM
Title : Structure of neuronal nitric oxide synthase heme domain in complex with 6-
{(3'R,4'R)-3'-[2''-(3'''-fluorophenethylamino)ethoxy]pyrrolidin-4'-yl}methyl}-
4-methylpyridin-2-amine
Authors : Li, H.; Delker, S.L.; Poulos, T.L.
Deposited on : 2010-06-21
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	860	-	-	-	X
5	ACT	B	860	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	2	0
			3343	2139	572	610	22			
1	B	411	Total	C	N	O	S	0	3	0
			3359	2149	575	613	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 6-{{[(3R,4R)-4-(2-{[2-(3-FLUOROPHENYL)ETHYL]AMINO}ETHOXY)PYRROLIDIN-3-YL]METHYL}-4-METHYLPYRIDIN-2-AMINE (three-letter code: JRR) (formula: $C_{21}H_{29}FN_4O$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			27	21	1	4	1		
3	B	1	Total	C	F	N	O	0	0
			27	21	1	4	1		

- Molecule 4 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			17	9	5	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	156	Total	O	0	0
			156	156		
7	B	225	Total	O	0	0
			225	225		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.05Å 111.12Å 164.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.11 – 1.85 37.01 – 1.83	Depositor EDS
% Data completeness (in resolution range)	94.3 (38.11-1.85) 94.3 (37.01-1.83)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.189 , 0.222 0.202 , 0.236	Depositor DCC
R_{free} test set	3880 reflections (5.17%)	DCC
Wilson B-factor (Å ²)	36.9	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 80275 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7266	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, ACT, H4B, JRR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/3439	0.64	0/4665
1	B	0.69	0/3461	0.67	1/4692 (0.0%)
All	All	0.67	0/6900	0.66	1/9357 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	570	MET	CA-CB-CG	-5.88	103.30	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3343	0	3253	29	0
1	B	3359	0	3276	16	0
2	A	43	0	30	1	0
2	B	43	0	30	2	0
3	A	27	0	29	2	0
3	B	27	0	29	2	0
4	A	17	0	15	0	0
4	B	17	0	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0
7	A	156	0	0	1	0
7	B	225	0	0	2	0
All	All	7266	0	6683	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:750:HEM:HBA2	3:A:800:JRR:H4	1.59	0.85
2:B:750:HEM:HBA2	3:B:800:JRR:H4	1.58	0.85
1:A:371:ARG:CG	1:A:371:ARG:HH21	1.95	0.78
1:A:371:ARG:HH21	1:A:371:ARG:HG2	1.53	0.73
1:A:371:ARG:HG2	1:A:371:ARG:NH2	2.03	0.72
1:B:299:ARG:HG2	1:B:317:HIS:NE2	2.07	0.70
1:B:299:ARG:HB3	1:B:318:LEU:HD21	1.79	0.65
1:A:628:GLN:HG2	1:B:631:VAL:HG11	1.79	0.64
1:A:321:THR:HG23	1:A:322:LEU:HG	1.87	0.56
1:A:569:ASN:O	1:A:707:GLN:HG2	2.07	0.54
1:B:717:LYS:NZ	7:B:1138:HOH:O	2.43	0.52
1:A:349:ARG:HA	1:A:353:GLN:NE2	2.24	0.51
1:B:403:TYR:CE1	1:B:407:HIS:CE1	2.99	0.51
1:A:380:ARG:HD3	1:A:400:GLU:OE1	2.12	0.50
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.94	0.50
1:A:569:ASN:HD22	1:A:569:ASN:H	1.58	0.50
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.92	0.49
1:B:300:PHE:HD2	1:B:313:THR:HG21	1.77	0.49
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.48	0.49
1:A:696:LEU:HB3	1:B:330:ILE:HD11	1.95	0.48
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.02	0.48
1:A:330:ILE:HD11	1:B:696:LEU:HD22	1.97	0.47
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.98	0.46
2:B:750:HEM:CBA	3:B:800:JRR:H4	2.38	0.46
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.98	0.45
1:A:335:ILE:CD1	1:B:694:GLU:HB3	2.47	0.45
1:A:299:ARG:HG3	1:A:318:LEU:HD21	1.98	0.45
1:A:524:LEU:O	1:A:531:PRO:HA	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:ARG:HG2	1:B:317:HIS:CE1	2.53	0.43
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.53	0.43
1:A:588:TYR:CD1	1:A:593:ILE:HD11	2.53	0.43
1:A:371:ARG:HH21	1:A:371:ARG:HG3	1.81	0.42
1:A:331:CYS:HB3	1:B:697:ASN:HB3	2.01	0.42
1:A:349:ARG:HA	1:A:353:GLN:HE22	1.84	0.42
1:A:335:ILE:HD12	1:B:694:GLU:HB3	2.02	0.42
1:A:436:HIS:ND1	7:A:1147:HOH:O	2.37	0.42
1:A:592[A]:GLU:OE1	3:A:800:JRR:H16	2.20	0.42
1:A:609:GLU:O	1:A:613:LYS:HG2	2.20	0.41
1:B:364:GLN:NE2	7:B:1214:HOH:O	2.52	0.41
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.55	0.41
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.51	0.40
1:A:317:HIS:O	1:A:320:SER:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/422 (96%)	400 (98%)	7 (2%)	0	100	100
1	B	410/422 (97%)	405 (99%)	5 (1%)	0	100	100
All	All	817/844 (97%)	805 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/377 (97%)	355 (97%)	12 (3%)	45	25
1	B	369/377 (98%)	361 (98%)	8 (2%)	60	43
All	All	736/754 (98%)	716 (97%)	20 (3%)	52	34

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	348	VAL
1	A	350	THR
1	A	352	ASP
1	A	353	GLN
1	A	371	ARG
1	A	380	ARG
1	A	389	GLU
1	A	489	ASP
1	A	547	ARG
1	A	569	ASN
1	A	645	LYS
1	A	705	GLU
1	B	353	GLN
1	B	382	GLU
1	B	389	GLU
1	B	390	SER
1	B	454	ASN
1	B	547	ARG
1	B	627	ASP
1	B	697	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	364	GLN
1	A	454	ASN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	628	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	697	ASN
1	B	364	GLN
1	B	385	ASN
1	B	454	ASN
1	B	507	GLN
1	B	535	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	750	1	30,50,50	2.21	7 (23%)	24,82,82	2.86	12 (50%)
4	H4B	A	760	-	13,18,18	0.81	0	11,26,26	2.43	6 (54%)
3	JRR	A	800	-	26,29,29	0.65	0	27,38,38	1.83	8 (29%)
5	ACT	A	860	-	1,3,3	1.24	0	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	750	1	30,50,50	2.17	6 (20%)	24,82,82	2.92	11 (45%)
4	H4B	B	760	-	13,18,18	1.10	1 (7%)	11,26,26	2.59	4 (36%)
3	JRR	B	800	-	26,29,29	0.74	0	27,38,38	1.83	8 (29%)
5	ACT	B	860	-	1,3,3	1.55	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8
4	H4B	A	760	-	-	0/8/17/17	0/2/2/2
3	JRR	A	800	-	-	0/13/23/23	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
4	H4B	B	760	-	-	0/8/17/17	0/2/2/2
3	JRR	B	800	-	-	0/13/23/23	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C3B-C4B	-6.85	1.45	1.51
2	A	750	HEM	C3B-C4B	-6.75	1.45	1.51
2	A	750	HEM	C3D-C4D	-5.57	1.44	1.51
2	B	750	HEM	C3D-C4D	-4.60	1.45	1.51
2	A	750	HEM	C2C-C1C	-3.64	1.45	1.52
2	B	750	HEM	C2C-C1C	-3.64	1.45	1.52
2	A	750	HEM	C2D-C1D	-2.40	1.44	1.51
2	B	750	HEM	C2B-C1B	-2.29	1.44	1.51
2	A	750	HEM	C2B-C1B	-2.02	1.45	1.51
4	B	760	H4B	C7-N8	2.03	1.48	1.46
2	A	750	HEM	CMA-C3A	2.11	1.56	1.51
2	B	750	HEM	CMA-C3A	2.51	1.56	1.51
2	B	750	HEM	FE-NC	3.94	2.11	1.95
2	A	750	HEM	FE-NC	4.52	2.13	1.95

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	CBA-CAA-C2A	-6.69	100.54	112.53
2	A	750	HEM	CBA-CAA-C2A	-6.36	101.13	112.53
2	B	750	HEM	CAA-C2A-C1A	-5.06	121.52	127.01
2	A	750	HEM	CBD-CAD-C3D	-4.47	100.54	113.55
3	A	800	JRR	C3-C4-C11	-3.48	105.61	112.83
2	B	750	HEM	CBD-CAD-C3D	-3.45	103.53	113.55
4	A	760	H4B	N3-C2-N1	-3.35	120.03	125.53
3	A	800	JRR	C14-C13-C12	-3.16	119.28	123.35
3	B	800	JRR	C14-C13-C12	-2.93	119.57	123.35
3	B	800	JRR	C3-C4-C11	-2.91	106.79	112.83
2	A	750	HEM	CAA-C2A-C1A	-2.90	123.86	127.01
4	B	760	H4B	N3-C2-N1	-2.84	120.88	125.53
3	B	800	JRR	C4A-C3A-C2A	-2.83	118.51	120.28
3	A	800	JRR	C4A-C3A-C2A	-2.75	118.56	120.28
2	A	750	HEM	CMA-C3A-C4A	-2.58	124.10	128.36
3	B	800	JRR	C3A-C2A-N1A	-2.29	120.38	122.96
3	A	800	JRR	C3A-C2A-N1A	-2.27	120.40	122.96
2	B	750	HEM	C3B-C4B-NB	-2.02	107.77	111.63
2	A	750	HEM	C3B-C4B-NB	-2.00	107.80	111.63
3	B	800	JRR	N6A-C6A-N1A	2.03	120.20	116.50
3	B	800	JRR	F13-C13-C14	2.06	121.96	118.52
3	A	800	JRR	F13-C13-C14	2.08	122.00	118.52
4	A	760	H4B	N2-C2-N1	2.10	120.67	117.20
3	A	800	JRR	N6A-C6A-N1A	2.10	120.33	116.50
4	B	760	H4B	C2-N1-C8A	2.17	119.41	114.54
2	B	750	HEM	C2D-C3D-C4D	2.24	105.30	101.50
4	A	760	H4B	C4A-C8A-N8	2.33	121.18	118.43
2	B	750	HEM	CMD-C2D-C3D	2.44	125.14	114.35
2	B	750	HEM	C2C-C1C-CHC	2.49	127.47	123.68
2	A	750	HEM	C2D-C3D-C4D	2.52	105.77	101.50
2	A	750	HEM	C3B-C4B-CHC	2.55	126.75	123.16
3	A	800	JRR	C3-N2-C2	2.82	123.35	113.35
2	A	750	HEM	CMD-C2D-C3D	2.96	127.44	114.35
4	A	760	H4B	C2-N1-C8A	3.04	121.38	114.54
4	A	760	H4B	C4-N3-C2	3.42	120.68	115.94
2	A	750	HEM	CMB-C2B-C3B	3.55	125.40	116.53
3	B	800	JRR	C11-C12-C13	3.55	121.79	118.84
2	B	750	HEM	CMC-C2C-C3C	4.04	126.63	116.53
4	B	760	H4B	C4-N3-C2	4.10	121.63	115.94
2	A	750	HEM	CAD-C3D-C4D	4.11	126.96	112.47
2	B	750	HEM	CAD-C3D-C2D	4.21	125.31	113.22
4	A	760	H4B	C4-C4A-C8A	4.52	118.65	114.56
2	B	750	HEM	CAD-C3D-C4D	4.80	129.38	112.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CAD-C3D-C2D	4.86	127.19	113.22
2	A	750	HEM	CMC-C2C-C3C	4.89	128.73	116.53
3	A	800	JRR	C6A-N1A-C2A	5.03	121.81	118.23
2	B	750	HEM	CMB-C2B-C3B	5.20	129.52	116.53
3	B	800	JRR	C6A-N1A-C2A	5.44	122.10	118.23
4	B	760	H4B	C4-C4A-C8A	5.88	119.89	114.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	1	0
3	A	800	JRR	2	0
2	B	750	HEM	2	0
3	B	800	JRR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	409/422 (96%)	1.04	87 (21%)	1 1	28, 47, 76, 92	0
1	B	411/422 (97%)	0.48	40 (9%)	10 9	26, 39, 64, 83	0
All	All	820/844 (97%)	0.76	127 (15%)	3 3	26, 43, 71, 92	0

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	9.7
1	A	488	PRO	8.5
1	A	348	VAL	7.5
1	A	486	LYS	6.9
1	B	348	VAL	6.3
1	A	373	GLY	6.1
1	A	352	ASP	5.8
1	A	300	PHE	5.7
1	A	716	TRP	5.6
1	A	355	PHE	5.3
1	A	299	ARG	4.8
1	A	351	LYS	4.7
1	B	715	VAL	4.7
1	A	491	SER	4.6
1	A	504	ILE	4.6
1	A	619	ARG	4.5
1	B	619	ARG	4.5
1	A	388	ILE	4.4
1	A	713	THR	4.3
1	A	715	VAL	4.1
1	A	490	GLY	4.0
1	B	299	ARG	4.0
1	A	507	GLN	3.9
1	A	492	THR	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	392	SER	3.9
1	B	350	THR	3.9
1	A	386	LYS	3.9
1	A	503	GLU	3.9
1	A	567	VAL	3.8
1	A	350	THR	3.7
1	B	718	GLY	3.6
1	A	680	VAL	3.6
1	A	479	LEU	3.5
1	A	390	SER	3.5
1	A	487	GLN	3.3
1	A	480	ILE	3.3
1	A	389	GLU	3.3
1	A	511	LYS	3.3
1	B	667	ARG	3.2
1	A	375	LYS	3.2
1	B	321	THR	3.2
1	A	667	ARG	3.1
1	A	678	TRP	3.1
1	B	351	LYS	3.1
1	B	566	ALA	3.0
1	A	592[A]	GLU	3.0
1	A	385	ASN	3.0
1	B	591	THR	3.0
1	B	677	VAL	2.9
1	A	566	ALA	2.8
1	A	704	PHE	2.8
1	B	713	THR	2.8
1	B	682	PRO	2.8
1	A	686	SER	2.7
1	B	572	LEU	2.7
1	A	415	CYS	2.7
1	A	677	VAL	2.7
1	B	691	PHE	2.7
1	A	682	PRO	2.7
1	B	680	VAL	2.6
1	B	592[A]	GLU	2.6
1	A	595	VAL	2.6
1	B	310	VAL	2.6
1	A	565	PRO	2.6
1	A	469	LYS	2.6
1	A	506	ILE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	552	ASP	2.6
1	A	353	GLN	2.6
1	A	391	THR	2.6
1	A	393	THR	2.6
1	A	302	LYS	2.6
1	A	514	ARG	2.6
1	B	683	MET	2.5
1	A	356	PRO	2.5
1	B	595	VAL	2.5
1	A	382	GLU	2.5
1	A	442	ILE	2.5
1	B	679	ILE	2.5
1	A	489	ASP	2.5
1	A	570	MET	2.4
1	B	352	ASP	2.4
1	A	583	PRO	2.4
1	A	584	PHE	2.4
1	B	328	GLU	2.4
1	A	591	THR	2.4
1	A	381	LEU	2.4
1	B	479	LEU	2.4
1	B	681	PRO	2.4
1	A	349	ARG	2.4
1	A	679	ILE	2.4
1	A	467	ASP	2.4
1	A	620	LYS	2.4
1	B	355	PHE	2.4
1	A	323	GLU	2.3
1	B	615	ASP	2.3
1	A	681	PRO	2.3
1	A	354	LEU	2.3
1	A	588	TYR	2.3
1	A	676	TRP	2.3
1	A	691	PHE	2.3
1	A	594	GLY	2.3
1	A	505	CYS	2.3
1	A	568	SER	2.3
1	A	485	TYR	2.3
1	B	561	TRP	2.2
1	A	714	HIS	2.2
1	B	302	LYS	2.2
1	A	371	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	676	TRP	2.2
1	B	620	LYS	2.2
1	A	572	LEU	2.2
1	A	508	GLN	2.2
1	B	630	LEU	2.2
1	A	712	ASN	2.1
1	A	330	ILE	2.1
1	B	301	LEU	2.1
1	A	416	VAL	2.1
1	B	593	ILE	2.1
1	B	567	VAL	2.1
1	A	617	ASP	2.1
1	A	556	ASP	2.1
1	B	616	LEU	2.1
1	B	685	GLY	2.1
1	A	478	GLN	2.1
1	B	704	PHE	2.0
1	A	683	MET	2.0
1	A	685	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	ACT	A	860	4/4	0.95	0.20	6.54	50,51,52,53	0
5	ACT	B	860	4/4	0.96	0.18	4.75	38,40,42,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	H4B	B	760	17/17	0.97	0.18	0.85	27,30,34,35	0
4	H4B	A	760	17/17	0.95	0.18	0.65	28,32,35,37	0
2	HEM	B	750	43/43	0.98	0.19	0.48	24,29,37,42	0
2	HEM	A	750	43/43	0.97	0.21	0.39	27,31,38,41	0
3	JRR	B	800	27/27	0.94	0.19	0.22	33,37,46,48	0
3	JRR	A	800	27/27	0.89	0.21	0.19	36,43,47,47	0
6	ZN	A	900	1/1	1.00	0.07	-1.29	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.