



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:03 PM GMT

PDB ID : 4NLE
Title : Crystal structure of apo Adenylosuccinate Lyase from Mycobacterium smegmatis
Authors : Banerjee, S.; Murthy, M.R.N.
Deposited on : 2013-11-14
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	<div> <div>9%</div> <div> <div></div> <div>90%</div> <div>• 5%</div> </div> </div>
1	B	488	<div> <div>7%</div> <div> <div></div> <div>90%</div> <div>• 8%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylosuccinate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	464	Total	C	N	O	S	0	3	0
			3542	2220	637	671	14			
1	B	451	Total	C	N	O	S	0	3	0
			3428	2149	611	656	12			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	EXPRESSION TAG	UNP A0R4I6
A	-12	ARG	-	EXPRESSION TAG	UNP A0R4I6
A	-11	GLY	-	EXPRESSION TAG	UNP A0R4I6
A	-10	SER	-	EXPRESSION TAG	UNP A0R4I6
A	-9	HIS	-	EXPRESSION TAG	UNP A0R4I6
A	-8	HIS	-	EXPRESSION TAG	UNP A0R4I6
A	-7	HIS	-	EXPRESSION TAG	UNP A0R4I6
A	-6	HIS	-	EXPRESSION TAG	UNP A0R4I6
A	-5	HIS	-	EXPRESSION TAG	UNP A0R4I6
A	-4	HIS	-	EXPRESSION TAG	UNP A0R4I6
A	-3	GLY	-	EXPRESSION TAG	UNP A0R4I6
A	-2	MET	-	EXPRESSION TAG	UNP A0R4I6
A	-1	ALA	-	EXPRESSION TAG	UNP A0R4I6
A	0	SER	-	EXPRESSION TAG	UNP A0R4I6
B	-13	MET	-	EXPRESSION TAG	UNP A0R4I6
B	-12	ARG	-	EXPRESSION TAG	UNP A0R4I6
B	-11	GLY	-	EXPRESSION TAG	UNP A0R4I6
B	-10	SER	-	EXPRESSION TAG	UNP A0R4I6
B	-9	HIS	-	EXPRESSION TAG	UNP A0R4I6
B	-8	HIS	-	EXPRESSION TAG	UNP A0R4I6
B	-7	HIS	-	EXPRESSION TAG	UNP A0R4I6
B	-6	HIS	-	EXPRESSION TAG	UNP A0R4I6
B	-5	HIS	-	EXPRESSION TAG	UNP A0R4I6
B	-4	HIS	-	EXPRESSION TAG	UNP A0R4I6
B	-3	GLY	-	EXPRESSION TAG	UNP A0R4I6

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	MET	-	EXPRESSION TAG	UNP A0R4I6
B	-1	ALA	-	EXPRESSION TAG	UNP A0R4I6
B	0	SER	-	EXPRESSION TAG	UNP A0R4I6

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	245	Total O 245 245	0	0
2	B	257	Total O 257 257	0	0

- Molecule 1: Adenylosuccinate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.60Å 176.83Å 73.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.86 – 2.16 37.86 – 2.16	Depositor EDS
% Data completeness (in resolution range)	98.6 (37.86-2.16) 98.7 (37.86-2.16)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.56 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.205 , 0.238 0.211 , 0.237	Depositor DCC
R_{free} test set	2658 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 36.9	EDS
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 52213 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7472	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3597	0.45	0/4876
1	B	0.26	0/3478	0.44	0/4715
All	All	0.26	0/7075	0.45	0/9591

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3542	0	3491	11	0
1	B	3428	0	3351	6	0
2	A	245	0	0	0	0
2	B	257	0	0	0	0
All	All	7472	0	6842	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:ARG:NH2	1:A:466:ALA:O	2.33	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ILE:HG22	1:A:84:ILE:O	2.01	0.60
1:A:87:PHE:HE1	1:A:96:VAL:HG22	1.72	0.54
1:A:103:ARG:HH11	1:A:231:GLN:HE21	1.55	0.53
1:B:311:LEU:HD11	1:B:326:VAL:HG22	1.92	0.52
1:A:115[A]:ARG:HG2	1:A:115[A]:ARG:HH11	1.76	0.50
1:A:87:PHE:CE1	1:A:96:VAL:HG22	2.49	0.48
1:A:53:VAL:HG12	1:A:93:HIS:CD2	2.49	0.47
1:B:331:LEU:HB2	1:B:332:PRO:HD3	1.96	0.47
1:B:97:HIS:CA	1:B:97:HIS:CG	2.98	0.46
1:A:115[A]:ARG:HG2	1:A:115[A]:ARG:NH1	2.31	0.46
1:B:103:ARG:HH11	1:B:231:GLN:HE21	1.65	0.45
1:A:331:LEU:HB2	1:A:332:PRO:HD3	1.98	0.44
1:B:111:GLN:HE22	1:B:186:ARG:H	1.67	0.43
1:A:321:VAL:CG1	1:A:321:VAL:O	2.67	0.43
1:B:78:HIS:HB3	1:B:80:VAL:HG12	2.02	0.40
1:A:87:PHE:CE1	1:A:94:GLU:HA	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/488 (94%)	443 (96%)	14 (3%)	4 (1%)	21	13
1	B	446/488 (91%)	435 (98%)	11 (2%)	0	100	100
All	All	907/976 (93%)	878 (97%)	25 (3%)	4 (0%)	39	34

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	276	GLN
1	A	84	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	275	GLY
1	A	273	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/382 (93%)	348 (98%)	7 (2%)	63	67
1	B	339/382 (89%)	334 (98%)	5 (2%)	72	78
All	All	694/764 (91%)	682 (98%)	12 (2%)	68	74

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	85	GLU
1	A	103	ARG
1	A	282	MET
1	A	322	PHE
1	A	361	ARG
1	A	407	GLN
1	B	103	ARG
1	B	285	LYS
1	B	322	PHE
1	B	326	VAL
1	B	407	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	93	HIS
1	A	97	HIS
1	A	113	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	231	GLN
1	A	257	HIS
1	A	265	HIS
1	A	317	ASN
1	A	341	GLN
1	A	407	GLN
1	B	111	GLN
1	B	113	GLN
1	B	150	ASN
1	B	231	GLN
1	B	265	HIS
1	B	341	GLN
1	B	407	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	464/488 (95%)	0.55	44 (9%) 10 16	14, 28, 62, 84	0
1	B	451/488 (92%)	0.38	35 (7%) 16 22	15, 27, 59, 77	0
All	All	915/976 (93%)	0.47	79 (8%) 13 19	14, 28, 61, 84	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	SER	9.7
1	A	274	PRO	8.8
1	A	280	SER	8.7
1	A	277	VAL	8.6
1	A	273	ALA	7.1
1	A	272	PHE	7.0
1	A	276	GLN	6.9
1	B	48	ALA	6.0
1	A	89	ALA	5.5
1	A	275	GLY	5.4
1	B	77	ARG	4.9
1	A	49	VAL	4.6
1	B	71	ALA	4.5
1	B	70	ALA	4.3
1	B	51	ASP	4.2
1	B	53	VAL	4.0
1	A	278	GLY	4.0
1	A	281	ALA	3.9
1	B	76	THR	3.7
1	B	74	ARG	3.7
1	A	87	PHE	3.6
1	B	78	HIS	3.6
1	B	84	ILE	3.4
1	B	428	VAL	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	78	HIS	3.3
1	B	286	MET	3.2
1	A	54	VAL	3.2
1	B	80	VAL	3.1
1	A	67	GLU	3.1
1	A	271	GLY	3.1
1	A	419	GLY	3.1
1	A	286	MET	3.1
1	A	71	ALA	3.0
1	B	79	ASP	3.0
1	B	61	LEU	3.0
1	A	287	ASN	2.9
1	A	285	LYS	2.8
1	B	56	ASP	2.8
1	A	398	VAL	2.8
1	B	316	TRP	2.8
1	B	88	ASN	2.8
1	A	88	ASN	2.8
1	A	85	GLU	2.7
1	A	60	VAL	2.7
1	A	433	ALA	2.7
1	B	95	HIS	2.7
1	A	53	VAL	2.6
1	B	75	VAL	2.6
1	A	70	ALA	2.6
1	A	387	GLU	2.6
1	B	390	HIS	2.5
1	B	285	LYS	2.5
1	B	52	GLY	2.5
1	A	282	MET	2.5
1	B	60	VAL	2.5
1	A	396	HIS	2.4
1	B	381	ARG	2.4
1	A	429	ALA	2.3
1	A	47	VAL	2.3
1	B	89	ALA	2.3
1	B	43	ALA	2.3
1	A	284	HIS	2.3
1	B	203	GLY	2.2
1	A	427	LYS	2.2
1	A	66	LEU	2.2
1	B	72	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	314	ALA	2.2
1	B	45	LEU	2.2
1	A	316	TRP	2.2
1	B	59	ARG	2.2
1	A	323	CYS	2.1
1	B	438	GLN	2.1
1	A	93	HIS	2.1
1	B	207	GLY	2.1
1	A	59	ARG	2.1
1	B	82	ALA	2.1
1	B	73	GLU	2.0
1	A	439	ALA	2.0
1	A	428	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.