



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:03 AM GMT

PDB ID : 3NNZ
Title : Structure of rat neuronal nitric oxide synthase heme domain complexed with 6-(((3S,4S)-4-(2-(3-Fluorophenethylamino)ethoxy)pyrrolidin-3-yl)methyl)pyridin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2010-06-24
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	8664 (2.00-1.96)
Clashscore	102246	9905 (2.00-1.96)
Ramachandran outliers	100387	9792 (2.00-1.96)
Sidechain outliers	100360	9791 (2.00-1.96)
RSRZ outliers	91569	8679 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>21%</div> <div> <div></div> <div>85%</div> <div>11%</div> <div>..</div> </div> </div>
1	B	422	<div> <div>10%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	A	860	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7312 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	1	0
			3316	2123	566	605	22			
1	B	411	Total	C	N	O	S	0	3	0
			3354	2146	574	612	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 17	C 9	N 5	O 3	0	0
3	B	1	Total 17	C 9	N 5	O 3	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is 6-[[[(3S,4S)-4-(2-{[2-(3-FLUOROPHENYL)ETHYL]AMINO}ETHOXY)PYRROLIDIN-3-YL]METHYL]PYRIDIN-2-AMINE (three-letter code: 59W) (formula: C₂₀H₂₇FN₄O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	0	0
			26	20	1	4	1		
5	B	1	Total	C	F	N	O	0	0
			26	20	1	4	1		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	199	Total	O	0	0
			199	199		
7	B	262	Total	O	0	0
			262	262		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.05Å 110.98Å 164.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.64 – 1.97 37.96 – 1.97	Depositor EDS
% Data completeness (in resolution range)	97.8 (38.64-1.97) 97.8 (37.96-1.97)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.179 , 0.213 0.197 , 0.228	Depositor DCC
R_{free} test set	3313 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.172	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 66815 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7312	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, H4B, 59W, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/3412	0.63	0/4629
1	B	0.66	0/3456	0.65	0/4685
All	All	0.63	0/6868	0.64	0/9314

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	378	MET	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3316	0	3226	28	0
1	B	3354	0	3274	16	0
2	A	43	0	30	1	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	26	0	27	1	0
5	B	26	0	27	2	0
6	A	1	0	0	0	0
7	A	199	0	0	0	0
7	B	262	0	0	2	0
All	All	7312	0	6650	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:706:TYR:OH	2:B:750:HEM:O1D	2.02	0.77
1:A:371:ARG:HH21	1:A:371:ARG:HG3	1.50	0.76
1:A:371:ARG:HH21	1:A:371:ARG:CG	2.07	0.67
1:B:717:LYS:NZ	7:B:1177:HOH:O	2.31	0.63
1:A:307:GLU:HG3	7:B:1260:HOH:O	1.98	0.62
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.82	0.62
1:A:499:VAL:O	1:A:503:GLU:HG3	2.03	0.59
1:A:380:ARG:HD3	1:A:400:GLU:OE1	2.04	0.58
1:A:351:LYS:HE2	1:A:392:SER:HB3	1.87	0.57
1:B:592:GLU:OE1	5:B:800:59W:H5'A	2.05	0.57
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.87	0.56
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.89	0.55
1:A:371:ARG:CG	1:A:371:ARG:NH2	2.68	0.53
1:A:487:GLN:HB3	1:A:488:PRO:HD2	1.91	0.53
2:B:750:HEM:HBB2	2:B:750:HEM:HHC	1.90	0.53
1:A:378:MET:HA	1:A:378:MET:CE	2.40	0.52
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.41	0.51
1:A:537:PRO:HB2	1:A:540:LEU:HG	1.94	0.49
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.01	0.49
1:A:706:TYR:OH	2:A:750:HEM:O1D	2.19	0.47
1:B:619:ARG:HE	1:B:619:ARG:HB2	1.56	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:361:PHE:O	1:A:364:GLN:HG2	2.15	0.46
1:A:487:GLN:HB3	1:A:488:PRO:CD	2.46	0.46
1:A:316:LEU:HD12	1:A:319:LYS:HD2	1.96	0.46
1:B:391:THR:O	1:B:392:SER:HB2	2.15	0.46
1:A:475:TRP:CE2	1:A:531:PRO:HG3	2.52	0.45
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.53	0.44
1:A:332:MET:CE	1:B:301:LEU:HD22	2.48	0.43
1:B:504:ILE:O	1:B:508:GLN:HG2	2.18	0.43
1:B:551:PHE:HE2	1:B:632:GLU:HG3	1.83	0.42
1:A:494:GLY:O	1:A:496:PRO:HD3	2.20	0.41
1:B:353:GLN:HE21	1:B:353:GLN:HB3	1.68	0.41
1:A:317:HIS:HB2	1:B:330:ILE:HD12	2.03	0.41
1:B:362:LEU:HD11	1:B:384:VAL:HG21	2.01	0.41
1:B:325:GLY:O	1:B:332:MET:HG3	2.21	0.41
2:B:750:HEM:HBA2	5:B:800:59W:H7AA	2.03	0.41
1:A:678:TRP:HH2	5:A:800:59W:H3	1.85	0.41
1:A:323:GLU:HG2	1:B:328:GLU:HB3	2.01	0.41
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.09	0.40
1:B:680:VAL:HA	1:B:681:PRO:HD3	1.93	0.40
1:A:645:LYS:HE3	1:A:645:LYS:HB2	1.89	0.40
1:A:362:LEU:HA	1:A:362:LEU:HD23	1.93	0.40
1:A:588:TYR:CD1	1:A:593:ILE:HD11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/422 (96%)	395 (98%)	8 (2%)	1 (0%)	52	47
1	B	410/422 (97%)	406 (99%)	4 (1%)	0	100	100
All	All	814/844 (96%)	801 (98%)	12 (2%)	1 (0%)	56	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/377 (97%)	348 (96%)	16 (4%)	35	27
1	B	369/377 (98%)	363 (98%)	6 (2%)	70	71
All	All	733/754 (97%)	711 (97%)	22 (3%)	50	44

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	338	PRO
1	A	350	THR
1	A	371	ARG
1	A	378	MET
1	A	380	ARG
1	A	454	ASN
1	A	489	ASP
1	A	507	GLN
1	A	511	LYS
1	A	547	ARG
1	A	555	LYS
1	A	569	ASN
1	A	609	GLU
1	A	620	LYS
1	A	645	LYS
1	A	715	VAL
1	B	353	GLN
1	B	389	GLU
1	B	540	LEU
1	B	547	ARG
1	B	602[A]	SER
1	B	602[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	454	ASN
1	A	569	ASN
1	A	601	ASN
1	A	605	ASN
1	A	697	ASN
1	B	364	GLN
1	B	454	ASN
1	B	507	GLN
1	B	601	ASN
1	B	605	ASN
1	B	642	GLN
1	B	697	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	750	1	30,50,50	2.46	8 (26%)	24,82,82	2.53	11 (45%)
3	H4B	A	760	-	13,18,18	0.89	0	11,26,26	2.92	6 (54%)
5	59W	A	800	-	25,28,28	0.71	1 (4%)	25,36,36	1.61	4 (16%)
4	ACT	A	860	-	1,3,3	1.11	0	0,3,3	0.00	-
2	HEM	B	750	1	30,50,50	2.17	10 (33%)	24,82,82	2.74	13 (54%)
3	H4B	B	760	-	13,18,18	1.05	0	11,26,26	2.58	6 (54%)
5	59W	B	800	-	25,28,28	0.84	0	25,36,36	1.59	3 (12%)
4	ACT	B	860	-	1,3,3	1.05	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
5	59W	A	800	-	-	0/13/23/23	0/3/3/3
4	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
5	59W	B	800	-	-	0/13/23/23	0/3/3/3
4	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3B-C4B	-7.73	1.45	1.51
2	A	750	HEM	C3D-C4D	-6.94	1.42	1.51
2	B	750	HEM	C3B-C4B	-5.40	1.47	1.51
2	B	750	HEM	C3D-C4D	-5.07	1.45	1.51
2	B	750	HEM	C2C-C1C	-4.07	1.44	1.52
2	A	750	HEM	C2C-C1C	-3.95	1.45	1.52
2	A	750	HEM	C2B-C1B	-2.12	1.44	1.51
2	B	750	HEM	C2D-C3D	-2.09	1.48	1.54
2	A	750	HEM	C2D-C1D	-2.05	1.45	1.51
2	A	750	HEM	C3B-CAB	2.03	1.55	1.51
2	B	750	HEM	C3B-CAB	2.05	1.55	1.51
5	A	800	59W	C14-C13	2.05	1.41	1.37
2	B	750	HEM	CMA-C3A	2.24	1.56	1.51
2	B	750	HEM	C3C-CAC	2.35	1.55	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C1C-NC	2.61	1.39	1.36
2	B	750	HEM	FE-NC	2.78	2.06	1.95
2	B	750	HEM	CAA-C2A	2.90	1.57	1.52
2	A	750	HEM	C1C-NC	2.93	1.39	1.36
2	A	750	HEM	FE-NC	3.22	2.08	1.95

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	CBA-CAA-C2A	-4.71	104.09	112.53
2	B	750	HEM	CAA-C2A-C1A	-4.08	122.58	127.01
2	A	750	HEM	CBA-CAA-C2A	-3.79	105.73	112.53
2	A	750	HEM	CBD-CAD-C3D	-3.79	102.53	113.55
3	A	760	H4B	N3-C2-N1	-3.73	119.43	125.53
2	B	750	HEM	CBD-CAD-C3D	-3.50	103.38	113.55
5	B	800	59W	C3A-C2A-N1A	-3.28	118.41	122.41
5	A	800	59W	C3A-C2A-N1A	-3.27	118.42	122.41
3	B	760	H4B	N3-C2-N1	-3.12	120.41	125.53
2	A	750	HEM	C3C-CAC-CBC	-3.01	119.84	124.46
2	B	750	HEM	CMA-C3A-C4A	-2.99	123.42	128.36
2	B	750	HEM	C3C-CAC-CBC	-2.97	119.90	124.46
5	A	800	59W	C14-C13-C12	-2.34	120.33	123.35
2	A	750	HEM	CAA-C2A-C1A	-2.26	124.55	127.01
5	B	800	59W	C14-C13-C12	-2.21	120.50	123.35
2	B	750	HEM	C3B-C4B-NB	-2.06	107.69	111.63
3	B	760	H4B	N2-C2-N3	2.09	120.67	117.20
3	B	760	H4B	C4A-C8A-N8	2.16	120.97	118.43
3	B	760	H4B	C2-N1-C8A	2.21	119.51	114.54
5	A	800	59W	F13-C13-C14	2.22	122.23	118.52
2	A	750	HEM	C3B-C4B-CHC	2.23	126.30	123.16
2	B	750	HEM	C3B-C4B-CHC	2.57	126.78	123.16
2	B	750	HEM	CMD-C2D-C3D	2.57	125.72	114.35
2	A	750	HEM	CMD-C2D-C3D	2.69	126.25	114.35
2	B	750	HEM	C2D-C3D-C4D	2.73	106.13	101.50
2	A	750	HEM	C2D-C3D-C4D	2.80	106.24	101.50
3	A	760	H4B	C2-N1-C8A	2.92	121.10	114.54
3	A	760	H4B	N2-C2-N3	2.97	122.12	117.20
2	A	750	HEM	CMB-C2B-C3B	3.15	124.41	116.53
3	A	760	H4B	C4A-C8A-N8	3.21	122.21	118.43
2	B	750	HEM	CAD-C3D-C2D	3.65	123.70	113.22
2	B	750	HEM	CMB-C2B-C3B	3.77	125.94	116.53
2	A	750	HEM	CAD-C3D-C4D	4.01	126.61	112.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	760	H4B	C4-N3-C2	4.63	122.37	115.94
3	B	760	H4B	C4-N3-C2	4.69	122.45	115.94
2	B	750	HEM	CMC-C2C-C3C	4.83	128.58	116.53
5	A	800	59W	C6A-N1A-C2A	4.83	121.67	118.23
2	A	750	HEM	CAD-C3D-C2D	4.84	127.12	113.22
2	B	750	HEM	CAD-C3D-C4D	4.99	130.08	112.47
3	B	760	H4B	C4-C4A-C8A	5.03	119.12	114.56
5	B	800	59W	C6A-N1A-C2A	5.14	121.88	118.23
2	A	750	HEM	CMC-C2C-C3C	5.41	130.04	116.53
3	A	760	H4B	C4-C4A-C8A	5.41	119.46	114.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	1	0
5	A	800	59W	1	0
2	B	750	HEM	3	0
5	B	800	59W	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	1.19	88 (21%) 1 1	24, 49, 90, 117	0
1	B	411/422 (97%)	0.55	43 (10%) 8 10	24, 37, 61, 78	0
All	All	818/844 (96%)	0.87	131 (16%) 3 3	24, 41, 83, 117	0

All (131) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	9.1
1	A	715	VAL	8.2
1	A	355	PHE	7.7
1	A	716	TRP	7.2
1	A	488	PRO	6.8
1	A	506	ILE	6.0
1	A	388	ILE	5.5
1	B	619	ARG	5.4
1	A	392	SER	5.3
1	A	486	LYS	5.2
1	A	503	GLU	5.2
1	A	491	SER	5.1
1	A	300	PHE	5.1
1	A	386	LYS	5.0
1	B	350	THR	5.0
1	A	350	THR	4.6
1	A	351	LYS	4.6
1	A	385	ASN	4.5
1	A	490	GLY	4.5
1	A	352	ASP	4.3
1	A	619	ARG	4.3
1	B	620	LYS	4.2
1	B	348	VAL	4.1
1	A	391	THR	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	354	LEU	3.9
1	A	507	GLN	3.9
1	A	470	HIS	3.8
1	A	567	VAL	3.8
1	A	682	PRO	3.7
1	A	677	VAL	3.6
1	B	479	LEU	3.6
1	A	713	THR	3.6
1	A	393	THR	3.5
1	A	712	ASN	3.5
1	A	584	PHE	3.4
1	A	714	HIS	3.4
1	A	479	LEU	3.4
1	A	489	ASP	3.2
1	B	667	ARG	3.2
1	A	680	VAL	3.1
1	B	616	LEU	3.1
1	A	643	SER	3.1
1	B	567	VAL	3.1
1	B	299	ARG	3.1
1	A	511	LYS	3.0
1	B	679	ILE	3.0
1	B	310	VAL	3.0
1	B	691	PHE	3.0
1	A	595	VAL	3.0
1	A	550	LYS	3.0
1	A	416	VAL	3.0
1	A	389	GLU	2.9
1	A	469	LYS	2.9
1	A	514	ARG	2.9
1	A	504	ILE	2.9
1	A	353	GLN	2.9
1	A	678	TRP	2.9
1	A	617	ASP	2.9
1	A	679	ILE	2.9
1	A	667	ARG	2.8
1	A	487	GLN	2.8
1	A	593	ILE	2.8
1	B	677	VAL	2.8
1	A	390	SER	2.8
1	B	682	PRO	2.8
1	B	680	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	356	PRO	2.8
1	A	676	TRP	2.8
1	A	480	ILE	2.7
1	A	321	THR	2.7
1	A	382	GLU	2.7
1	B	355	PHE	2.7
1	B	595	VAL	2.7
1	A	552	ASP	2.7
1	A	681	PRO	2.7
1	B	718	GLY	2.7
1	A	642	GLN	2.6
1	B	301	LEU	2.6
1	B	681	PRO	2.6
1	A	508	GLN	2.6
1	A	415	CYS	2.6
1	A	683	MET	2.6
1	B	615	ASP	2.6
1	A	299	ARG	2.6
1	A	704	PHE	2.5
1	B	591	THR	2.5
1	A	594	GLY	2.5
1	B	683	MET	2.5
1	A	553	TRP	2.5
1	B	327	THR	2.5
1	A	551	PHE	2.4
1	B	311	VAL	2.4
1	A	685	GLY	2.4
1	A	338	PRO	2.4
1	A	485	TYR	2.4
1	B	351	LYS	2.4
1	A	328	GLU	2.3
1	B	416	VAL	2.3
1	B	352	ASP	2.3
1	A	612	LYS	2.3
1	B	550	LYS	2.3
1	B	480	ILE	2.3
1	B	328	GLU	2.3
1	A	505	CYS	2.3
1	B	593	ILE	2.3
1	B	687	ILE	2.3
1	B	329	HIS	2.2
1	A	630	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	564	LEU	2.2
1	B	321	THR	2.2
1	B	338	PRO	2.2
1	B	676	TRP	2.2
1	A	591	THR	2.2
1	A	381	LEU	2.2
1	A	373	GLY	2.2
1	A	561	TRP	2.2
1	B	678	TRP	2.2
1	A	609	GLU	2.2
1	A	710	PRO	2.1
1	A	468	GLY	2.1
1	A	500	GLN	2.1
1	A	493	LEU	2.1
1	A	375	LYS	2.1
1	B	584	PHE	2.1
1	A	528	GLY	2.1
1	A	588	TYR	2.1
1	A	311	VAL	2.1
1	A	620	LYS	2.1
1	A	582	CYS	2.1
1	B	562	TYR	2.0
1	B	685	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ACT	A	860	4/4	0.90	0.17	3.81	53,55,56,56	0
5	59W	B	800	26/26	0.88	0.23	1.73	32,41,53,55	0
5	59W	A	800	26/26	0.91	0.23	1.00	31,40,57,60	0
2	HEM	B	750	43/43	0.97	0.18	0.89	21,26,36,44	0
3	H4B	B	760	17/17	0.97	0.19	0.88	24,29,31,34	0
3	H4B	A	760	17/17	0.96	0.21	0.79	27,29,33,33	0
2	HEM	A	750	43/43	0.97	0.20	0.67	23,29,39,41	0
4	ACT	B	860	4/4	0.97	0.12	0.64	40,42,44,44	0
6	ZN	A	900	1/1	1.00	0.04	-2.33	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.