



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:08 PM GMT

PDB ID : 1NOL
Title : OXYGENATED HEMOCYANIN (SUBUNIT TYPE II)
Authors : Hazes, B.; Hol, W.G.J.
Deposited on : 1995-10-17
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	628	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOCYANIN (SUBUNIT TYPE II).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	607	Total	C	N	O	S	0	0	0
			4933	3145	862	904	22			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	ILE	VAL	CONFLICT	UNP P04253

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		

- Molecule 5 is PEROXIDE ION (three-letter code: PER) (formula: O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	212	Total 212	O 212	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HEMOCYANIN (SUBUNIT TYPE II)



4 Data and refinement statistics

Property	Value	Source
Space group	R 3 2	Depositor
Cell constants a, b, c, α , β , γ	117.00Å 117.00Å 117.00Å 60.02° 60.02° 60.02°	Depositor
Resolution (Å)	8.00 – 2.40 33.79 – 2.41	Depositor EDS
% Data completeness (in resolution range)	83.4 (8.00-2.40) 79.5 (33.79-2.41)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	X-PLOR 2.1	Depositor
R, R_{free}	0.181 , (Not available) 0.168 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	(Not available)	Xtriage
Anisotropy	(Not available)	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 67.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtriage
Outliers	(Not available)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5154	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PER, CU, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/5077	1.59	66/6880 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	481	ARG	NE-CZ-NH2	-13.55	113.53	120.30
1	A	273	ARG	NE-CZ-NH1	11.37	125.99	120.30
1	A	100	ARG	NE-CZ-NH1	10.66	125.63	120.30
1	A	481	ARG	NE-CZ-NH1	10.30	125.45	120.30
1	A	195	ARG	NE-CZ-NH1	8.46	124.53	120.30
1	A	460	ARG	NE-CZ-NH1	8.19	124.39	120.30
1	A	163	TYR	CB-CG-CD1	-8.14	116.12	121.00
1	A	411	ARG	NE-CZ-NH1	8.12	124.36	120.30
1	A	273	ARG	NE-CZ-NH2	-7.89	116.35	120.30
1	A	365	ARG	NE-CZ-NH1	7.72	124.16	120.30
1	A	174	TRP	CE2-CD2-CG	-7.67	101.16	107.30
1	A	363	TRP	CD1-CG-CD2	7.60	112.38	106.30
1	A	176	TRP	CD1-CG-CD2	7.59	112.37	106.30
1	A	195	ARG	NE-CZ-NH2	-7.57	116.51	120.30
1	A	365	ARG	NE-CZ-NH2	-7.52	116.54	120.30
1	A	33	ARG	NE-CZ-NH1	7.52	124.06	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	TRP	CD1-CG-CD2	7.41	112.23	106.30
1	A	162	TYR	CB-CG-CD2	-7.35	116.59	121.00
1	A	538	TRP	CE2-CD2-CG	-7.20	101.54	107.30
1	A	326	TRP	CD1-CG-CD2	7.17	112.03	106.30
1	A	356	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	A	538	TRP	CD1-CG-CD2	7.16	112.03	106.30
1	A	563	TRP	CD1-CG-CD2	7.12	112.00	106.30
1	A	163	TYR	CB-CG-CD2	7.05	125.23	121.00
1	A	326	TRP	CE2-CD2-CG	-7.03	101.67	107.30
1	A	176	TRP	CE2-CD2-CG	-6.94	101.75	107.30
1	A	363	TRP	CE2-CD2-CG	-6.77	101.88	107.30
1	A	65	TRP	CE2-CD2-CG	-6.69	101.94	107.30
1	A	602	ARG	NE-CZ-NH1	6.64	123.62	120.30
1	A	174	TRP	CG-CD2-CE3	6.59	139.83	133.90
1	A	174	TRP	CD1-CG-CD2	6.49	111.49	106.30
1	A	100	ARG	NE-CZ-NH2	-6.43	117.08	120.30
1	A	58	ARG	NE-CZ-NH1	6.32	123.46	120.30
1	A	480	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	A	164	ARG	NE-CZ-NH1	6.18	123.39	120.30
1	A	309	GLU	CA-C-N	6.16	130.75	117.20
1	A	184	TRP	CE2-CD2-CG	-6.13	102.40	107.30
1	A	411	ARG	NE-CZ-NH2	-6.01	117.29	120.30
1	A	356	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	A	432	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	A	501	VAL	CG1-CB-CG2	-5.87	101.50	110.90
1	A	563	TRP	CE2-CD2-CG	-5.82	102.65	107.30
1	A	31	SER	N-CA-C	-5.79	95.36	111.00
1	A	145	VAL	CG1-CB-CG2	-5.70	101.78	110.90
1	A	184	TRP	CD1-CG-CD2	5.57	110.75	106.30
1	A	326	TRP	CG-CD2-CE3	5.48	138.83	133.90
1	A	164	ARG	NE-CZ-NH2	-5.47	117.57	120.30
1	A	580	VAL	CB-CA-C	-5.43	101.08	111.40
1	A	481	ARG	CG-CD-NE	-5.41	100.44	111.80
1	A	308	VAL	CG1-CB-CG2	-5.39	102.28	110.90
1	A	174	TRP	CB-CG-CD1	-5.37	120.02	127.00
1	A	120	ILE	CA-CB-CG1	-5.34	100.85	111.00
1	A	61	TYR	CA-CB-CG	5.30	123.47	113.40
1	A	230	ASP	CB-CA-C	-5.22	99.96	110.40
1	A	514	GLN	CA-CB-CG	5.19	124.81	113.40
1	A	298	LEU	CA-CB-CG	5.18	127.22	115.30
1	A	478	GLU	CA-CB-CG	5.18	124.79	113.40
1	A	568	VAL	CG1-CB-CG2	-5.18	102.62	110.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	TYR	CB-CG-CD2	-5.16	117.90	121.00
1	A	33	ARG	NE-CZ-NH2	-5.13	117.73	120.30
1	A	139	ASP	CA-CB-CG	5.11	124.64	113.40
1	A	8	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	58	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	602	ARG	NE-CZ-NH2	-5.02	117.79	120.30
1	A	340	ARG	NE-CZ-NH1	5.02	122.81	120.30
1	A	52	ASP	CB-CG-OD1	5.02	122.81	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	434	TYR	Sidechain
1	A	525	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4933	0	4680	54	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
4	A	4	0	0	1	0
5	A	2	0	0	0	0
6	A	212	0	0	7	0
All	All	5154	0	4680	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:MET:HB3	6:A:808:HOH:O	1.80	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:GLN:H	1:A:112:GLN:HE21	1.35	0.75
1:A:559:MET:HG3	1:A:617:MET:HG2	1.70	0.73
1:A:211:TYR:HE2	1:A:223:MET:HG3	1.53	0.72
1:A:144:LEU:HD21	1:A:432:ARG:HH21	1.55	0.70
1:A:359:ILE:HA	1:A:362:ASN:HD22	1.58	0.69
1:A:442:SER:HB3	1:A:503:HIS:HD2	1.63	0.63
1:A:95:VAL:HG23	1:A:178:LEU:HD22	1.81	0.62
1:A:9:ILE:HD12	1:A:106:LEU:HD13	1.83	0.61
1:A:259:ARG:HG3	6:A:809:HOH:O	2.03	0.57
1:A:481:ARG:HD2	6:A:690:HOH:O	2.04	0.57
1:A:144:LEU:HD21	1:A:432:ARG:NH2	2.22	0.55
1:A:259:ARG:HE	1:A:337:PRO:HG3	1.73	0.54
1:A:128:ALA:HB1	1:A:429:ILE:HG21	1.89	0.54
1:A:574:VAL:HA	6:A:807:HOH:O	2.08	0.54
1:A:233:LEU:HD23	1:A:359:ILE:HG12	1.90	0.53
1:A:359:ILE:HD12	1:A:362:ASN:HD22	1.74	0.52
1:A:80:ALA:O	1:A:84:VAL:HG22	2.10	0.51
1:A:576:CYS:HB2	1:A:586:ARG:HB2	1.93	0.50
1:A:524:VAL:O	1:A:526:LEU:HG	2.12	0.50
1:A:103:CYS:HA	1:A:106:LEU:HD12	1.95	0.48
1:A:404:ASN:ND2	1:A:620:LYS:HE2	2.29	0.48
1:A:112:GLN:H	1:A:112:GLN:NE2	2.08	0.47
1:A:387:ASN:ND2	6:A:786:HOH:O	2.47	0.47
1:A:514:GLN:HG3	6:A:720:HOH:O	2.13	0.47
1:A:475:LYS:HD2	1:A:607:GLU:HG2	1.97	0.47
1:A:327:GLY:HA3	1:A:363:TRP:CZ2	2.51	0.46
1:A:590:TYR:HA	1:A:591:PRO:HD3	1.67	0.46
1:A:315:VAL:HG13	6:A:784:HOH:O	2.14	0.46
1:A:359:ILE:HD12	1:A:362:ASN:ND2	2.31	0.45
1:A:170:ASN:HB3	1:A:352:SER:O	2.16	0.45
1:A:519:ASP:HA	1:A:524:VAL:HB	1.99	0.45
1:A:394:GLN:OE1	1:A:497:LYS:HE2	2.17	0.44
1:A:333:ARG:NH2	4:A:633:NO3:O3	2.51	0.44
1:A:574:VAL:CG1	1:A:585:ALA:HB1	2.48	0.43
1:A:120:ILE:HA	1:A:121:PRO:HD3	1.77	0.43
1:A:532:GLU:OE1	1:A:576:CYS:HA	2.18	0.43
1:A:101:ASP:O	1:A:104:LYS:HB2	2.19	0.43
1:A:15:GLN:NE2	1:A:418:LYS:O	2.53	0.42
1:A:568:VAL:HG23	1:A:593:LYS:HG3	2.02	0.42
1:A:19:ALA:HB2	1:A:81:ARG:HH21	1.84	0.42
1:A:231:GLU:HG3	1:A:232:PRO:HD2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:GLN:HA	1:A:119:PHE:CD1	2.54	0.42
1:A:81:ARG:HB3	1:A:89:PHE:CE1	2.55	0.41
1:A:76:ILE:HA	1:A:76:ILE:HD12	1.84	0.41
1:A:359:ILE:HA	1:A:359:ILE:HD12	1.95	0.41
1:A:142:PRO:HG3	1:A:427:ARG:HE	1.85	0.41
1:A:473:GLU:OE1	1:A:475:LYS:HE2	2.21	0.41
1:A:131:LYS:HE3	1:A:143:ILE:HG23	2.02	0.41
1:A:475:LYS:HD3	1:A:475:LYS:HA	1.88	0.41
1:A:548:ASN:O	1:A:625:LYS:HA	2.20	0.40
1:A:185:ASN:HA	1:A:186:PRO:HD3	1.83	0.40
1:A:466:LYS:HG3	1:A:555:HIS:CE1	2.56	0.40
1:A:436:LEU:O	1:A:541:HIS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	597/628 (95%)	554 (93%)	38 (6%)	5 (1%)	24	35

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	31	SER
1	A	140	GLU
1	A	245	SER
1	A	148	GLN
1	A	423	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	530/558 (95%)	496 (94%)	34 (6%)	22	34

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	HIS
1	A	37	VAL
1	A	41	GLN
1	A	61	TYR
1	A	101	ASP
1	A	104	LYS
1	A	112	GLN
1	A	120	ILE
1	A	139	ASP
1	A	150	THR
1	A	153	ILE
1	A	169	ILE
1	A	187	LYS
1	A	231	GLU
1	A	278	ASP
1	A	352	SER
1	A	382	ASP
1	A	387	ASN
1	A	388	PHE
1	A	401	ARG
1	A	423	PRO
1	A	427	ARG
1	A	428	SER
1	A	453	SER
1	A	475	LYS
1	A	492	ASP
1	A	514	GLN
1	A	518	GLU
1	A	526	LEU
1	A	531	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	574	VAL
1	A	580	VAL
1	A	610	SER
1	A	627	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	112	GLN
1	A	239	HIS
1	A	362	ASN
1	A	372	HIS
1	A	387	ASN
1	A	399	HIS
1	A	413	GLN
1	A	503	HIS
1	A	514	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PER	A	631	2	0,1,1	0.00	-	0,0,0	0.00	-
4	NO3	A	633	-	3,3,3	0.25	0	3,3,3	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PER	A	631	2	-	0/0/0/0	0/0/0/0
4	NO3	A	633	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	633	NO3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	607/628 (96%)	-0.78	8 (1%) 79 79	4, 15, 31, 45	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	21	VAL	4.4
1	A	525	GLY	2.8
1	A	139	ASP	2.6
1	A	244	ALA	2.4
1	A	150	THR	2.3
1	A	524	VAL	2.2
1	A	152	ASN	2.2
1	A	133	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NO3	A	633	4/4	0.99	0.08	-0.67	15,16,16,16	0
2	CU	A	630	1/1	0.99	0.10	-0.69	11,11,11,11	0
5	PER	A	631	2/2	1.00	0.10	-0.85	7,7,7,7	0
3	CA	A	632	1/1	0.98	0.07	-1.39	21,21,21,21	0
2	CU	A	629	1/1	1.00	0.10	-1.49	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.