



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:08 PM GMT

PDB ID : 1NOS
Title : MURINE INDUCIBLE NITRIC OXIDE SYNTHASE OXYGENASE DOMAIN (DELTA 114), IMIDAZOLE COMPLEX
Authors : Crane, B.R.; Arvai, A.S.; Getzoff, E.D.; Stuehr, D.J.; Tainer, J.A.
Deposited on : 1997-09-28
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	388	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INDUCIBLE NITRIC OXIDE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2613	1672	446	480	15			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	A	1	Total	C	N	0	0
			5	3	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	255	Total	O	1	0
			255	255		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: INDUCIBLE NITRIC OXIDE SYNTHASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.00 Å 73.80 Å 92.80 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10	Depositor
% Data completeness (in resolution range)	99.7 (30.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.205 , 0.261	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2921	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2684	0.74	8/3647 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	336	LEU	N-CA-C	-8.14	89.03	111.00
1	A	377	PHE	N-CA-C	7.61	131.56	111.00
1	A	371	GLU	N-CA-C	5.88	126.86	111.00
1	A	378	CYS	N-CA-C	-5.60	95.87	111.00
1	A	501	HIS	N-CA-C	-5.26	96.81	111.00
1	A	337	GLY	N-CA-C	5.21	126.12	113.10
1	A	376	ASP	N-CA-C	-5.20	96.97	111.00
1	A	292	ARG	N-CA-C	-5.14	97.13	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2613	0	2504	63	0
2	A	43	0	30	3	0
3	A	10	0	9	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	255	0	0	9	0
All	All	2921	0	2543	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:ASP:H	1:A:496:GLN:NE2	1.83	0.76
1:A:286:ASP:HB3	1:A:501:HIS:O	1.86	0.74
1:A:271:ILE:HG12	1:A:272:ARG:N	2.07	0.70
1:A:210:ASN:HD22	1:A:210:ASN:H	1.39	0.69
1:A:223:ARG:HD3	4:A:1035:HOH:O	1.91	0.69
1:A:141:ILE:HD11	1:A:163:VAL:HG21	1.73	0.69
1:A:415:ALA:O	1:A:419:SER:HB2	1.94	0.68
1:A:141:ILE:HG12	4:A:1047:HOH:O	1.94	0.68
1:A:377:PHE:HB2	1:A:379:ASP:H	1.61	0.65
1:A:271:ILE:HD11	1:A:273:GLY:O	1.97	0.65
1:A:309:ASP:H	1:A:496:GLN:HE21	1.48	0.60
1:A:414:VAL:HG13	1:A:415:ALA:H	1.67	0.60
1:A:495:TRP:CE2	1:A:502:HIS:HB3	2.38	0.59
1:A:366:TRP:H	2:A:901:HEM:HAB	1.67	0.58
1:A:194:CYS:O	1:A:197:ARG:HD3	2.04	0.58
1:A:115:ASN:N	1:A:116:PRO:HD2	2.19	0.57
1:A:435:GLU:O	1:A:439:LYS:HG3	2.06	0.56
1:A:410:THR:HG22	1:A:410:THR:O	2.06	0.56
1:A:414:VAL:HG13	1:A:415:ALA:N	2.22	0.55
1:A:226:LEU:HD23	1:A:319:LEU:CD2	2.37	0.55
1:A:478:VAL:HG11	4:A:956:HOH:O	2.06	0.55
1:A:152:LYS:HD3	1:A:155:GLU:HG2	1.90	0.54
1:A:324:THR:H	1:A:423:GLN:HE22	1.55	0.54
1:A:151:ALA:HB3	4:A:1103:HOH:O	2.08	0.53
1:A:160:LEU:HD21	4:A:1060:HOH:O	2.09	0.52
1:A:372:ILE:HG13	1:A:372:ILE:O	2.08	0.52
2:A:901:HEM:HBD1	2:A:901:HEM:HHA	1.91	0.52
1:A:337:GLY:O	1:A:338:LEU:HD23	2.12	0.50
1:A:115:ASN:N	1:A:116:PRO:CD	2.75	0.49
1:A:241:PHE:HB3	1:A:242:PRO:CD	2.43	0.48
1:A:495:TRP:CZ2	1:A:502:HIS:HB3	2.48	0.48
1:A:254:TRP:HE3	1:A:283:LEU:HD21	1.77	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:LEU:HD23	4:A:1046:HOH:O	2.14	0.47
1:A:129:PRO:HB2	1:A:131:GLU:OE2	2.13	0.47
1:A:342:ALA:HB1	1:A:425:VAL:HG11	1.97	0.47
1:A:142:ASN:HD22	1:A:156:HIS:HE1	1.63	0.47
1:A:272:ARG:HH11	1:A:272:ARG:HG3	1.81	0.46
1:A:371:GLU:O	1:A:373:GLY:N	2.49	0.46
1:A:195:ILE:O	1:A:198:ILE:HG12	2.17	0.45
1:A:149:LYS:HB3	1:A:149:LYS:HE3	1.75	0.45
1:A:256:SER:HA	1:A:348:ASN:ND2	2.31	0.45
1:A:438:MET:O	1:A:442:GLN:HG3	2.17	0.45
1:A:379:ASP:C	1:A:381:GLN:N	2.69	0.45
1:A:271:ILE:HD12	1:A:275:ALA:HB2	1.98	0.45
1:A:210:ASN:ND2	1:A:210:ASN:H	2.12	0.44
1:A:130:LEU:HD22	1:A:167:ILE:HG22	1.98	0.44
1:A:278:LEU:HD23	1:A:278:LEU:HA	1.85	0.44
1:A:257:GLN:HA	1:A:345:ALA:O	2.17	0.44
1:A:364:ASN:O	2:A:901:HEM:HMC2	2.18	0.44
1:A:418:HIS:O	1:A:422:LYS:HG2	2.19	0.43
1:A:337:GLY:C	1:A:338:LEU:HD23	2.39	0.43
1:A:239:THR:O	1:A:361:CYS:HA	2.18	0.43
1:A:414:VAL:HG23	4:A:977:HOH:O	2.18	0.43
1:A:272:ARG:NH1	1:A:272:ARG:HG3	2.34	0.42
1:A:250:ASP:HB2	4:A:980:HOH:O	2.18	0.42
1:A:190:ASN:O	1:A:192:PRO:HD3	2.20	0.42
1:A:132:GLU:O	1:A:136:HIS:HD2	2.03	0.42
1:A:223:ARG:NH2	4:A:1112:HOH:O	2.52	0.41
1:A:152:LYS:HA	1:A:155:GLU:OE1	2.20	0.41
1:A:283:LEU:O	1:A:287:LEU:HG	2.20	0.41
1:A:253:LEU:HD22	1:A:347:ALA:HB1	2.03	0.41
1:A:241:PHE:HB3	1:A:242:PRO:HD2	2.03	0.41
1:A:374:VAL:HG12	1:A:374:VAL:O	2.21	0.40
1:A:374:VAL:C	1:A:376:ASP:H	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	315/388 (81%)	286 (91%)	22 (7%)	7 (2%)	8 3

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	149	LYS
1	A	372	ILE
1	A	373	GLY
1	A	499	HIS
1	A	379	ASP
1	A	378	CYS
1	A	497	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	278/340 (82%)	252 (91%)	26 (9%)	11 7

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	117	LYS
1	A	150	GLU
1	A	153	ILE
1	A	177	LEU
1	A	178	ASP
1	A	199	GLN
1	A	210	ASN
1	A	226	LEU
1	A	271	ILE
1	A	318	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	325	MET
1	A	338	LEU
1	A	370	THR
1	A	371	GLU
1	A	376	ASP
1	A	377	PHE
1	A	381	GLN
1	A	413	ASN
1	A	417	LEU
1	A	418	HIS
1	A	419	SER
1	A	427	ILE
1	A	438	MET
1	A	497	ASN
1	A	498	GLU
1	A	501	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	HIS
1	A	143	GLN
1	A	156	HIS
1	A	199	GLN
1	A	202	ASN
1	A	210	ASN
1	A	230	ASN
1	A	257	GLN
1	A	421	GLN
1	A	443	ASN
1	A	496	GLN
1	A	502	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	901	1,3	30,50,50	2.74	8 (26%)	24,82,82	2.32	8 (33%)
3	IMD	A	902	2	3,5,5	0.21	0	4,5,5	0.84	0
3	IMD	A	903	-	3,5,5	0.15	0	4,5,5	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	901	1,3	-	0/10/54/54	0/0/8/8
3	IMD	A	902	2	-	0/0/0/0	0/1/1/1
3	IMD	A	903	-	-	0/0/0/0	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	HEM	C2D-C3D	-7.11	1.33	1.54
2	A	901	HEM	C3D-C4D	-6.86	1.42	1.51
2	A	901	HEM	C3B-CAB	-6.28	1.39	1.51
2	A	901	HEM	C3C-CAC	-5.28	1.41	1.51
2	A	901	HEM	C3B-C4B	-3.92	1.48	1.51
2	A	901	HEM	C2C-C1C	-3.12	1.46	1.52
2	A	901	HEM	C2B-C1B	-2.79	1.42	1.51
2	A	901	HEM	C2D-C1D	-2.75	1.42	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	901	HEM	CMD-C2D-C3D	2.61	125.88	114.35
2	A	901	HEM	C3B-C4B-CHC	3.13	127.57	123.16
2	A	901	HEM	C2D-C3D-C4D	3.16	106.86	101.50
2	A	901	HEM	CMC-C2C-C3C	3.93	126.35	116.53
2	A	901	HEM	C3C-CAC-CBC	3.98	130.57	124.46
2	A	901	HEM	CAD-C3D-C4D	4.10	126.93	112.47
2	A	901	HEM	CAD-C3D-C2D	4.52	126.20	113.22
2	A	901	HEM	CMB-C2B-C3B	4.85	128.64	116.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.