



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2016 – 03:12 PM EDT

PDB ID : 4NOE
Title : Crystal structure of DdrB bound to 30b ssDNA
Authors : Sugiman-Marangos, S.N.; Weiss, Y.M.; Junop, M.S.
Deposited on : 2013-11-19
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027257
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027257


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	<div> <div>86%</div> <div>7% • 6%</div> </div>
1	B	148	<div> <div>80%</div> <div>9% • 9%</div> </div>
1	C	148	<div> <div>81%</div> <div>9% • 9%</div> </div>
1	D	148	<div> <div>%</div> <div>81%</div> <div>9% 10%</div> </div>
1	E	148	<div> <div>86%</div> <div>9% •</div> </div>
2	F	30	<div> <div>10%</div> <div>37%</div> <div>53%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single-stranded DNA-binding protein DdrB.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	0	0	0
			1035	655	192	188			
1	B	134	Total	C	N	O	0	0	0
			1034	658	198	178			
1	C	134	Total	C	N	O	0	0	0
			1012	646	193	173			
1	D	133	Total	C	N	O	0	0	0
			1002	638	185	179			
1	E	142	Total	C	N	O	0	0	0
			1062	675	197	190			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ASP	-	EXPRESSION TAG	UNP Q9RY80
A	-2	PRO	-	EXPRESSION TAG	UNP Q9RY80
A	-1	PHE	-	EXPRESSION TAG	UNP Q9RY80
A	0	THR	-	EXPRESSION TAG	UNP Q9RY80
B	-3	ASP	-	EXPRESSION TAG	UNP Q9RY80
B	-2	PRO	-	EXPRESSION TAG	UNP Q9RY80
B	-1	PHE	-	EXPRESSION TAG	UNP Q9RY80
B	0	THR	-	EXPRESSION TAG	UNP Q9RY80
C	-3	ASP	-	EXPRESSION TAG	UNP Q9RY80
C	-2	PRO	-	EXPRESSION TAG	UNP Q9RY80
C	-1	PHE	-	EXPRESSION TAG	UNP Q9RY80
C	0	THR	-	EXPRESSION TAG	UNP Q9RY80
D	-3	ASP	-	EXPRESSION TAG	UNP Q9RY80
D	-2	PRO	-	EXPRESSION TAG	UNP Q9RY80
D	-1	PHE	-	EXPRESSION TAG	UNP Q9RY80
D	0	THR	-	EXPRESSION TAG	UNP Q9RY80
E	-3	ASP	-	EXPRESSION TAG	UNP Q9RY80
E	-2	PRO	-	EXPRESSION TAG	UNP Q9RY80
E	-1	PHE	-	EXPRESSION TAG	UNP Q9RY80

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	0	THR	-	EXPRESSION TAG	UNP Q9RY80

- Molecule 2 is a DNA chain called 5'-D(*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	30	Total 610	C 290	N 100	O 190	P 30	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Ca 1	0	0
3	D	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	110	Total 110	O 110	0	0
4	B	104	Total 104	O 104	0	0
4	C	102	Total 102	O 102	0	0
4	D	106	Total 106	O 106	0	0
4	E	91	Total 91	O 91	0	0
4	F	33	Total 33	O 33	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Single-stranded DNA-binding protein DdrB

Chain A: 




- Molecule 1: Single-stranded DNA-binding protein DdrB

Chain B: 




- Molecule 1: Single-stranded DNA-binding protein DdrB

Chain C: 



- Molecule 1: Single-stranded DNA-binding protein DdrB

Chain D: 



- Molecule 1: Single-stranded DNA-binding protein DdrB

Chain E: 



- Molecule 2: 5'-D(*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP*TP*TP*GP*CP*GP*CP)-3'

Chain F: 

T1	T2	G3	C4	G5	G6	T7	T8	G9	C10	G11	C12	T13	T14	G15	C16	G17	C18	T19	T20	G21	C22	G23	C24	T25	T26	G27	C28	G29	C30
----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	131.76Å 131.76Å 102.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.88 – 2.20 68.88 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (65.88-2.20) 100.0 (68.88-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.190 , 0.232 0.188 , 0.228	Depositor DCC
R_{free} test set	4452 reflections (5.10%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6303	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	0/1056	0.56	0/1434
1	B	0.37	0/1057	0.56	1/1431 (0.1%)
1	C	0.39	0/1035	0.57	1/1405 (0.1%)
1	D	0.50	0/1024	0.56	0/1391
1	E	0.36	0/1086	0.53	0/1476
2	F	1.69	12/679 (1.8%)	1.33	11/1046 (1.1%)
All	All	0.69	12/5937 (0.2%)	0.70	13/8183 (0.2%)

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	3	DG	O3'-P	-10.00	1.49	1.61
2	F	19	DT	O3'-P	-8.89	1.50	1.61
2	F	7	DT	O3'-P	-8.34	1.51	1.61
2	F	13	DT	O3'-P	-7.77	1.51	1.61
2	F	25	DT	O3'-P	-7.74	1.51	1.61

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	9	DG	O5'-P-OP1	-18.38	88.65	110.70
2	F	27	DG	O5'-P-OP1	-12.09	94.82	105.70
2	F	21	DG	O5'-P-OP1	-8.47	98.08	105.70
2	F	9	DG	O5'-P-OP2	7.99	120.29	110.70
2	F	6	DC	C1'-O4'-C4'	-7.40	102.70	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1035	0	957	7	0
1	B	1034	0	989	13	0
1	C	1012	0	953	9	0
1	D	1002	0	931	19	0
1	E	1062	0	988	10	0
2	F	610	0	341	44	4
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	A	110	0	0	1	0
4	B	104	0	0	5	0
4	C	102	0	0	2	0
4	D	106	0	0	4	0
4	E	91	0	0	5	0
4	F	33	0	0	6	0
All	All	6303	0	5159	81	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:GLU:HG3	1:A:136:ARG:HB2	1.53	0.91
1:B:13:ARG:NH1	4:B:393:HOH:O	2.02	0.90
2:F:1:DT:H5"	2:F:2:DT:H72	1.61	0.81
1:C:83:ARG:NH1	4:C:246:HOH:O	2.18	0.76
1:E:120:ASP:OD2	4:E:205:HOH:O	2.03	0.76

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:12:DC:O2	2:F:27:DG:N2[8_666]	1.37	0.83
2:F:15:DG:N2	2:F:24:DC:O2[8_666]	1.73	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:9:DG:N2	2:F:30:DC:O2[8_666]	1.81	0.39
2:F:3:DG:N2	2:F:6:DC:O2[8_666]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/148 (91%)	134 (99%)	1 (1%)	0	100	100
1	B	128/148 (86%)	128 (100%)	0	0	100	100
1	C	128/148 (86%)	128 (100%)	0	0	100	100
1	D	127/148 (86%)	125 (98%)	2 (2%)	0	100	100
1	E	138/148 (93%)	135 (98%)	3 (2%)	0	100	100
All	All	656/740 (89%)	650 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/123 (76%)	90 (97%)	3 (3%)	46	57
1	B	96/123 (78%)	94 (98%)	2 (2%)	61	74
1	C	91/123 (74%)	90 (99%)	1 (1%)	80	89
1	D	91/123 (74%)	90 (99%)	1 (1%)	80	89

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	96/123 (78%)	94 (98%)	2 (2%)	61	74
All	All	467/615 (76%)	458 (98%)	9 (2%)	65	77

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	87	LEU
1	E	25	LEU
1	D	25	LEU
1	A	97	LEU
1	C	13	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9
1	A	139/148 (93%)	-0.31	0	100100	12, 22, 42, 62	0
1	B	134/148 (90%)	-0.19	0	100100	11, 26, 46, 63	0
1	C	134/148 (90%)	-0.23	0	100100	12, 25, 46, 56	0
1	D	133/148 (89%)	-0.29	1 (0%)	8787	12, 24, 41, 58	0
1	E	142/148 (95%)	-0.19	0	100100	10, 26, 47, 61	0
2	F	30/30 (100%)	0.13	0	100100	26, 45, 57, 60	0
All	All	712/770 (92%)	-0.23	1 (0%)	9595	10, 25, 48, 63	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	66	TRP	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CA	D	201	1/1	0.98	0.06	-3.73	34,34,34,34	0
3	CA	B	201	1/1	0.96	0.07	-	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.