



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:03 AM GMT

PDB ID : 3NXD
Title : JC polyomavirus VP1 in complex with LSTc
Authors : Neu, U.; Stroeh, L.J.; Stehle, T.
Deposited on : 2010-07-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>5%</div> </div> </div>
1	B	272	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>5%</div> </div> </div>
1	C	272	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>5%</div> </div> </div>
1	D	272	<div> <div></div> <div> <div></div> <div>91%</div> <div>5%</div> </div> </div>
1	E	272	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	3	-	-	-	X
3	GOL	A	4	-	-	-	X
3	GOL	B	290	-	-	-	X
3	GOL	C	290	-	-	-	X
3	GOL	D	2	-	-	-	X
3	GOL	E	3	-	-	-	X
4	EDO	A	5	-	-	-	X
4	EDO	A	6	-	-	-	X
4	EDO	C	291	-	-	-	X
4	EDO	D	290	-	-	-	X
4	EDO	E	290	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major capsid protein VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	261	Total	C	N	O	S	0	4	0
			2055	1295	352	397	11			
1	B	258	Total	C	N	O	S	0	5	0
			2023	1277	345	389	12			
1	A	259	Total	C	N	O	S	0	5	0
			2039	1286	348	394	11			
1	D	261	Total	C	N	O	S	0	3	0
			2043	1287	350	395	11			
1	E	258	Total	C	N	O	S	0	5	0
			2040	1286	350	393	11			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	18	GLY	-	EXPRESSION TAG	UNP P03089
C	19	SER	-	EXPRESSION TAG	UNP P03089
C	20	HIS	-	EXPRESSION TAG	UNP P03089
C	21	MET	-	EXPRESSION TAG	UNP P03089
B	18	GLY	-	EXPRESSION TAG	UNP P03089
B	19	SER	-	EXPRESSION TAG	UNP P03089
B	20	HIS	-	EXPRESSION TAG	UNP P03089
B	21	MET	-	EXPRESSION TAG	UNP P03089
A	18	GLY	-	EXPRESSION TAG	UNP P03089
A	19	SER	-	EXPRESSION TAG	UNP P03089
A	20	HIS	-	EXPRESSION TAG	UNP P03089
A	21	MET	-	EXPRESSION TAG	UNP P03089
D	18	GLY	-	EXPRESSION TAG	UNP P03089
D	19	SER	-	EXPRESSION TAG	UNP P03089
D	20	HIS	-	EXPRESSION TAG	UNP P03089
D	21	MET	-	EXPRESSION TAG	UNP P03089
E	18	GLY	-	EXPRESSION TAG	UNP P03089
E	19	SER	-	EXPRESSION TAG	UNP P03089
E	20	HIS	-	EXPRESSION TAG	UNP P03089

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	21	MET	-	EXPRESSION TAG	UNP P03089

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	5	Total	C	N	O	0	0
			68	37	2	29		
2	B	5	Total	C	N	O	0	0
			68	37	2	29		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	206	Total O 206 206	0	0
5	B	181	Total O 181 181	0	0
5	A	170	Total O 170 170	0	0
5	D	200	Total O 200 200	0	0
5	E	189	Total O 189 189	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Major capsid protein VP1




- Molecule 1: Major capsid protein VP1




- Molecule 1: Major capsid protein VP1



- Molecule 1: Major capsid protein VP1



- Molecule 1: Major capsid protein VP1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.67Å 96.34Å 128.22Å 90.00° 110.28° 90.00°	Depositor
Resolution (Å)	35.27 – 2.00 44.27 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (35.27-2.00) 99.6 (44.27-2.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.178 , 0.206 0.179 , 0.206	Depositor DCC
R_{free} test set	5768 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 114938 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11336	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BGC, NAG, EDO, SIA, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2099	0.50	0/2852
1	B	0.36	0/2083	0.49	0/2830
1	C	0.36	0/2112	0.50	0/2868
1	D	0.36	0/2097	0.51	0/2849
1	E	0.35	0/2100	0.50	0/2852
All	All	0.36	0/10491	0.50	0/14251

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2039	0	2001	5	0
1	B	2023	0	1983	6	0
1	C	2055	0	2016	6	0
1	D	2043	0	1993	6	0
1	E	2040	0	2001	5	0
2	B	68	0	58	0	0
2	C	68	0	58	0	0
3	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	D	6	0	8	0	0
3	E	6	0	8	0	0
4	A	8	0	12	0	0
4	C	4	0	6	0	0
4	D	4	0	6	0	0
4	E	8	0	12	0	0
5	A	170	0	0	0	0
5	B	181	0	0	0	0
5	C	206	0	0	1	0
5	D	200	0	0	0	0
5	E	189	0	0	0	0
All	All	11336	0	10186	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:25:GLU:HB3	1:C:288:LYS:HB3	1.88	0.56
1:E:25:GLU:HB3	1:E:288:LYS:HB3	1.94	0.49
1:A:25:GLU:HB3	1:A:288:LYS:HB3	1.93	0.49
1:D:157:LEU:O	1:D:179:GLN:HA	2.14	0.48
1:B:30:LYS:HD3	1:B:100:LEU:HD22	1.96	0.48
1:B:112:ILE:HD11	1:B:277:TYR:HB2	1.94	0.47
1:E:157:LEU:O	1:E:179:GLN:HA	2.16	0.46
1:D:103:GLU:OE1	1:D:285:ARG:HD2	2.16	0.46
1:C:46:GLU:HG3	5:C:334:HOH:O	2.15	0.46
1:D:27:LEU:HD21	1:D:288:LYS:HB2	1.99	0.45
1:C:220:GLU:H	1:C:220:GLU:HG3	1.47	0.45
1:A:27:LEU:HB3	1:A:28:GLU:H	1.70	0.45
1:D:220:GLU:HG3	1:D:220:GLU:H	1.57	0.44
1:C:157:LEU:O	1:C:179:GLN:HA	2.18	0.44
1:C:103:GLU:OE1	1:C:285:ARG:HD2	2.18	0.44
1:B:30:LYS:HE3	1:B:286:ARG:HB2	2.00	0.43
1:A:29:VAL:HG11	1:A:283:ARG:HD2	2.01	0.43
1:D:171:PRO:HB3	1:D:185:HIS:CG	2.54	0.43
1:B:182:ASN:HB3	1:B:185:HIS:CD2	2.54	0.42
1:B:220:GLU:HG3	1:B:220:GLU:H	1.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:152:GLU:HB3	1:E:206[B]:ARG:CZ	2.51	0.41
1:B:157:LEU:O	1:B:179:GLN:HA	2.20	0.41
1:E:182:ASN:HB3	1:E:185:HIS:CD2	2.56	0.41
1:A:157:LEU:O	1:A:179:GLN:HA	2.20	0.41
1:D:159:ASN:ND2	1:D:161:ARG:H	2.20	0.40
1:A:171:PRO:HB3	1:A:185:HIS:CG	2.56	0.40
1:C:171:PRO:HB3	1:C:185:HIS:CG	2.56	0.40
1:E:152:GLU:OE2	1:E:206[B]:ARG:HD2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/272 (96%)	252 (97%)	8 (3%)	0	100	100
1	B	259/272 (95%)	250 (96%)	9 (4%)	0	100	100
1	C	261/272 (96%)	251 (96%)	10 (4%)	0	100	100
1	D	260/272 (96%)	254 (98%)	6 (2%)	0	100	100
1	E	259/272 (95%)	253 (98%)	6 (2%)	0	100	100
All	All	1299/1360 (96%)	1260 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/237 (98%)	228 (98%)	4 (2%)	68	71
1	B	229/237 (97%)	226 (99%)	3 (1%)	76	79
1	C	233/237 (98%)	230 (99%)	3 (1%)	76	79
1	D	230/237 (97%)	225 (98%)	5 (2%)	60	62
1	E	231/237 (98%)	228 (99%)	3 (1%)	76	79
All	All	1155/1185 (98%)	1137 (98%)	18 (2%)	70	73

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	159	ASN
1	C	216	LEU
1	C	220	GLU
1	B	30	LYS
1	B	159	ASN
1	B	220	GLU
1	A	27	LEU
1	A	67	PHE
1	A	159	ASN
1	A	220	GLU
1	D	76	MET
1	D	94	LEU
1	D	159	ASN
1	D	216	LEU
1	D	220	GLU
1	E	30	LYS
1	E	159	ASN
1	E	220	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	137	GLN
1	C	159	ASN
1	B	159	ASN
1	A	159	ASN
1	D	123	ASN
1	D	137	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	159	ASN
1	E	137	GLN
1	E	159	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SIA	B	1	2	16,20,21	0.41	0	18,28,31	0.47	0
2	GAL	B	2	2	11,11,12	0.62	0	14,15,17	0.70	0
2	NAG	B	3	2	14,14,15	0.51	0	15,19,21	0.79	1 (6%)
2	GAL	B	4	2	11,11,12	0.55	0	14,15,17	0.92	1 (7%)
2	BGC	B	5	2	12,12,12	0.52	0	17,17,17	0.72	0
2	SIA	C	1	2	16,20,21	0.41	0	18,28,31	0.44	0
2	GAL	C	2	2	11,11,12	0.65	0	14,15,17	0.69	0
2	NAG	C	3	2	14,14,15	0.47	0	15,19,21	0.84	1 (6%)
2	GAL	C	4	2	11,11,12	0.59	0	14,15,17	0.94	1 (7%)
2	BGC	C	5	2	12,12,12	0.51	0	17,17,17	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SIA	B	1	2	-	0/14/34/38	0/1/1/1
2	GAL	B	2	2	-	0/2/19/22	0/1/1/1
2	NAG	B	3	2	-	0/6/23/26	0/1/1/1
2	GAL	B	4	2	-	0/2/19/22	0/1/1/1
2	BGC	B	5	2	-	0/2/22/22	0/1/1/1
2	SIA	C	1	2	-	0/14/34/38	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	NAG	C	3	2	-	0/6/23/26	0/1/1/1
2	GAL	C	4	2	-	0/2/19/22	0/1/1/1
2	BGC	C	5	2	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	NAG	C1-O5-C5	2.00	114.79	112.25
2	C	3	NAG	C1-O5-C5	2.25	115.11	112.25
2	B	4	GAL	C1-C2-C3	2.26	112.22	109.54
2	C	4	GAL	C1-C2-C3	2.69	112.72	109.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	4	-	5,5,5	0.35	0	5,5,5	0.32	0
4	EDO	A	5	-	3,3,3	0.47	0	2,2,2	0.45	0
4	EDO	A	6	-	3,3,3	0.47	0	2,2,2	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	290	-	5,5,5	0.34	0	5,5,5	0.28	0
3	GOL	C	290	-	5,5,5	0.36	0	5,5,5	0.26	0
4	EDO	C	291	-	3,3,3	0.48	0	2,2,2	0.48	0
3	GOL	D	2	-	5,5,5	0.35	0	5,5,5	0.26	0
4	EDO	D	290	-	3,3,3	0.48	0	2,2,2	0.48	0
4	EDO	E	290	-	3,3,3	0.47	0	2,2,2	0.47	0
3	GOL	E	3	-	5,5,5	0.40	0	5,5,5	0.31	0
4	EDO	E	4	-	3,3,3	0.49	0	2,2,2	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	4	-	-	0/4/4/4	0/0/0/0
4	EDO	A	5	-	-	0/1/1/1	0/0/0/0
4	EDO	A	6	-	-	0/1/1/1	0/0/0/0
3	GOL	B	290	-	-	0/4/4/4	0/0/0/0
3	GOL	C	290	-	-	0/4/4/4	0/0/0/0
4	EDO	C	291	-	-	0/1/1/1	0/0/0/0
3	GOL	D	2	-	-	0/4/4/4	0/0/0/0
4	EDO	D	290	-	-	0/1/1/1	0/0/0/0
4	EDO	E	290	-	-	0/1/1/1	0/0/0/0
3	GOL	E	3	-	-	0/4/4/4	0/0/0/0
4	EDO	E	4	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/272 (95%)	-0.37	4 (1%) 76 77	9, 15, 29, 45	0
1	B	258/272 (94%)	-0.23	8 (3%) 52 53	10, 16, 32, 49	0
1	C	261/272 (95%)	-0.40	5 (1%) 70 70	8, 15, 31, 49	0
1	D	261/272 (95%)	-0.34	1 (0%) 93 93	8, 14, 29, 49	0
1	E	258/272 (94%)	-0.44	2 (0%) 87 88	8, 14, 27, 43	0
All	All	1297/1360 (95%)	-0.36	20 (1%) 76 77	8, 15, 30, 49	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	33	VAL	7.3
1	B	33	VAL	5.8
1	D	33	VAL	5.2
1	E	33	VAL	5.0
1	B	91	ASN	4.6
1	B	99	ILE	4.3
1	A	24	VAL	3.7
1	C	289	ASN	3.6
1	C	92	GLU	3.5
1	A	33	VAL	3.5
1	B	89	ASN	3.2
1	B	289	ASN	2.8
1	C	94	LEU	2.7
1	B	90	LEU	2.5
1	B	288	LYS	2.5
1	A	289	ASN	2.5
1	E	289	ASN	2.5
1	A	65	ASP	2.4
1	C	34	ASP	2.3
1	B	36	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	C	3	14/15	0.92	0.14	2.05	18,21,23,24	0
2	NAG	B	3	14/15	0.95	0.13	1.79	20,22,24,24	0
2	SIA	C	1	20/21	0.97	0.10	0.59	14,15,16,16	0
2	SIA	B	1	20/21	0.98	0.07	-1.09	15,18,18,18	0
2	GAL	C	2	11/12	0.96	0.18	-	18,20,21,22	0
2	GAL	B	2	11/12	0.96	0.08	-	18,20,22,23	0
2	BGC	B	5	12/12	0.71	0.29	-	35,42,44,44	0
2	GAL	C	4	11/12	0.91	0.17	-	25,27,29,30	0
2	BGC	C	5	12/12	0.80	0.27	-	34,40,41,41	0
2	GAL	B	4	11/12	0.94	0.13	-	26,28,30,31	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	C	291	4/4	0.88	0.21	14.44	37,37,37,39	0
4	EDO	D	290	4/4	0.82	0.23	12.02	43,43,43,43	0
4	EDO	A	5	4/4	0.90	0.16	11.12	38,38,39,39	0
4	EDO	E	290	4/4	0.90	0.22	8.78	39,40,40,40	0
3	GOL	D	2	6/6	0.78	0.16	5.20	35,36,37,38	0
3	GOL	E	3	6/6	0.84	0.23	5.18	37,37,38,38	0
3	GOL	B	290	6/6	0.82	0.17	4.43	42,43,43,43	0
3	GOL	C	290	6/6	0.89	0.15	3.58	31,33,35,37	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	4	6/6	0.87	0.20	2.77	34,35,35,36	0
4	EDO	A	6	4/4	0.88	0.18	2.12	27,29,29,29	0
4	EDO	E	4	4/4	0.81	0.16	-	49,49,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.