



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:13 PM GMT

PDB ID : 4O4L
Title : Tubulin-Peloruside A-Epothilone A complex
Authors : Prota, A.E.; Bargsten, K.; Northcote, P.T.; Marsh, M.; Altmann, K.H.; Miller, J.H.; Diaz, J.F.; Steinmetz, M.O.
Deposited on : 2013-12-18
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	 90% 7%
1	C	451	 89% 8%
2	B	445	 87% 9%
2	D	445	 86% 11%
3	E	143	 82% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	384	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	EP	D	504	-	-	-	X

2 Entry composition i

There are 12 unique types of molecules in this entry. The entry contains 18663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	439	Total	C	N	O	S	0	13	0
			3489	2217	585	662	25			
1	C	440	Total	C	N	O	S	0	22	0
			3536	2245	588	676	27			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	428	Total	C	N	O	S	0	15	0
			3435	2164	580	662	29			
2	D	431	Total	C	N	O	S	0	6	0
			3411	2143	580	660	28			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	123	Total	C	N	O	S	0	6	0
			1046	647	188	205	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	ILE	CLONING ARTIFACT	UNP P63043
E	4	ALA	SER	CLONING ARTIFACT	UNP P63043


- Molecule 4 is a protein called Tubulin-tyrosine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	F	351	Total	C	N	O	S	0	9	0
			2902	1870	490	528	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	380	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	381	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	382	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	383	HIS	-	EXPRESSION TAG	UNP E1BQ43
F	384	HIS	-	EXPRESSION TAG	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	B	1	Total	Mg	0	0
			1	1		
6	A	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	2	Total Ca 2 2	0	0
7	A	1	Total Ca 1 1	0	0

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C N O P 28 10 5 11 2	0	0
8	D	1	Total C N O P 28 10 5 11 2	0	0

- Molecule 9 is PELORUSIDE A (three-letter code: POU) (formula: C₂₇H₄₈O₁₁).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			38	27	11		
9	D	1	Total	C	O	0	0
			38	27	11		

- Molecule 10 is EPOTHILONE A (three-letter code: EP) (formula: $C_{26}H_{39}NO_6S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
10	B	1	Total	C	N	O	S	0	0
			34	26	1	6	1		
10	D	1	Total	C	N	O	S	0	0
			34	26	1	6	1		

- Molecule 11 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
11	B	1	12	6	1	4	1	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	130	Total	O	0	0
			130	130		
12	B	127	Total	O	0	0
			127	127		
12	C	204	Total	O	0	0
			204	204		
12	D	44	Total	O	0	0
			44	44		
12	E	30	Total	O	0	0
			30	30		
12	F	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain



- Molecule 2: Tubulin beta-2B chain




- Molecule 2: Tubulin beta-2B chain







● Molecule 3: Stathmin-4



● Molecule 4: Tubulin-tyrosine ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.40Å 156.33Å 180.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.10 – 2.20 78.23 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (59.10-2.20) 99.9 (78.23-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.2_1309)	Depositor
R, R_{free}	0.186 , 0.224 0.197 , 0.230	Depositor DCC
R_{free} test set	7516 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	53.7	Xtrriage
Anisotropy	0.217	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	2 of 149694 reflections (0.001%)	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18663	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, CA, POU, GTP, MES, EP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/3606	0.42	0/4896
1	C	0.23	0/3678	0.43	0/4997
2	B	0.23	0/3552	0.42	0/4809
2	D	0.22	0/3504	0.40	0/4748
3	E	0.21	0/1073	0.36	0/1425
4	F	0.21	0/2992	0.41	0/4042
All	All	0.22	0/18405	0.41	0/24917

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3489	0	3442	18	0
1	C	3536	0	3485	22	0
2	B	3435	0	3355	23	0
2	D	3411	0	3306	28	0
3	E	1046	0	1072	3	0
4	F	2902	0	2921	33	0
5	A	32	0	12	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
8	B	28	0	12	0	0
8	D	28	0	12	1	0
9	B	38	0	48	1	0
9	D	38	0	48	3	0
10	B	34	0	39	1	0
10	D	34	0	39	2	0
11	B	12	0	12	2	0
12	A	130	0	0	0	0
12	B	127	0	0	2	0
12	C	204	0	0	2	0
12	D	44	0	0	2	0
12	E	30	0	0	0	0
12	F	26	0	0	0	0
All	All	18663	0	17815	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 128 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:286:LEU:HD13	2:B:373:MET:HG2	1.65	0.76
4:F:138:ARG:NH1	4:F:143:GLU:OE1	2.20	0.74
2:D:240[B]:THR:HG21	2:D:320:ARG:HD2	1.69	0.73
2:D:311:ARG:NH1	2:D:436:GLN:O	2.22	0.73
2:B:90:ASP:OD1	2:B:90:ASP:N	2.21	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/451 (100%)	436 (97%)	14 (3%)	0	100	100
1	C	460/451 (102%)	452 (98%)	8 (2%)	0	100	100
2	B	441/445 (99%)	427 (97%)	14 (3%)	0	100	100
2	D	435/445 (98%)	420 (97%)	14 (3%)	1 (0%)	52	59
3	E	125/143 (87%)	120 (96%)	5 (4%)	0	100	100
4	F	353/384 (92%)	314 (89%)	35 (10%)	4 (1%)	17	14
All	All	2264/2319 (98%)	2169 (96%)	90 (4%)	5 (0%)	52	59

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	F	156	LYS
4	F	243	HIS
4	F	155	ALA
4	F	232	ASN
2	D	73	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/379 (101%)	378 (99%)	5 (1%)	76	87
1	C	393/379 (104%)	387 (98%)	6 (2%)	72	84
2	B	385/383 (100%)	373 (97%)	12 (3%)	47	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	378/383 (99%)	370 (98%)	8 (2%)	61	74
3	E	116/127 (91%)	113 (97%)	3 (3%)	54	66
4	F	322/342 (94%)	309 (96%)	13 (4%)	38	47
All	All	1977/1993 (99%)	1930 (98%)	47 (2%)	57	69

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	347[B]	CYS
2	D	110	GLU
4	F	258	GLU
2	D	1	MET
2	D	139	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	301	GLN
2	B	293	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 7 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GTP	A	501	6	25,34,34	0.95	1 (4%)	34,54,54	1.62	5 (14%)
8	GDP	B	501	6	23,30,30	1.20	2 (8%)	30,47,47	1.76	7 (23%)
9	POU	B	503	-	36,39,39	2.04	4 (11%)	33,57,57	1.88	9 (27%)
10	EP	B	504	-	32,36,36	1.28	3 (9%)	35,53,53	2.42	11 (31%)
11	MES	B	507	-	11,12,12	0.61	0	14,16,16	2.29	3 (21%)
5	GTP	C	501	6	25,34,34	0.94	1 (4%)	34,54,54	1.63	5 (14%)
8	GDP	D	501	6	23,30,30	1.18	2 (8%)	30,47,47	1.87	8 (26%)
9	POU	D	503	-	36,39,39	2.22	6 (16%)	33,57,57	2.17	9 (27%)
10	EP	D	504	-	32,36,36	1.30	4 (12%)	35,53,53	2.36	11 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	501	6	-	0/18/38/38	0/3/3/3
8	GDP	B	501	6	-	0/12/32/32	0/3/3/3
9	POU	B	503	-	-	1/54/76/76	0/0/2/2
10	EP	B	504	-	-	1/49/55/55	0/1/3/3
11	MES	B	507	-	-	0/6/14/14	0/1/1/1
5	GTP	C	501	6	-	0/18/38/38	0/3/3/3
8	GDP	D	501	6	-	0/12/32/32	0/3/3/3
9	POU	D	503	-	-	1/54/76/76	0/0/2/2
10	EP	D	504	-	-	1/49/55/55	0/1/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	D	503	POU	C2-C1	-4.85	1.38	1.52
9	B	503	POU	C2-C1	-4.68	1.39	1.52
9	D	503	POU	C12-C11	-2.21	1.50	1.53
10	D	504	EP	C59-C57	-2.01	1.51	1.54
10	B	504	EP	C13-S1	2.01	1.73	1.70

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	D	503	POU	C13-C12-C11	-7.83	104.39	114.35
10	B	504	EP	C32-C27-C24	-6.69	114.36	123.40
10	D	504	EP	C32-C27-C24	-6.40	114.75	123.40
5	C	501	GTP	N3-C2-N1	-5.06	119.73	127.44
5	A	501	GTP	N3-C2-N1	-4.79	120.14	127.44

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	B	504	EP	C5-C10-C12-N20
10	D	504	EP	C5-C10-C12-N20
9	B	503	POU	C23-C22-C21-C15
9	D	503	POU	C23-C22-C21-C15

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	501	GTP	1	0
9	B	503	POU	1	0
10	B	504	EP	1	0
11	B	507	MES	2	0
8	D	501	GDP	1	0
9	D	503	POU	3	0
10	D	504	EP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/451 (97%)	0.04	3 (0%) 89 88	31, 48, 83, 158	0
1	C	440/451 (97%)	0.11	2 (0%) 91 91	25, 38, 66, 97	0
2	B	428/445 (96%)	0.22	9 (2%) 67 65	25, 44, 83, 159	2 (0%)
2	D	431/445 (96%)	0.26	16 (3%) 45 44	35, 67, 103, 140	6 (1%)
3	E	123/143 (86%)	0.40	6 (4%) 33 33	38, 65, 107, 137	0
4	F	351/384 (91%)	2.58	173 (49%) 0 0	51, 110, 185, 210	0
All	All	2212/2319 (95%)	0.56	209 (9%) 11 10	25, 54, 140, 210	8 (0%)

The worst 5 of 209 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	133	ALA	16.3
4	F	173	ILE	15.2
4	F	89	GLU	12.3
4	F	177	GLY	12.0
2	B	280	SER	11.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
10	EP	D	504	34/34	0.90	0.25	2.61	49,67,73,82	0
9	POU	D	503	38/38	0.88	0.30	1.33	43,84,113,114	0
7	CA	A	503	1/1	0.98	0.15	1.16	66,66,66,66	0
8	GDP	B	501	28/28	0.98	0.19	0.73	23,29,36,39	0
11	MES	B	507	12/12	0.94	0.14	0.23	51,61,73,81	0
5	GTP	C	501	32/32	0.98	0.14	0.16	17,27,40,53	0
9	POU	B	503	38/38	0.92	0.13	0.08	37,56,88,96	0
5	GTP	A	501	32/32	0.98	0.14	-0.09	24,32,39,49	0
10	EP	B	504	34/34	0.87	0.17	-0.27	53,68,78,108	0
8	GDP	D	501	28/28	0.96	0.14	-0.46	40,59,72,75	0
6	MG	D	502	1/1	0.97	0.06	-	64,64,64,64	0
6	MG	C	502	1/1	0.90	0.19	-	36,36,36,36	0
6	MG	B	502	1/1	0.96	0.23	-	24,24,24,24	0
7	CA	B	505	1/1	0.75	0.20	-	97,97,97,97	0
6	MG	A	502	1/1	0.98	0.13	-	33,33,33,33	0
7	CA	B	506	1/1	0.80	0.21	-	99,99,99,99	0

6.5 Other polymers [i](#)

There are no such residues in this entry.