



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:46 AM GMT

PDB ID : 2O60  
Title : Calmodulin bound to peptide from neuronal nitric oxide synthase  
Authors : Ng, H.L.; Alber, T.; Wand, A.J.  
Deposited on : 2006-12-06  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1665 (1.58-1.54)
Clashscore	102246	1014 (1.56-1.56)
Ramachandran outliers	100387	1704 (1.58-1.54)
Sidechain outliers	100360	1702 (1.58-1.54)
RSRZ outliers	91569	1668 (1.58-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	<div> <div>49%</div> <div>90%</div> <div>9%</div> <div>..</div> </div>
2	B	24	<div> <div>46%</div> <div>83%</div> <div>13%</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	0	7	0
			1188	728	190	260	10			

- Molecule 2 is a protein called Peptide corresponding to calmodulin binding domain of neuronal nitric oxide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	21	Total	C	N	O	S	0	2	1
			156	103	28	24	1			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Ca	0	0
			4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	155	Total	O	0	0
			155	155		
4	B	10	Total	O	0	0
			10	10		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Calmodulin



#### • Molecule 2: Peptide corresponding to calmodulin binding domain of neuronal nitric oxide synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.13Å 32.93Å 73.83Å 90.00° 93.30° 90.00°	Depositor
Resolution (Å)	74.54 – 1.55 10.43 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.4 (74.54-1.55) 99.8 (10.43-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.58 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.3.0021	Depositor
R, $R_{free}$	0.202 , 0.244 0.215 , 0.253	Depositor DCC
$R_{free}$ test set	1208 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.5	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 23602 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	1513	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/1237	0.85	0/1654
2	B	0.74	0/168	0.65	0/218
All	All	0.75	0/1405	0.83	0/1872

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1188	0	1115	11	0
2	B	156	0	175	4	0
3	A	4	0	0	0	0
4	A	155	0	0	2	0
4	B	10	0	0	0	0
All	All	1513	0	1290	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13[B]:LYS:NZ	4:A:457:HOH:O	1.86	1.08
2:B:15[B]:LYS:NZ	2:B:15[B]:LYS:HB2	1.97	0.77
1:A:66:PRO:O	1:A:70[A]:THR:HG23	1.85	0.77
2:B:15[B]:LYS:HZ2	2:B:15[B]:LYS:HB2	1.51	0.75
1:A:85:ILE:HG23	1:A:141:PHE:HE2	1.50	0.74
1:A:85:ILE:HG23	1:A:141:PHE:CE2	2.34	0.63
1:A:78:ASP:O	1:A:79:THR:HG22	2.02	0.59
1:A:79:THR:HG23	1:A:80:ASP:N	2.26	0.50
1:A:81:SER:OG	1:A:82:GLU:N	2.49	0.46
1:A:75:LYS:HE3	4:A:461:HOH:O	2.16	0.44
1:A:79:THR:HG23	1:A:80:ASP:HB2	2.01	0.42
1:A:1:ALA:O	1:A:2:ASP:C	2.58	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/148 (103%)	145 (95%)	5 (3%)	2 (1%)	15	1
2	B	21/24 (88%)	20 (95%)	1 (5%)	0	100	100
All	All	173/172 (101%)	165 (95%)	6 (4%)	2 (1%)	16	2

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	ASP
1	A	78	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/126 (106%)	130 (98%)	3 (2%)	58	25
2	B	16/17 (94%)	16 (100%)	0	100	100
All	All	149/143 (104%)	146 (98%)	3 (2%)	61	31

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	ASP
1	A	83	GLU
1	A	84	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
2	B	23	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	148/148 (100%)	2.64	72 (48%) <b>0</b> <b>0</b>	30, 36, 52, 61	7 (4%)
2	B	20/24 (83%)	2.69	11 (55%) <b>0</b> <b>0</b>	32, 35, 41, 41	2 (10%)
All	All	168/172 (97%)	2.65	83 (49%) <b>0</b> <b>0</b>	30, 36, 51, 61	9 (5%)

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	81	SER	13.2
1	A	125[A]	ILE	11.2
1	A	78	ASP	10.5
1	A	142	VAL	8.9
2	B	4	ALA	8.9
1	A	79	THR	7.7
1	A	138	TYR	7.4
1	A	86	ARG	7.2
1	A	77	LYS	7.2
1	A	85	ILE	6.9
1	A	72[A]	MET	6.6
1	A	80	ASP	6.4
1	A	1	ALA	6.4
1	A	84	GLU	6.1
1	A	13[A]	LYS	5.9
1	A	148[A]	LYS	5.4
2	B	15[A]	LYS	5.3
1	A	130	ILE	5.3
2	B	23	GLN	5.1
2	B	19[A]	LYS	5.1
1	A	83	GLU	4.8
1	A	87	GLU	4.7
1	A	7[A]	GLU	4.4
1	A	2	ASP	4.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	70[A]	THR	4.2
1	A	76	MET	4.2
1	A	116	LEU	4.2
1	A	139[A]	GLU	4.0
1	A	121	VAL	3.8
1	A	89	PHE	3.7
1	A	147	ALA	3.7
2	B	14	VAL	3.7
1	A	107	HIS	3.7
1	A	52	ILE	3.6
1	A	136	VAL	3.6
1	A	145	MET	3.5
1	A	82	GLU	3.3
1	A	88	ALA	3.3
1	A	100	ILE	3.3
1	A	93	ASP	3.2
1	A	42	ASN	3.2
1	A	16	PHE	3.2
1	A	99	TYR	3.2
1	A	91	VAL	3.1
1	A	111	ASN	3.0
1	A	63	ILE	3.0
1	A	112	LEU	2.9
1	A	90	ARG	2.9
1	A	126	ARG	2.9
1	A	3	GLN	2.9
1	A	15	ALA	2.8
1	A	123	GLU	2.8
1	A	108	VAL	2.8
1	A	27	ILE	2.8
1	A	110	THR	2.7
1	A	143	GLN	2.7
1	A	39	LEU	2.7
1	A	48	LEU	2.7
1	A	8	GLN	2.6
1	A	127	GLU	2.5
1	A	146	THR	2.5
1	A	46	ALA	2.5
1	A	33	GLY	2.4
1	A	35	VAL	2.4
1	A	141	PHE	2.4
1	A	19	PHE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	115	LYS	2.3
1	A	132	GLY	2.3
1	A	62	THR	2.3
2	B	22	GLY	2.3
1	A	20	ASP	2.2
1	A	117	THR	2.2
1	A	55	VAL	2.2
1	A	94	LYS	2.2
2	B	5	ILE	2.2
2	B	20	LEU	2.2
1	A	119	GLU	2.2
2	B	16	PHE	2.2
2	B	8	LYS	2.1
1	A	5	THR	2.1
1	A	73	ALA	2.1
2	B	7	PHE	2.1
1	A	34	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CA	A	402	1/1	0.99	0.19	1.26	32,32,32,32	0
3	CA	A	403	1/1	0.99	0.14	-0.41	32,32,32,32	0
3	CA	A	404	1/1	0.96	0.17	-0.53	34,34,34,34	0
3	CA	A	401	1/1	0.96	0.14	-1.56	35,35,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.