



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:13 PM GMT

PDB ID : 4O6L
Title : Crystal Structure of TTK kinase domain with an inhibitor: 401498 (N-[(1R)-1-(2-chlorophenyl)propyl]-3-{4-[(1-methylpiperidin-4-yl)oxy]phenyl}-1H-indazole-5-carboxamide)
Authors : Qiu, W.; Plotnikova, O.; Feher, M.; Awrey, D.E.; Chirgadze, N.Y.
Deposited on : 2013-12-22
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	<div> <div>5%</div> <div>68%</div> <div>23%</div> <div>7%</div> </div>
1	B	281	<div> <div>7%</div> <div>72%</div> <div>18%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1002	-	-	X	X
3	GOL	B	1002	-	-	X	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4375 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity protein kinase TTK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	P	S	0	0	0
			2133	1361	352	404	3	13			
1	B	256	Total	C	N	O	S		0	0	0
			2091	1345	348	385	13				

- Molecule 2 is N-[(1R)-1-(2-CHLOROPHENYL)PROPYL]-3-{4-[(1-METHYLPIPERIDI N-4-YL)OXY]PHENYL}-1H-INDAZOLE-5-CARBOXAMIDE (three-letter code: 2QK) (formula: C₂₉H₃₁ClN₄O₂).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total	O	0	0
			32	32		
4	B	35	Total	O	0	0
			35	35		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dual specificity protein kinase TTK



• Molecule 1: Dual specificity protein kinase TTK



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	153.23 Å 70.26 Å 106.89 Å 90.00° 133.14° 90.00°	Depositor
Resolution (Å)	38.27 – 2.38 39.00 – 2.38	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.27-2.38) 94.1 (39.00-2.38)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 2.39 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.210 , 0.250 0.220 , 0.262	Depositor DCC
R_{free} test set	1047 reflections (3.26%)	DCC
Wilson B-factor (Å ²)	54.6	Xtriage
Anisotropy	0.564	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.3	EDS
Estimated twinning fraction	0.001 for h+2*l,k,-h-l 0.009 for h,-k,-h-l 0.387 for -h-2*l,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 33137 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4375	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, GOL, 2QK, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/2147	0.58	0/2901
1	B	0.43	0/2137	0.59	0/2887
All	All	0.44	0/4284	0.58	0/5788

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2133	0	2125	37	0
1	B	2091	0	2106	32	0
2	A	36	0	31	4	0
2	B	36	0	31	2	0
3	A	6	0	8	6	0
3	B	6	0	8	9	0
4	A	32	0	0	0	0
4	B	35	0	0	0	0
All	All	4375	0	4309	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:713:PRO:HB2	3:B:1002:GOL:H2	1.52	0.92
1:A:713:PRO:HB2	3:A:1002:GOL:H2	1.64	0.79
1:B:781:SER:HB2	3:B:1002:GOL:H31	1.62	0.79
1:B:587:ARG:HG2	1:B:587:ARG:HH21	1.58	0.68
1:A:773:LYS:HG3	1:A:779:ARG:HA	1.77	0.66
1:B:585:ILE:O	1:B:587:ARG:NH2	2.31	0.63
1:B:620:ASP:HB3	1:B:623:GLU:HB2	1.81	0.61
1:B:574:TYR:CD1	1:B:643:ILE:HD11	2.36	0.61
1:A:574:TYR:CD1	1:A:643:ILE:HD11	2.36	0.60
1:A:593:ILE:HG23	1:A:598:ILE:HG12	1.82	0.60
1:A:672:GLN:HG2	1:A:673:PRO:HD2	1.84	0.60
1:A:571:GLU:OE2	1:A:665:PHE:N	2.30	0.59
1:B:713:PRO:CB	3:B:1002:GOL:H2	2.31	0.58
1:A:733:PRO:O	1:A:736:GLN:NE2	2.36	0.57
1:B:534:GLY:O	1:B:670:GLN:NE2	2.36	0.57
1:A:609:LEU:HD22	1:A:660:LEU:HD11	1.85	0.56
1:A:613:LEU:HB3	1:A:726:TYR:HE2	1.70	0.56
1:A:781:SER:HB2	3:A:1002:GOL:H31	1.87	0.55
1:B:582:SER:HB2	1:B:637:THR:HG21	1.87	0.54
1:A:783:PRO:HD3	3:A:1002:GOL:H12	1.90	0.54
1:B:782:ILE:HB	3:B:1002:GOL:H12	1.89	0.54
1:A:664:ASP:CG	2:A:1001:2QK:H31	2.28	0.53
1:A:789:PRO:HB3	1:A:793:ILE:HD12	1.90	0.53
1:B:664:ASP:CG	2:B:1001:2QK:H31	2.29	0.53
1:A:537:SER:HB3	1:A:555:VAL:HG13	1.92	0.53
1:B:617:LYS:HD3	1:B:617:LYS:H	1.75	0.51
1:A:676:TPO:O3P	1:B:529:LYS:NZ	2.43	0.51
1:B:587:ARG:HG2	1:B:587:ARG:NH2	2.25	0.50
1:A:675:TPO:HB	2:A:1001:2QK:H20	1.94	0.49
1:B:664:ASP:OD2	2:B:1001:2QK:H31	2.14	0.47
1:B:625:LYS:HE3	1:B:790:TYR:O	2.15	0.47
1:B:788:HIS:CG	1:B:789:PRO:HD2	2.50	0.47
1:B:783:PRO:HD3	3:B:1002:GOL:H11	1.96	0.47
1:B:615:LYS:HB3	1:B:617:LYS:HE2	1.96	0.46
1:A:583:ASP:HA	1:A:587:ARG:HH22	1.80	0.46
1:B:741:ILE:HD12	1:B:741:ILE:H	1.79	0.46
1:A:693:GLU:CG	1:A:776:PRO:HG3	2.46	0.45
1:A:713:PRO:CB	3:A:1002:GOL:H2	2.41	0.45
1:A:577:LYS:NZ	1:A:641:HIS:HD2	2.14	0.45
1:B:782:ILE:H	3:B:1002:GOL:H12	1.82	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:653:PHE:HB3	1:A:660:LEU:HD22	1.98	0.45
1:A:783:PRO:CD	3:A:1002:GOL:H12	2.47	0.45
1:B:686:THR:OG1	1:B:687:VAL:N	2.49	0.45
1:A:670:GLN:HG2	1:A:671:MET:O	2.17	0.44
1:B:763:ASP:HB3	1:B:790:TYR:HB2	2.00	0.44
1:A:693:GLU:HG2	1:A:776:PRO:HG3	1.99	0.44
1:A:544:ASN:OD1	1:A:548:GLN:N	2.39	0.44
1:A:781:SER:CB	3:A:1002:GOL:H31	2.47	0.44
1:A:529:LYS:HE2	1:A:529:LYS:HB2	1.58	0.44
1:B:541:GLN:HG2	1:B:549:ILE:HD11	2.00	0.43
1:A:737:ILE:O	1:A:743:LYS:HE3	2.17	0.43
1:A:539:VAL:HG21	2:A:1001:2QK:H5	2.00	0.43
1:B:781:SER:CB	3:B:1002:GOL:H31	2.42	0.43
1:B:653:PHE:HB3	1:B:660:LEU:HD22	2.00	0.43
1:B:572:ILE:O	1:B:576:ASN:HB2	2.18	0.42
1:B:580:GLN:H	1:B:580:GLN:HG3	1.42	0.42
1:B:749:ASP:HA	1:B:750:PRO:HD2	1.95	0.42
1:A:763:ASP:OD1	1:A:764:LEU:N	2.53	0.42
1:A:629:LYS:HB2	1:A:629:LYS:HE3	1.78	0.41
1:B:783:PRO:HD3	3:B:1002:GOL:C1	2.51	0.41
1:A:567:SER:OG	1:A:668:ALA:HA	2.19	0.41
1:A:660:LEU:HA	1:A:660:LEU:HD23	1.87	0.41
1:A:664:ASP:OD2	2:A:1001:2QK:H31	2.21	0.41
1:A:516:GLU:HB3	1:A:527:ILE:HD12	2.02	0.41
1:A:583:ASP:HA	1:A:587:ARG:NH2	2.36	0.41
1:B:556:ASN:HD21	1:B:558:GLU:HB2	1.85	0.41
1:A:767:VAL:HG23	1:A:788:HIS:CE1	2.56	0.40
1:B:783:PRO:CD	3:B:1002:GOL:H11	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/281 (90%)	249 (99%)	3 (1%)	0	100	100
1	B	250/281 (89%)	246 (98%)	4 (2%)	0	100	100
All	All	502/562 (89%)	495 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/257 (93%)	216 (91%)	22 (9%)	11	15
1	B	236/257 (92%)	215 (91%)	21 (9%)	12	16
All	All	474/514 (92%)	431 (91%)	43 (9%)	12	15

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	529	LYS
1	A	536	SER
1	A	545	GLU
1	A	547	LYS
1	A	559	GLU
1	A	561	ASP
1	A	565	LEU
1	A	566	ASP
1	A	578	LEU
1	A	592	GLU
1	A	593	ILE
1	A	617	LYS
1	A	620	ASP
1	A	622	TRP
1	A	625	LYS
1	A	687	VAL
1	A	699	SER
1	A	736	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	741	ILE
1	A	773	LYS
1	A	782	ILE
1	A	784	GLU
1	B	515	ASN
1	B	516	GLU
1	B	521	LYS
1	B	526	SER
1	B	545	GLU
1	B	559	GLU
1	B	563	GLN
1	B	565	LEU
1	B	566	ASP
1	B	576	ASN
1	B	580	GLN
1	B	613	LEU
1	B	617	LYS
1	B	619	ILE
1	B	622	TRP
1	B	671	MET
1	B	740	GLN
1	B	753	GLU
1	B	762	LYS
1	B	782	ILE
1	B	791	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	581	HIS
1	A	641	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	TPO	A	675	1	8,10,11	1.14	0	7,14,16	0.85	0
1	TPO	A	676	1	8,10,11	1.55	1 (12%)	7,14,16	1.27	1 (14%)
1	SEP	A	677	1	8,9,10	1.61	2 (25%)	8,12,14	1.44	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	675	1	-	1/8/11/13	0/0/0/0
1	TPO	A	676	1	-	0/8/11/13	0/0/0/0
1	SEP	A	677	1	-	0/6/8/10	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	677	SEP	P-O2P	2.07	1.62	1.54
1	A	677	SEP	P-O1P	3.22	1.61	1.51
1	A	676	TPO	P-O1P	3.22	1.61	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	676	TPO	CG2-CB-CA	-2.36	108.37	113.17
1	A	677	SEP	OG-CB-CA	2.99	110.82	108.27

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	675	TPO	OG1-CB-CA-N

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	675	TPO	1	0
1	A	676	TPO	1	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2QK	A	1001	-	37,40,40	1.47	5 (13%)	44,56,56	2.55	12 (27%)
3	GOL	A	1002	-	5,5,5	0.42	0	5,5,5	0.69	0
2	2QK	B	1001	-	37,40,40	1.50	5 (13%)	44,56,56	2.18	11 (25%)
3	GOL	B	1002	-	5,5,5	0.37	0	5,5,5	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2QK	A	1001	-	-	0/22/32/32	0/5/5/5
3	GOL	A	1002	-	-	0/4/4/4	0/0/0/0
2	2QK	B	1001	-	-	0/22/32/32	0/5/5/5
3	GOL	B	1002	-	-	0/4/4/4	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	2QK	C14-C13	-4.09	1.44	1.49
2	B	1001	2QK	C14-C13	-3.33	1.45	1.49
2	B	1001	2QK	C07-C05	2.04	1.54	1.50
2	B	1001	2QK	C09-C08	2.09	1.41	1.36
2	A	1001	2QK	C35-C30	2.12	1.42	1.39
2	A	1001	2QK	C09-C08	2.17	1.41	1.36
2	A	1001	2QK	C29-C07	2.18	1.41	1.37
2	B	1001	2QK	C29-C07	2.89	1.42	1.37
2	A	1001	2QK	C05-N04	4.43	1.44	1.34
2	B	1001	2QK	C05-N04	4.69	1.44	1.34

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	2QK	C31-C30-C03	-4.15	110.58	120.24
2	B	1001	2QK	C34-C35-C30	-3.97	118.33	121.97
2	A	1001	2QK	C34-C35-C30	-3.67	118.61	121.97
2	B	1001	2QK	C31-C30-C03	-3.54	111.99	120.24
2	A	1001	2QK	C17-O18-C19	-3.07	114.14	119.06
2	B	1001	2QK	C17-O18-C19	-2.43	115.17	119.06
2	B	1001	2QK	C08-C09-C10	-2.00	118.71	120.88
2	B	1001	2QK	C02-C03-N04	2.16	114.64	110.78
2	A	1001	2QK	C31-C30-C35	2.33	119.31	116.79
2	B	1001	2QK	C31-C30-C35	2.37	119.36	116.79
2	A	1001	2QK	C02-C03-N04	2.44	115.14	110.78
2	B	1001	2QK	C21-N22-C24	2.75	113.14	109.53
2	A	1001	2QK	C21-C20-C19	2.99	113.13	110.12
2	B	1001	2QK	C23-N22-C21	3.78	116.52	110.63
2	A	1001	2QK	C20-C21-N22	4.01	116.90	111.32
2	A	1001	2QK	C23-N22-C21	4.17	117.13	110.63
2	B	1001	2QK	C30-C35-CL	4.82	125.73	120.42
2	B	1001	2QK	C35-C30-C03	4.99	128.65	121.71
2	A	1001	2QK	C21-N22-C24	5.11	116.23	109.53
2	A	1001	2QK	C30-C35-CL	5.56	126.55	120.42
2	A	1001	2QK	C35-C30-C03	6.04	130.11	121.71
2	B	1001	2QK	C23-N22-C24	7.68	122.61	110.63
2	A	1001	2QK	C23-N22-C24	8.43	123.78	110.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	2QK	4	0
3	A	1002	GOL	6	0
2	B	1001	2QK	2	0
3	B	1002	GOL	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	257/281 (91%)	0.74	15 (5%)	26 31	32, 53, 77, 98	0
1	B	256/281 (91%)	0.74	20 (7%)	16 18	31, 54, 77, 93	0
All	All	513/562 (91%)	0.74	35 (6%)	20 23	31, 53, 77, 98	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	673	PRO	4.6
1	B	695	ILE	4.0
1	A	559	GLU	3.8
1	B	737	ILE	3.7
1	A	741	ILE	3.4
1	A	622	TRP	2.9
1	B	744	LEU	2.9
1	B	572	ILE	2.9
1	A	618	SER	2.8
1	A	726	TYR	2.7
1	A	616	LYS	2.7
1	A	687	VAL	2.7
1	A	555	VAL	2.6
1	B	559	GLU	2.5
1	A	697	ASP	2.5
1	A	655	ILE	2.5
1	B	520	VAL	2.4
1	B	622	TRP	2.4
1	B	617	LYS	2.3
1	B	785	LEU	2.3
1	B	698	MET	2.3
1	B	690	MET	2.2
1	B	593	ILE	2.2
1	B	791	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	738	ILE	2.1
1	A	734	PHE	2.1
1	B	726	TYR	2.1
1	A	695	ILE	2.1
1	B	538	LYS	2.1
1	A	575	LEU	2.1
1	B	736	GLN	2.1
1	A	547	LYS	2.0
1	A	754	ILE	2.0
1	B	748	ILE	2.0
1	B	598	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	675	11/12	0.83	0.16	-	69,79,85,86	0
1	TPO	A	676	11/12	0.92	0.15	-	74,87,90,92	0
1	SEP	A	677	10/11	0.72	0.15	-	87,93,104,110	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	B	1002	6/6	0.68	0.39	10.03	39,45,50,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	1002	6/6	0.82	0.27	4.04	36,42,51,53	0
2	2QK	B	1001	36/36	0.93	0.16	-0.12	32,41,58,61	0
2	2QK	A	1001	36/36	0.95	0.15	-1.18	32,43,58,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.