



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 27, 2016 – 04:48 AM BST

PDB ID : 2O8K
Title : NMR Structure of the Sigma-54 RpoN Domain Bound to the-24 Promoter Element
Authors : Doucleff, M.; Pelton, J.G.; Lee, P.S.; Wemmer, D.E.
Deposited on : 2006-12-12

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 62%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	14	
2	C	14	
3	A	63	

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:16-A:74 (59)	0.19	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 6, 9, 10, 12, 13, 16, 17
2	5, 19, 20
3	15, 18
Single-model clusters	7; 8; 11; 14

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1969 atoms, of which 878 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(*TP*TP*TP*TP*GP*GP*CP*AP*CP*GP*TP*TP*TP*C)-3'.

Mol	Chain	Residues	Atoms						Trace
1	B	14	Total	C	H	N	O	P	0
			444	137	163	43	88	13	

- Molecule 2 is a DNA chain called 5'-D(*GP*AP*AP*AP*CP*GP*TP*GP*CP*CP*AP*AP*AP*A)-3'.

Mol	Chain	Residues	Atoms						Trace
2	C	14	Total	C	H	N	O	P	0
			444	137	157	61	76	13	

- Molecule 3 is a protein called RNA polymerase sigma factor RpoN.

Mol	Chain	Residues	Atoms						Trace
3	A	63	Total	C	H	N	O	S	0
			1081	327	558	98	95	3	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	HIS	-	CLONING ARTIFACT	UNP O66858
A	15	MET	-	CLONING ARTIFACT	UNP O66858

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: 5'-D(*TP*TP*TP*TP*GP*GP*CP*AP*CP*GP*TP*TP*TP*C)-3'

Chain B: 



- Molecule 2: 5'-D(*GP*AP*AP*AP*CP*GP*TP*GP*CP*CP*AP*AP*AP*A)-3'

Chain C: 



- Molecule 3: RNA polymerase sigma factor RpoN

Chain A: 




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 6. Colouring as in section 4.1 above.

- Molecule 1: 5'-D(*TP*TP*TP*TP*GP*GP*CP*AP*CP*GP*TP*TP*TP*C)-3'


Chain B: 



- Molecule 2: 5'-D(*GP*AP*AP*AP*CP*GP*TP*GP*CP*CP*AP*AP*AP*A)-3'



• Molecule 3: RNA polymerase sigma factor RpoN



5 Refinement protocol and experimental data overview

The models were refined using the following method: *docking, rigid-body minimization, simulated annealing*.

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *20 structures with lowest energy and no restraint violations greater than 0.5 Å and 5 degrees for distance and dihedral restraints, respectively*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
X-PLOR	refinement	2.14

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 15083
Number of chemical shift lists	1
Total number of shifts	948
Number of shifts mapped to atoms	948
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	62%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	1.37±0.02	1±2/312 (0.5±0.5%)	1.64±0.01	10±1/480 (2.0±0.2%)
2	C	1.20±0.02	0±0/324 (0.1±0.1%)	1.51±0.02	2±0/498 (0.4±0.1%)
3	A	0.74±0.00	0±0/490 (0.0±0.0%)	0.75±0.01	0±0/652 (0.0±0.0%)
All	All	1.08	34/22520 (0.2%)	1.31	240/32600 (0.7%)

5 of 8 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	111	DT	C5-C7	5.63	1.53	1.50	20	7
2	C	121	DT	C5-C7	5.48	1.53	1.50	4	4
1	B	101	DT	C5-C7	5.45	1.53	1.50	9	5
1	B	112	DT	C5-C7	5.41	1.53	1.50	9	7
1	B	102	DT	C5-C7	5.29	1.53	1.50	20	2

5 of 15 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	C	121	DT	C6-C5-C7	-6.63	118.92	122.90	20	20
1	B	113	DT	C6-C5-C7	-6.19	119.18	122.90	16	20
1	B	103	DT	C6-C5-C7	-6.12	119.23	122.90	19	20
1	B	104	DT	C6-C5-C7	-5.94	119.33	122.90	9	20
2	C	121	DT	C4-C5-C6	5.91	121.55	118.00	12	20

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	281	163	163	12±4
2	C	287	157	157	16±5
3	A	485	515	515	45±5
All	All	21060	16700	16700	1253

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

5 of 284 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
3:A:25:ILE:HD11	3:A:66:LEU:HD13	0.90	1.44	9	17
3:A:25:ILE:HD11	3:A:66:LEU:CD1	0.76	2.11	5	14
3:A:17:THR:HG23	3:A:20:GLU:OE2	0.76	1.81	14	8
3:A:25:ILE:O	3:A:29:VAL:HG22	0.74	1.82	4	1
3:A:23:LYS:C	3:A:24:LEU:HD22	0.72	2.05	20	20

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	59/63 (94%)	45±2 (77±3%)	8±2 (14±3%)	6±1 (10±2%)	2	10
All	All	1180/1260 (94%)	903 (77%)	160 (14%)	117 (10%)	2	10

5 of 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	51	GLY	20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
3	A	67	GLY	20
3	A	53	LYS	20
3	A	40	ASP	20
3	A	17	THR	15

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	53/57 (93%)	46±1 (87±3%)	7±1 (13±3%)	9	50
All	All	1060/1140 (93%)	920 (87%)	140 (13%)	9	50

5 of 20 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	59	VAL	20
3	A	53	LYS	18
3	A	43	ILE	18
3	A	66	LEU	14
3	A	31	ASN	14

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 62% for the well-defined parts and 62% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 15083

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	948
Number of shifts mapped to atoms	948
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	10

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	57	-0.68 ± 0.13	Should be applied
$^{13}\text{C}_\beta$	60	0.02 ± 0.18	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	—
^{15}N	60	-0.23 ± 0.28	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 62%, i.e. 835 atoms were assigned a chemical shift out of a possible 1350. 8 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	226/291 (78%)	116/116 (100%)	53/118 (45%)	57/57 (100%)
Sidechain	386/484 (80%)	246/286 (86%)	139/168 (83%)	1/30 (3%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	12/25 (48%)	12/13 (92%)	0/12 (0%)	0/0 (—%)
Overall	835/1350 (62%)	585/741 (79%)	192/488 (39%)	58/121 (48%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	37	PRO	CD	26.26	55.31 – 45.41	-24.3
1	A	37	PRO	CG	51.55	32.66 – 21.76	22.3
1	A	52	PHE	CA	4.92	70.99 – 45.29	-20.7
1	A	53	LYS	CD	53.04	34.86 – 23.06	20.4
1	A	65	MET	CG	55.27	38.33 – 25.73	18.4
1	A	56	ARG	CA	14.36	68.35 – 45.25	-18.4
1	A	55	ALA	CB	44.86	28.03 – 9.93	14.3
1	A	63	ARG	CB	53.58	39.81 – 21.51	12.5
1	A	68	ILE	CA	35.19	75.08 – 48.18	-9.8
1	A	29	VAL	CA	43.00	76.93 – 48.03	-6.7

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

