



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:33 PM GMT

PDB ID : 3O8A  
Title : Crystal structure of Plasmodium falciparum dihydroorotate dehydrogenase bound with novel Inhibitor Genz667348  
Authors : Deng, X.; Booker, M.L.; Phillips, M.A.  
Deposited on : 2010-08-02  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	<div> <div>12%</div> <div>78%</div> <div>12%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	LDA	A	1004	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3095 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydroorotate dehydrogenase homolog, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	375	2975	1902	498	560	15	0	1	0

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	125	MET	-	EXPRESSION TAG	UNP Q08210
A	126	GLY	-	EXPRESSION TAG	UNP Q08210
A	127	SER	-	EXPRESSION TAG	UNP Q08210
A	128	SER	-	EXPRESSION TAG	UNP Q08210
A	129	HIS	-	EXPRESSION TAG	UNP Q08210
A	130	HIS	-	EXPRESSION TAG	UNP Q08210
A	131	HIS	-	EXPRESSION TAG	UNP Q08210
A	132	HIS	-	EXPRESSION TAG	UNP Q08210
A	133	HIS	-	EXPRESSION TAG	UNP Q08210
A	134	HIS	-	EXPRESSION TAG	UNP Q08210
A	135	SER	-	EXPRESSION TAG	UNP Q08210
A	136	SER	-	EXPRESSION TAG	UNP Q08210
A	137	GLY	-	EXPRESSION TAG	UNP Q08210
A	138	LEU	-	EXPRESSION TAG	UNP Q08210
A	139	VAL	-	EXPRESSION TAG	UNP Q08210
A	140	PRO	-	EXPRESSION TAG	UNP Q08210
A	141	ARG	-	EXPRESSION TAG	UNP Q08210
A	142	GLY	-	EXPRESSION TAG	UNP Q08210
A	143	SER	-	EXPRESSION TAG	UNP Q08210
A	144	HIS	-	EXPRESSION TAG	UNP Q08210
A	145	MET	-	EXPRESSION TAG	UNP Q08210
A	146	ALA	-	EXPRESSION TAG	UNP Q08210
A	147	SER	-	EXPRESSION TAG	UNP Q08210
A	148	MET	-	EXPRESSION TAG	UNP Q08210
A	149	THR	-	EXPRESSION TAG	UNP Q08210
A	150	GLY	-	EXPRESSION TAG	UNP Q08210
A	151	GLY	-	EXPRESSION TAG	UNP Q08210

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	152	GLN	-	EXPRESSION TAG	UNP Q08210
A	153	GLN	-	EXPRESSION TAG	UNP Q08210
A	154	GLY	-	EXPRESSION TAG	UNP Q08210
A	155	ARG	-	EXPRESSION TAG	UNP Q08210
A	156	ASP	-	EXPRESSION TAG	UNP Q08210
A	157	PRO	-	EXPRESSION TAG	UNP Q08210

- Molecule 2 is N-CYCLOPROPYL-5-[2-METHYL-5-(TRIFLUOROMETHOXY)-1H-BENZIMIDAZOL-1-YL]THIOPHENE-2-CARBOXAMIDE (three-letter code: O8A) (formula:  $C_{17}H_{14}F_3N_3O_2S$ ).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			26	17	3	3	2	1		

- Molecule 3 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula:  $C_{17}H_{21}N_4O_9P$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 4 is OROTIC ACID (three-letter code: ORO) (formula:  $C_5H_4N_2O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	5	2	4		

- Molecule 5 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula:  $\text{C}_{14}\text{H}_{31}\text{NO}$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			16	14	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	36	Total	O	0	0
			36	36		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase homolog, mitochondrial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.29 Å 85.29 Å 138.72 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.30 39.19 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (40.00-2.30) 99.6 (39.19-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 2.29 Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.205 , 0.229 0.219 , 0.242	Depositor DCC
$R_{free}$ test set	1295 reflections (5.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	63.4	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 54.5	EDS
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 25407 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3095	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ORO, LDA, O8A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	2/3030 (0.1%)	0.73	2/4080 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	283	CYS	CB-SG	-6.95	1.70	1.82
1	A	530	CYS	CB-SG	-5.77	1.72	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296	ARG	NE-CZ-NH2	-8.80	115.90	120.30
1	A	296	ARG	NE-CZ-NH1	6.68	123.64	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2975	0	3018	35	0
2	A	26	0	14	1	0
3	A	31	0	19	1	0
4	A	11	0	3	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	16	0	31	5	0
6	A	36	0	0	2	0
All	All	3095	0	3085	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1004:LDA:C9	5:A:1004:LDA:H51	1.81	1.08
5:A:1004:LDA:H51	5:A:1004:LDA:H92	1.39	1.03
1:A:162:ASN:OD1	1:A:163:PRO:HD2	1.72	0.89
1:A:326:LYS:HG2	1:A:370:GLU:HG3	1.63	0.80
1:A:191:LEU:HD22	1:A:196:ILE:HD11	1.68	0.74
1:A:268:GLU:OE1	1:A:268:GLU:C	2.30	0.70
1:A:564:ARG:O	1:A:565:LYS:C	2.30	0.70
1:A:537:LYS:HE2	1:A:541:GLN:NE2	2.07	0.69
1:A:222:GLY:HA3	1:A:244:PHE:CE2	2.30	0.67
1:A:537:LYS:HE2	1:A:541:GLN:HE22	1.60	0.66
1:A:162:ASN:OD1	1:A:163:PRO:CD	2.46	0.64
1:A:326:LYS:HG2	1:A:370:GLU:CG	2.27	0.64
5:A:1004:LDA:C9	5:A:1004:LDA:C5	2.61	0.57
1:A:223:VAL:HG22	1:A:245:ILE:HG13	1.89	0.54
1:A:162:ASN:ND2	1:A:164:GLU:HB2	2.22	0.53
1:A:244:PHE:HB3	1:A:307:ILE:HB	1.92	0.50
1:A:357:ALA:HB2	1:A:441:GLU:HG2	1.93	0.50
1:A:489:ILE:HG23	1:A:502:ILE:HD13	1.93	0.50
1:A:268:GLU:OE1	1:A:268:GLU:O	2.30	0.49
1:A:226:GLY:HA3	3:A:1002:FMN:N5	2.27	0.49
5:A:1004:LDA:H21	5:A:1004:LDA:HM11	1.43	0.48
1:A:222:GLY:HA3	1:A:244:PHE:CD2	2.48	0.47
1:A:348:THR:HG23	6:A:630:HOH:O	2.13	0.47
1:A:485:SER:OG	1:A:517:LYS:NZ	2.49	0.46
1:A:564:ARG:O	1:A:564:ARG:HG2	2.16	0.45
1:A:349:PRO:HA	1:A:350:GLY:HA2	1.81	0.45
1:A:457:SER:N	1:A:505:SER:O	2.43	0.45
1:A:463:ILE:HD13	1:A:480:LYS:HB3	1.98	0.45
1:A:382:ASP:O	1:A:383:GLU:C	2.54	0.44
1:A:326:LYS:CG	1:A:370:GLU:HG3	2.41	0.44
1:A:415:LEU:HD23	1:A:416:TRP:NE1	2.32	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:PHE:CE2	4:A:1003:ORO:H5	2.53	0.43
1:A:163:PRO:O	1:A:166:PHE:HB3	2.18	0.43
1:A:229:LYS:NZ	6:A:621:HOH:O	2.50	0.43
1:A:255:GLN:OE1	1:A:316:LYS:N	2.46	0.42
1:A:414:PHE:N	1:A:414:PHE:CD1	2.87	0.42
5:A:1004:LDA:H91	5:A:1004:LDA:C5	2.48	0.42
1:A:263:ILE:HG21	2:A:1001:O8A:H16A	2.02	0.41
1:A:247:ILE:HD11	1:A:310:VAL:HG22	2.01	0.41
1:A:364:ILE:O	1:A:368:LYS:HG3	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/415 (90%)	361 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/365 (91%)	330 (99%)	3 (1%)	84	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	235	ASP
1	A	255	GLN
1	A	316	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	195	ASN
1	A	330	ASN
1	A	450	ASN
1	A	464	ASN
1	A	541	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	O8A	A	1001	-	20,29,29	1.67	8 (40%)	21,44,44	2.52	6 (28%)
3	FMN	A	1002	-	32,33,33	1.34	5 (15%)	34,50,50	2.21	7 (20%)
4	ORO	A	1003	-	6,11,11	1.07	1 (16%)	6,15,15	3.65	3 (50%)
5	LDA	A	1004	-	15,15,15	5.15	4 (26%)	16,17,17	0.87	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	O8A	A	1001	-	-	0/8/19/19	0/3/4/4
3	FMN	A	1002	-	-	0/18/18/18	0/3/3/3
4	ORO	A	1003	-	-	0/0/4/4	0/1/1/1
5	LDA	A	1004	-	-	0/13/13/13	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1004	LDA	O1-N1	-18.65	1.21	1.39
5	A	1004	LDA	C1-N1	-5.47	1.41	1.51
5	A	1004	LDA	CM1-N1	-3.35	1.44	1.49
5	A	1004	LDA	CM2-N1	-2.95	1.44	1.49
2	A	1001	O8A	C10-S1	-2.92	1.68	1.74
2	A	1001	O8A	C2-N2	-2.85	1.35	1.39
2	A	1001	O8A	C6-C1	-2.52	1.37	1.41
2	A	1001	O8A	O2-C14	-2.39	1.18	1.23
2	A	1001	O8A	C3-C2	-2.15	1.36	1.41
4	A	1003	ORO	C2-N1	-2.08	1.33	1.38
2	A	1001	O8A	C10-N2	-2.05	1.42	1.46
3	A	1002	FMN	C4-C4A	-2.02	1.37	1.41
2	A	1001	O8A	C15-N3	-2.02	1.42	1.46
2	A	1001	O8A	C2-C1	2.25	1.44	1.40
3	A	1002	FMN	C4-N3	2.58	1.37	1.33
3	A	1002	FMN	C5A-N5	2.72	1.39	1.35
3	A	1002	FMN	C1'-N10	2.82	1.51	1.48
3	A	1002	FMN	C4A-N5	3.66	1.39	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	O8A	C16-C15-N3	-6.57	108.55	118.71
2	A	1001	O8A	C17-C15-N3	-5.91	109.57	118.71
3	A	1002	FMN	N3-C2-N1	-5.38	118.63	127.69
4	A	1003	ORO	C5-C4-N3	-4.61	119.15	124.02
4	A	1003	ORO	N3-C2-N1	-3.67	121.51	127.69
3	A	1002	FMN	C4A-C4-N3	-2.83	119.83	123.52
2	A	1001	O8A	C6-C1-C2	-2.48	118.56	121.10
5	A	1004	LDA	O1-N1-C1	-2.20	107.79	110.27
3	A	1002	FMN	C4A-C10-N10	-2.13	118.97	120.52
2	A	1001	O8A	C6-C1-N1	2.01	136.63	130.72
3	A	1002	FMN	C4A-N5-C5A	2.07	119.16	116.72
3	A	1002	FMN	C1'-N10-C9A	2.18	121.36	118.83
2	A	1001	O8A	C11-C14-N3	2.34	120.32	115.09
3	A	1002	FMN	C5A-C9A-N10	5.02	121.34	117.58
2	A	1001	O8A	C9-C7-N1	5.26	134.19	124.12
4	A	1003	ORO	C4-N3-C2	6.53	121.09	114.21
3	A	1002	FMN	C4-N3-C2	8.19	121.99	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	O8A	1	0
3	A	1002	FMN	1	0
4	A	1003	ORO	1	0
5	A	1004	LDA	5	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	375/415 (90%)	0.60	48 (12%) 5 7	23, 74, 106, 117	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	161	TYR	9.4
1	A	551	TYR	5.6
1	A	223	VAL	5.3
1	A	525	CYS	5.2
1	A	556	TYR	5.1
1	A	455	ILE	4.7
1	A	225	ALA	4.7
1	A	224	ALA	4.6
1	A	564	ARG	4.2
1	A	222	GLY	3.9
1	A	503	ILE	3.7
1	A	504	ALA	3.7
1	A	464	ASN	3.7
1	A	350	GLY	3.7
1	A	340	ALA	3.6
1	A	527	LEU	3.4
1	A	311	SER	3.3
1	A	245	ILE	3.2
1	A	310	VAL	3.1
1	A	309	GLY	3.1
1	A	244	PHE	2.9
1	A	249	THR	2.9
1	A	526	GLN	2.8
1	A	349	PRO	2.8
1	A	162	ASN	2.8
1	A	316	LYS	2.8
1	A	524	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	227	PHE	2.7
1	A	307	ILE	2.7
1	A	353	ASP	2.6
1	A	247	ILE	2.6
1	A	505	SER	2.6
1	A	226	GLY	2.6
1	A	469	PHE	2.6
1	A	456	ILE	2.5
1	A	166	PHE	2.5
1	A	338	TYR	2.4
1	A	207	TYR	2.4
1	A	339	ILE	2.3
1	A	454	MET	2.3
1	A	467	LYS	2.3
1	A	528	TYR	2.3
1	A	427	PHE	2.2
1	A	308	VAL	2.1
1	A	552	GLN	2.1
1	A	444	ASP	2.0
1	A	246	GLU	2.0
1	A	238	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	LDA	A	1004	16/16	0.81	0.27	6.63	72,86,97,97	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FMN	A	1002	31/31	0.97	0.31	0.94	58,64,73,75	0
4	ORO	A	1003	11/11	0.96	0.15	0.09	55,59,66,69	0
2	O8A	A	1001	26/26	0.97	0.09	-0.96	52,57,67,69	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.