



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:17 PM GMT

PDB ID : 1O9I
Title : CRYSTAL STRUCTURE OF THE Y42F MUTANT OF MANGANESE CATALASE FROM LACTOBACILLUS PLANTARUM AT 1.33Å RESOLUTION
Authors : Barynin, V.V.; Whittaker, M.M.; Whittaker, J.W.
Deposited on : 2002-12-13
Resolution : 1.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1806 (1.38-1.30)
Ramachandran outliers	100387	1749 (1.38-1.30)
Sidechain outliers	100360	1749 (1.38-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	266	
1	B	266	
1	C	266	
1	D	266	
1	E	266	
1	F	266	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14325 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PSEUDOCATALASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	9	0
			2123	1322	364	419	18			
1	B	266	Total	C	N	O	S	0	6	0
			2107	1310	363	416	18			
1	C	266	Total	C	N	O	S	0	13	0
			2140	1329	367	426	18			
1	D	266	Total	C	N	O	S	0	14	0
			2146	1334	370	424	18			
1	E	266	Total	C	N	O	S	0	14	0
			2144	1334	372	420	18			
1	F	266	Total	C	N	O	S	0	8	0
			2120	1318	365	419	18			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	PHE	TYR	ENGINEERED MUTATION	UNP BAA13239
B	42	PHE	TYR	ENGINEERED MUTATION	UNP BAA13239
C	42	PHE	TYR	ENGINEERED MUTATION	UNP BAA13239
D	42	PHE	TYR	ENGINEERED MUTATION	UNP BAA13239
E	42	PHE	TYR	ENGINEERED MUTATION	UNP BAA13239
F	42	PHE	TYR	ENGINEERED MUTATION	UNP BAA13239

- Molecule 2 is MANGANESE (III) ION (three-letter code: MN3) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	2	Total	Mn	0	0
			2	2		
2	E	2	Total	Mn	0	0
			2	2		
2	B	2	Total	Mn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	2	Total 2	Mn 2	0	0
2	A	2	Total 2	Mn 2	0	0
2	F	2	Total 2	Mn 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total 1	Ca 1	0	0
3	E	1	Total 1	Ca 1	0	0
3	B	1	Total 1	Ca 1	0	0
3	C	1	Total 1	Ca 1	0	0
3	A	1	Total 1	Ca 1	0	0
3	F	1	Total 1	Ca 1	0	0

- Molecule 4 is OXYGEN ATOM (three-letter code: O) (formula: O).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total 1	O 1	0	0
4	E	1	Total 1	O 1	0	0
4	B	1	Total 1	O 1	0	0
4	C	1	Total 1	O 1	0	0
4	A	1	Total 1	O 1	0	0
4	F	1	Total 1	O 1	0	0

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Na	0	0
			2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	232	Total	O	0	0
			232	232		
7	B	242	Total	O	0	0
			242	242		
7	C	242	Total	O	0	0
			242	242		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	243	Total 243	O 243	0	0
7	E	248	Total 248	O 248	0	0
7	F	252	Total 252	O 252	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: PSEUDOCATALASE

Chain A: 



• Molecule 1: PSEUDOCATALASE

Chain B: 




• Molecule 1: PSEUDOCATALASE

Chain C: 



• Molecule 1: PSEUDOCATALASE

Chain D: 



• Molecule 1: PSEUDOCATALASE

Chain E: 



• Molecule 1: PSEUDOCATALASE

Chain F: 

H1	F2	N11	M39	M40	M67	E89	M120	P121	V138	N144	R147	E148	E163	L192	K234	K238	E246	M247	R260	G266
----	----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.49 Å 95.27 Å 105.00 Å 90.00° 106.55° 90.00°	Depositor
Resolution (Å)	50.00 – 1.33	Depositor
% Data completeness (in resolution range)	89.1 (50.00-1.33)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.115 , 0.145	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14325	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA, MN3, MES, O

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2204	0.81	2/2964 (0.1%)
1	B	0.63	0/2175	0.81	2/2926 (0.1%)
1	C	0.65	0/2236	0.81	2/3007 (0.1%)
1	D	0.65	0/2248	0.83	4/3020 (0.1%)
1	E	0.64	0/2249	0.81	4/3022 (0.1%)
1	F	0.63	0/2196	0.79	1/2953 (0.0%)
All	All	0.64	0/13308	0.81	15/17892 (0.1%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	147	ARG	NE-CZ-NH1	7.17	123.88	120.30
1	C	147	ARG	NE-CZ-NH1	6.71	123.66	120.30
1	D	93[A]	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	D	93[B]	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	E	93	ARG	NE-CZ-NH2	-5.84	117.38	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2123	0	2035	14	0
1	B	2107	0	2016	9	0
1	C	2140	0	2036	19	0
1	D	2146	0	2057	14	0
1	E	2144	0	2053	18	0
1	F	2120	0	2027	15	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	A	12	0	13	0	0
5	E	24	0	25	2	0
5	F	24	0	26	0	0
6	A	2	0	0	0	0
7	A	232	0	0	5	0
7	B	242	0	0	3	0
7	C	242	0	0	8	0
7	D	243	0	0	3	0
7	E	248	0	0	8	0
7	F	252	0	0	8	0
All	All	14325	0	12288	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:260[B]:ARG:NH2	1:E:89:GLU:OE1	1.76	1.17
1:A:11[B]:ASN:ND2	7:A:2009:HOH:O	1.82	1.10
1:C:23[B]:ARG:NH2	7:C:2025:HOH:O	1.92	1.03
1:C:122[A]:ASN:OD1	7:C:2131:HOH:O	1.77	1.02
1:D:260[B]:ARG:CZ	1:E:89:GLU:OE1	2.15	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/266 (103%)	267 (98%)	6 (2%)	0	100	100
1	B	270/266 (102%)	265 (98%)	5 (2%)	0	100	100
1	C	277/266 (104%)	270 (98%)	7 (2%)	0	100	100
1	D	278/266 (104%)	274 (99%)	4 (1%)	0	100	100
1	E	278/266 (104%)	273 (98%)	5 (2%)	0	100	100
1	F	272/266 (102%)	266 (98%)	6 (2%)	0	100	100
All	All	1648/1596 (103%)	1615 (98%)	33 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/222 (104%)	230 (100%)	1 (0%)	93	80
1	B	228/222 (103%)	226 (99%)	2 (1%)	84	57
1	C	235/222 (106%)	234 (100%)	1 (0%)	93	80
1	D	236/222 (106%)	234 (99%)	2 (1%)	86	62
1	E	236/222 (106%)	234 (99%)	2 (1%)	86	62
1	F	230/222 (104%)	229 (100%)	1 (0%)	93	80
All	All	1396/1332 (105%)	1387 (99%)	9 (1%)	88	70

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	147	ARG
1	F	147	ARG
1	E	15	ASP
1	B	216	LEU
1	D	246	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	253	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 31 ligands modelled in this entry, 26 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MES	A	273	-	11,12,12	0.49	0	14,16,16	1.89	4 (28%)
5	MES	E	273	-	11,12,12	0.47	0	14,16,16	1.25	2 (14%)
5	MES	E	274	-	11,12,12	0.50	0	14,16,16	1.99	6 (42%)
5	MES	F	273	-	11,12,12	0.64	0	14,16,16	1.71	4 (28%)
5	MES	F	274	-	11,12,12	0.57	0	14,16,16	1.13	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MES	A	273	-	-	0/6/14/14	0/1/1/1
5	MES	E	273	-	-	0/6/14/14	0/1/1/1
5	MES	E	274	-	-	0/6/14/14	0/1/1/1
5	MES	F	273	-	-	0/6/14/14	0/1/1/1
5	MES	F	274	-	-	0/6/14/14	0/1/1/1

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	274	MES	O2S-S-C8	-3.32	104.07	106.91
5	E	274	MES	O1-C2-C3	-2.15	106.92	111.84
5	A	273	MES	C7-N4-C5	-2.03	106.05	111.27
5	F	273	MES	C7-N4-C5	-2.02	106.09	111.27
5	E	274	MES	C6-O1-C2	2.03	116.73	109.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	273	MES	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.