



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:16 AM GMT

PDB ID : 3O95  
Title : Crystal Structure of Human DPP4 Bound to TAK-100  
Authors : Yano, J.K.; Aertgeerts, K.  
Deposited on : 2010-08-03  
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2228 (2.90-2.82)
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)
RSRZ outliers	91569	2236 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	740	<div> <div>4%</div> <div>77%</div> <div>19%</div> <div>..</div> </div>
1	B	740	<div> <div>%</div> <div>78%</div> <div>19%</div> <div>..</div> </div>
1	C	740	<div> <div>8%</div> <div>78%</div> <div>19%</div> <div>..</div> </div>
1	D	740	<div> <div>4%</div> <div>76%</div> <div>20%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	901	X	-	-	-
3	NAG	B	3211	-	-	-	X
3	NAG	D	5201	X	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 24870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	727	Total	C	N	O	S	0	0	0
			5957	3824	981	1126	26			
1	B	733	Total	C	N	O	S	0	0	0
			6013	3857	997	1133	26			
1	C	726	Total	C	N	O	S	0	0	0
			5946	3818	977	1125	26			
1	D	727	Total	C	N	O	S	0	0	0
			5957	3824	981	1126	26			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	ALA	-	EXPRESSION TAG	UNP P27487
A	28	ASP	-	EXPRESSION TAG	UNP P27487
A	29	PRO	-	EXPRESSION TAG	UNP P27487
A	30	GLY	-	EXPRESSION TAG	UNP P27487
A	31	GLY	-	EXPRESSION TAG	UNP P27487
A	32	SER	-	EXPRESSION TAG	UNP P27487
A	33	HIS	-	EXPRESSION TAG	UNP P27487
A	34	HIS	-	EXPRESSION TAG	UNP P27487
A	35	HIS	-	EXPRESSION TAG	UNP P27487
A	36	HIS	-	EXPRESSION TAG	UNP P27487
A	37	HIS	-	EXPRESSION TAG	UNP P27487
A	38	HIS	-	EXPRESSION TAG	UNP P27487
B	27	ALA	-	EXPRESSION TAG	UNP P27487
B	28	ASP	-	EXPRESSION TAG	UNP P27487
B	29	PRO	-	EXPRESSION TAG	UNP P27487
B	30	GLY	-	EXPRESSION TAG	UNP P27487
B	31	GLY	-	EXPRESSION TAG	UNP P27487
B	32	SER	-	EXPRESSION TAG	UNP P27487
B	33	HIS	-	EXPRESSION TAG	UNP P27487
B	34	HIS	-	EXPRESSION TAG	UNP P27487
B	35	HIS	-	EXPRESSION TAG	UNP P27487

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	36	HIS	-	EXPRESSION TAG	UNP P27487
B	37	HIS	-	EXPRESSION TAG	UNP P27487
B	38	HIS	-	EXPRESSION TAG	UNP P27487
C	27	ALA	-	EXPRESSION TAG	UNP P27487
C	28	ASP	-	EXPRESSION TAG	UNP P27487
C	29	PRO	-	EXPRESSION TAG	UNP P27487
C	30	GLY	-	EXPRESSION TAG	UNP P27487
C	31	GLY	-	EXPRESSION TAG	UNP P27487
C	32	SER	-	EXPRESSION TAG	UNP P27487
C	33	HIS	-	EXPRESSION TAG	UNP P27487
C	34	HIS	-	EXPRESSION TAG	UNP P27487
C	35	HIS	-	EXPRESSION TAG	UNP P27487
C	36	HIS	-	EXPRESSION TAG	UNP P27487
C	37	HIS	-	EXPRESSION TAG	UNP P27487
C	38	HIS	-	EXPRESSION TAG	UNP P27487
D	27	ALA	-	EXPRESSION TAG	UNP P27487
D	28	ASP	-	EXPRESSION TAG	UNP P27487
D	29	PRO	-	EXPRESSION TAG	UNP P27487
D	30	GLY	-	EXPRESSION TAG	UNP P27487
D	31	GLY	-	EXPRESSION TAG	UNP P27487
D	32	SER	-	EXPRESSION TAG	UNP P27487
D	33	HIS	-	EXPRESSION TAG	UNP P27487
D	34	HIS	-	EXPRESSION TAG	UNP P27487
D	35	HIS	-	EXPRESSION TAG	UNP P27487
D	36	HIS	-	EXPRESSION TAG	UNP P27487
D	37	HIS	-	EXPRESSION TAG	UNP P27487
D	38	HIS	-	EXPRESSION TAG	UNP P27487

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	D	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is [5-(AMINOMETHYL)-6-(2,2-DIMETHYLPROPYL)-2-ETHYL-4-(4-METHYLPHENYL)PYRIDIN-3-YL]ACETIC ACID (three-letter code: 01T) (formula:  $C_{22}H_{30}N_2O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			26	22	2	2		
4	B	1	Total	C	N	O	0	0
			26	22	2	2		
4	C	1	Total	C	N	O	0	0
			26	22	2	2		
4	D	1	Total	C	N	O	0	0
			26	22	2	2		

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	3	Total	C	N	O	0	0
			42	24	3	15		

- Molecule 6 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	145	Total 145	O 145	0	0
6	B	134	Total 134	O 134	0	0
6	C	76	Total 76	O 76	0	0
6	D	132	Total 132	O 132	0	0









● Molecule 1: Dipeptidyl peptidase 4



● Molecule 1: Dipeptidyl peptidase 4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.72Å 123.33Å 144.42Å 90.00° 114.78° 90.00°	Depositor
Resolution (Å)	35.00 – 2.85 34.83 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.2 (35.00-2.85) 99.2 (34.83-2.85)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.85Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.187 , 0.248 0.242 , 0.288	Depositor DCC
$R_{free}$ test set	4489 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.2	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 42.8	EDS
Estimated twinning fraction	0.012 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 89524 reflections	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	24870	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 01T, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	1/6129 (0.0%)	0.62	0/8336
1	B	0.52	0/6190	0.61	0/8419
1	C	0.49	0/6118	0.59	0/8322
1	D	0.54	0/6129	0.61	0/8336
All	All	0.52	1/24566 (0.0%)	0.61	0/33413

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	1	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	697	GLN	CD-OE1	5.07	1.35	1.24

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	901	NAG	C1

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5957	0	5673	87	0
1	B	6013	0	5714	81	0
1	C	5946	0	5663	81	0
1	D	5957	0	5674	87	0
2	A	112	0	100	0	0
2	B	56	0	50	2	0
2	D	28	0	25	1	0
3	A	28	0	26	0	0
3	B	42	0	39	0	0
3	C	42	0	39	0	0
3	D	56	0	52	1	0
4	A	26	0	29	2	0
4	B	26	0	29	1	0
4	C	26	0	29	0	0
4	D	26	0	29	1	0
5	C	42	0	37	1	0
6	A	145	0	0	1	0
6	B	134	0	0	3	0
6	C	76	0	0	2	0
6	D	132	0	0	6	0
All	All	24870	0	23208	332	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 332 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:193:ILE:HG22	1:D:194:ILE:HD13	1.49	0.94
1:C:253:ARG:HH21	1:D:253:ARG:HH21	0.94	0.91
3:D:2811:NAG:H2	6:D:845:HOH:O	1.70	0.89
1:C:511:SER:HB3	6:C:812:HOH:O	1.74	0.88
1:B:73:GLU:HA	6:B:866:HOH:O	1.79	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	725/740 (98%)	674 (93%)	47 (6%)	4 (1%)	30	63
1	B	731/740 (99%)	682 (93%)	47 (6%)	2 (0%)	46	76
1	C	724/740 (98%)	667 (92%)	54 (8%)	3 (0%)	39	71
1	D	725/740 (98%)	675 (93%)	44 (6%)	6 (1%)	24	56
All	All	2905/2960 (98%)	2698 (93%)	192 (7%)	15 (0%)	34	67

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	73	GLU
1	B	85	ASN
1	C	73	GLU
1	D	85	ASN
1	D	140	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	652/662 (98%)	607 (93%)	45 (7%)	19	45
1	B	658/662 (99%)	608 (92%)	50 (8%)	16	40
1	C	651/662 (98%)	617 (95%)	34 (5%)	29	60
1	D	652/662 (98%)	598 (92%)	54 (8%)	14	36
All	All	2613/2648 (99%)	2430 (93%)	183 (7%)	19	44

5 of 183 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	514	LEU
1	C	184	ARG
1	D	460	SER
1	B	597	ARG
1	B	728	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	508	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

17 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	1501	1,2	14,14,15	0.63	0	15,19,21	2.14	3 (20%)
2	NAG	A	1502	2	14,14,15	0.53	0	15,19,21	0.85	0
2	NAG	A	1503	1,2	14,14,15	0.64	0	15,19,21	1.56	1 (6%)
2	NAG	A	1504	2	14,14,15	0.68	0	15,19,21	1.48	2 (13%)
2	NAG	A	2291	1,2	14,14,15	0.59	0	15,19,21	0.97	0
2	NAG	A	2292	2	14,14,15	0.60	0	15,19,21	1.43	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	2811	1,2	14,14,15	0.73	0	15,19,21	1.71	3 (20%)
2	NAG	A	2812	2	14,14,15	0.53	0	15,19,21	0.75	0
2	NAG	B	2291	1,2	14,14,15	0.58	0	15,19,21	1.39	1 (6%)
2	NAG	B	2292	2	14,14,15	0.65	0	15,19,21	1.50	2 (13%)
2	NAG	B	901	1,2	14,14,15	0.56	0	15,19,21	1.23	0
2	NAG	B	902	2	14,14,15	0.48	0	15,19,21	0.89	1 (6%)
5	NAG	C	2291	1,5	14,14,15	0.41	0	15,19,21	1.27	1 (6%)
5	NAG	C	2292	5	14,14,15	0.53	0	15,19,21	3.06	6 (40%)
5	NAG	C	2293	5	14,14,15	0.37	0	15,19,21	1.08	1 (6%)
2	NAG	D	2291	1,2	14,14,15	0.55	0	15,19,21	1.45	3 (20%)
2	NAG	D	2292	2	14,14,15	0.49	0	15,19,21	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1501	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1502	2	-	0/6/23/26	0/1/1/1
2	NAG	A	1503	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1504	2	-	0/6/23/26	0/1/1/1
2	NAG	A	2291	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2292	2	-	0/6/23/26	0/1/1/1
2	NAG	A	2811	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2812	2	-	0/6/23/26	0/1/1/1
2	NAG	B	2291	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2292	2	-	0/6/23/26	0/1/1/1
2	NAG	B	901	1,2	1/1/5/7	0/6/23/26	0/1/1/1
2	NAG	B	902	2	-	0/6/23/26	0/1/1/1
5	NAG	C	2291	1,5	-	0/6/23/26	0/1/1/1
5	NAG	C	2292	5	-	0/6/23/26	0/1/1/1
5	NAG	C	2293	5	-	0/6/23/26	0/1/1/1
2	NAG	D	2291	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2292	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	2292	NAG	C4-C3-C2	-5.32	102.97	111.23
2	A	1501	NAG	C4-C3-C2	-3.52	105.75	111.23
5	C	2292	NAG	C3-C4-C5	-2.29	106.20	110.20
5	C	2292	NAG	O7-C7-C8	-2.02	118.35	122.06
2	B	902	NAG	C1-O5-C5	2.02	114.82	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	901	NAG	C1

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2291	NAG	1	0
2	B	2292	NAG	1	0
2	B	901	NAG	1	0
2	B	902	NAG	1	0
5	C	2291	NAG	1	0
2	D	2291	NAG	1	0

## 5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	01T	A	1	-	24,27,27	0.44	0	28,39,39	1.69	5 (17%)
3	NAG	A	2191	1	14,14,15	0.57	0	15,19,21	1.48	2 (13%)
3	NAG	A	3211	1	14,14,15	0.57	0	15,19,21	1.60	1 (6%)
4	01T	B	1	-	24,27,27	0.60	0	28,39,39	1.62	4 (14%)
3	NAG	B	1501	1	14,14,15	0.45	0	15,19,21	1.16	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	B	2811	1	14,14,15	0.54	0	15,19,21	1.78	1 (6%)
3	NAG	B	3211	1	14,14,15	0.66	0	15,19,21	1.57	4 (26%)
4	01T	C	1	-	24,27,27	0.56	0	28,39,39	1.83	6 (21%)
3	NAG	C	1501	1	14,14,15	0.59	0	15,19,21	2.05	4 (26%)
3	NAG	C	2811	1	14,14,15	0.65	0	15,19,21	1.49	2 (13%)
3	NAG	C	3211	1	14,14,15	0.73	0	15,19,21	1.32	3 (20%)
4	01T	D	1	-	24,27,27	0.68	0	28,39,39	1.70	3 (10%)
3	NAG	D	1501	1	14,14,15	0.67	0	15,19,21	2.16	4 (26%)
3	NAG	D	2191	1	14,14,15	0.62	0	15,19,21	1.61	3 (20%)
3	NAG	D	2811	1	14,14,15	0.80	1 (7%)	15,19,21	1.22	2 (13%)
3	NAG	D	5201	1	14,14,15	0.52	0	15,19,21	1.11	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	01T	A	1	-	-	0/14/17/17	0/2/2/2
3	NAG	A	2191	1	-	0/6/23/26	0/1/1/1
3	NAG	A	3211	1	-	0/6/23/26	0/1/1/1
4	01T	B	1	-	-	0/14/17/17	0/2/2/2
3	NAG	B	1501	1	-	0/6/23/26	0/1/1/1
3	NAG	B	2811	1	-	0/6/23/26	0/1/1/1
3	NAG	B	3211	1	-	0/6/23/26	0/1/1/1
4	01T	C	1	-	-	0/14/17/17	0/2/2/2
3	NAG	C	1501	1	-	0/6/23/26	0/1/1/1
3	NAG	C	2811	1	-	0/6/23/26	0/1/1/1
3	NAG	C	3211	1	-	0/6/23/26	0/1/1/1
4	01T	D	1	-	-	0/14/17/17	0/2/2/2
3	NAG	D	1501	1	-	0/6/23/26	0/1/1/1
3	NAG	D	2191	1	-	0/6/23/26	0/1/1/1
3	NAG	D	2811	1	-	0/6/23/26	0/1/1/1
3	NAG	D	5201	1	1/1/5/7	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2811	NAG	C1-C2	2.14	1.55	1.52

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	C	1	01T	C15-C3-N4	-5.85	118.58	123.35
4	D	1	01T	C11-C5-N4	-5.28	119.05	123.35
4	A	1	01T	C11-C5-N4	-4.97	119.30	123.35
4	A	1	01T	C15-C3-N4	-4.78	119.46	123.35
4	D	1	01T	C15-C3-N4	-4.68	119.54	123.35

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	5201	NAG	C1

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	01T	2	0
4	B	1	01T	1	0
4	D	1	01T	1	0
3	D	2811	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	727/740 (98%)	0.29	27 (3%) 45 38	37, 47, 63, 78	0
1	B	733/740 (99%)	0.05	10 (1%) 78 75	35, 47, 61, 80	0
1	C	726/740 (98%)	0.43	60 (8%) 14 9	39, 48, 67, 80	0
1	D	727/740 (98%)	0.31	33 (4%) 37 31	37, 47, 62, 71	0
All	All	2913/2960 (98%)	0.27	130 (4%) 37 31	35, 47, 64, 80	0

The worst 5 of 130 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	138	ASN	5.9
1	D	87	SER	5.2
1	C	333	SER	5.0
1	C	83	TYR	4.7
1	D	91	GLU	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	D	2291	14/15	0.90	0.22	0.65	60,63,67,71	0
2	NAG	A	2291	14/15	0.93	0.21	-0.08	55,57,59,61	0
2	NAG	B	2291	14/15	0.89	0.17	-0.95	58,61,63,67	0
2	NAG	A	1503	14/15	0.79	0.19	-1.33	86,88,90,92	0
5	NAG	C	2291	14/15	0.93	0.14	-1.42	60,63,65,68	0
2	NAG	A	1504	14/15	0.74	0.33	-	94,94,96,96	0
2	NAG	A	2811	14/15	0.74	0.18	-	64,68,70,73	0
2	NAG	B	2292	14/15	0.82	0.31	-	71,72,73,73	0
2	NAG	D	2292	14/15	0.80	0.32	-	74,76,79,80	0
2	NAG	A	2292	14/15	0.74	0.32	-	61,63,65,66	0
5	NAG	C	2292	14/15	0.84	0.21	-	70,73,76,80	0
2	NAG	A	1502	14/15	0.79	0.40	-	86,88,89,89	0
5	NAG	C	2293	14/15	0.76	0.26	-	83,85,87,88	0
2	NAG	A	2812	14/15	0.82	0.26	-	76,78,78,78	0
2	NAG	B	902	14/15	0.75	0.33	-	95,97,97,97	0
2	NAG	A	1501	14/15	0.81	0.32	-	73,78,80,84	0
2	NAG	B	901	14/15	0.69	0.20	-	87,89,91,93	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	D	5201	14/15	0.67	0.35	16.26	75,78,81,82	0
3	NAG	B	3211	14/15	0.80	0.42	7.21	59,62,63,63	0
3	NAG	A	3211	14/15	0.86	0.27	1.62	55,57,62,62	0
4	01T	B	1	26/26	0.95	0.28	0.90	45,48,51,55	0
4	01T	D	1	26/26	0.94	0.25	0.59	45,48,55,58	0
3	NAG	C	3211	14/15	0.71	0.29	0.40	66,69,75,76	0
4	01T	A	1	26/26	0.95	0.29	0.39	46,49,56,59	0
4	01T	C	1	26/26	0.95	0.21	0.33	44,47,50,51	0
3	NAG	D	2811	14/15	0.80	0.19	-	63,65,67,67	0
3	NAG	D	1501	14/15	0.75	0.28	-	63,66,66,66	0
3	NAG	A	2191	14/15	0.78	0.36	-	61,65,70,70	0
3	NAG	C	2811	14/15	0.91	0.15	-	63,65,67,67	0
3	NAG	B	1501	14/15	0.88	0.23	-	61,62,63,64	0
3	NAG	D	2191	14/15	0.87	0.29	-	60,64,67,68	0
3	NAG	C	1501	14/15	0.82	0.19	-	64,66,67,67	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	B	2811	14/15	0.88	0.12	-	57,59,60,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.