



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:15 PM GMT

PDB ID : 4O99  
Title : Crystal structure of Beta-ketothiolase (PhaA) from Ralstonia eutropha H16  
Authors : Kim, E.J.; Kim, J.; Kim, S.; Kim, K.J.  
Deposited on : 2014-01-02  
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	<div> <div>2%</div> <div>92%</div> <div>7%</div> </div>
1	B	392	<div> <div>%</div> <div>91%</div> <div>7%</div> </div>
1	C	392	<div> <div>%</div> <div>90%</div> <div>10%</div> </div>
1	D	392	<div> <div>2%</div> <div>89%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	401	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12325 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetyl-CoA acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			2827	1769	501	537	20			
1	B	392	Total	C	N	O	S	0	0	0
			2827	1769	501	537	20			
1	C	392	Total	C	N	O	S	0	0	0
			2826	1769	501	536	20			
1	D	392	Total	C	N	O	S	0	0	0
			2827	1769	501	537	20			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	254	Total 254	O 254	0	0
3	B	249	Total 249	O 249	0	0
3	C	255	Total 255	O 255	0	0
3	D	248	Total 248	O 248	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acetyl-CoA acetyltransferase




- Molecule 1: Acetyl-CoA acetyltransferase




- Molecule 1: Acetyl-CoA acetyltransferase



- Molecule 1: Acetyl-CoA acetyltransferase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.38Å 105.47Å 106.91Å 90.00° 106.18° 90.00°	Depositor
Resolution (Å)	30.62 – 1.96 30.62 – 1.96	Depositor EDS
% Data completeness (in resolution range)	94.4 (30.62-1.96) 94.4 (30.62-1.96)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	15.71 (at 1.96Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.171 , 0.230 0.183 , 0.234	Depositor DCC
$R_{free}$ test set	4918 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.8	Xtriage
Anisotropy	1.094	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.7	EDS
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 98056 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12325	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.89	1/2867 (0.0%)	0.94	10/3876 (0.3%)
1	B	0.92	2/2867 (0.1%)	0.91	5/3876 (0.1%)
1	C	0.87	0/2866	0.90	3/3876 (0.1%)
1	D	0.88	1/2867 (0.0%)	0.90	5/3876 (0.1%)
All	All	0.89	4/11467 (0.0%)	0.91	23/15504 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	88	CYS	CB-SG	-6.71	1.70	1.82
1	A	88	CYS	CB-SG	-5.97	1.72	1.81
1	B	198	GLU	CB-CG	-5.27	1.42	1.52
1	B	88	CYS	CB-SG	-5.09	1.73	1.81

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	368	ARG	NE-CZ-NH2	8.10	124.35	120.30
1	D	136	ASP	CB-CG-OD1	8.00	125.50	118.30
1	C	375	LEU	CA-CB-CG	7.67	132.93	115.30
1	A	334	ASP	CB-CG-OD1	6.93	124.54	118.30
1	C	357	ARG	NE-CZ-NH1	6.85	123.73	120.30
1	D	136	ASP	CB-CG-OD2	-6.72	112.25	118.30
1	A	129	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	C	357	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	40	ARG	NE-CZ-NH2	-6.19	117.21	120.30
1	B	198	GLU	CB-CA-C	-6.15	98.09	110.40
1	D	368	ARG	NE-CZ-NH1	-5.99	117.31	120.30
1	B	136	ASP	CB-CG-OD2	-5.95	112.95	118.30
1	D	299	ARG	NE-CZ-NH1	5.89	123.24	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ASP	CB-CG-OD2	-5.82	113.06	118.30
1	A	196	ASP	CB-CG-OD1	5.79	123.51	118.30
1	A	136	ASP	CB-CG-OD1	5.68	123.42	118.30
1	D	299	ARG	NE-CZ-NH2	-5.55	117.53	120.30
1	B	368	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	B	221	ARG	NE-CZ-NH1	5.47	123.04	120.30
1	A	334	ASP	CB-CG-OD2	-5.35	113.48	118.30
1	B	378	LEU	CA-CB-CG	5.34	127.58	115.30
1	A	129	ARG	CG-CD-NE	-5.12	101.05	111.80
1	A	88	CYS	CA-CB-SG	-5.03	104.95	114.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2827	0	2895	17	0
1	B	2827	0	2895	22	0
1	C	2826	0	2895	18	0
1	D	2827	0	2895	21	0
2	A	6	0	8	0	0
2	C	6	0	8	0	0
3	A	254	0	0	1	0
3	B	249	0	0	2	0
3	C	255	0	0	1	0
3	D	248	0	0	2	0
All	All	12325	0	11596	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:103:MET:HG2	3:D:572:HOH:O	1.35	1.22
1:D:140:VAL:HG11	1:D:145:VAL:HG21	1.68	0.75
1:B:328:HIS:HD2	1:B:333:TRP:HE1	1.35	0.73
1:D:140:VAL:CG1	1:D:145:VAL:HG21	2.21	0.70
1:C:45:PRO:HB3	1:C:75:LEU:HD23	1.77	0.66
1:B:328:HIS:CD2	1:B:333:TRP:HE1	2.14	0.64
1:B:393:LYS:H	1:B:393:LYS:HE2	1.62	0.64
1:C:149:TRP:HE1	1:C:154:GLN:NE2	1.97	0.63
1:C:88:CYS:SG	1:C:349:HIS:NE2	2.72	0.63
1:B:393:LYS:CE	1:B:393:LYS:H	2.10	0.62
1:B:329:GLN:HE21	1:B:330:GLN:HE21	1.49	0.60
1:D:88:CYS:SG	1:D:349:HIS:CE1	2.95	0.60
1:A:23:LYS:NZ	1:B:130:ASP:OD2	2.36	0.58
1:B:198:GLU:HG2	1:B:360:VAL:HG12	1.86	0.58
1:C:88:CYS:SG	1:C:349:HIS:CE1	2.96	0.58
1:A:251:ASN:ND2	1:A:349:HIS:H	2.03	0.56
1:D:3:ASP:OD1	1:D:262:ALA:HB2	2.05	0.56
1:B:349:HIS:HD2	1:B:354:SER:OG	1.91	0.54
1:B:251:ASN:ND2	1:B:349:HIS:H	2.06	0.53
1:A:133:ARG:NH1	3:A:559:HOH:O	2.33	0.53
1:C:44:LYS:H	1:C:47:GLN:NE2	2.07	0.53
1:C:313:LEU:HD21	1:C:365:GLU:HG3	1.92	0.52
1:A:251:ASN:HD22	1:A:349:HIS:H	1.58	0.52
1:B:299:ARG:HG3	3:B:508:HOH:O	2.10	0.51
1:A:88:CYS:SG	1:A:349:HIS:NE2	2.81	0.50
1:A:393:LYS:CE	1:A:393:LYS:HA	2.42	0.49
1:B:355:GLY:HA2	1:B:378:LEU:HD11	1.94	0.49
1:C:364:HIS:HE1	3:C:565:HOH:O	1.95	0.49
1:A:40:ARG:NH2	1:A:200:VAL:HG12	2.28	0.49
1:B:251:ASN:HD22	1:B:349:HIS:H	1.60	0.48
1:B:73:ALA:O	3:B:538:HOH:O	2.20	0.48
1:D:44:LYS:H	1:D:47:GLN:HE21	1.60	0.48
1:D:88:CYS:SG	1:D:349:HIS:NE2	2.87	0.47
1:B:320:PHE:H	1:B:323:GLN:HE21	1.62	0.47
1:C:88:CYS:HG	1:C:349:HIS:CE1	2.28	0.47
1:A:40:ARG:NH2	1:A:200:VAL:CG1	2.77	0.47
1:A:372:LYS:O	1:A:391:GLU:HA	2.15	0.46
1:C:140:VAL:CG1	1:C:145:VAL:HG21	2.46	0.46
1:A:349:HIS:HD2	1:A:354:SER:OG	1.99	0.46
1:C:140:VAL:HG11	1:C:145:VAL:HG21	1.97	0.45
1:D:378:LEU:N	1:D:378:LEU:HD12	2.32	0.45
1:A:28:GLU:HG2	1:A:72:LYS:HD3	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:88:CYS:HG	1:D:349:HIS:CE1	2.34	0.45
1:A:88:CYS:SG	1:A:349:HIS:CE1	3.10	0.44
1:D:9:ALA:HB1	1:D:360:VAL:HG22	1.98	0.44
1:D:314:MET:HB2	1:D:338:VAL:HG22	1.98	0.44
1:D:72:LYS:HG3	3:D:442:HOH:O	2.19	0.43
1:D:118:MET:N	1:D:252:ASP:OD1	2.50	0.43
1:B:315:GLU:O	1:B:376:ALA:HA	2.18	0.43
1:A:119:SER:O	1:B:129:ARG:HD2	2.18	0.43
1:C:318:GLU:CD	1:C:343:GLY:HA3	2.39	0.43
1:A:328:HIS:HD2	1:A:333:TRP:HE1	1.65	0.42
1:D:26:ALA:HB3	1:D:27:PRO:HD3	2.02	0.42
1:B:140:VAL:CG1	1:B:145:VAL:HG21	2.49	0.42
1:D:229:MET:HG2	1:D:245:ALA:HB1	2.01	0.42
1:C:360:VAL:O	1:C:364:HIS:HD2	2.02	0.42
1:B:351:ILE:HG21	1:B:351:ILE:HD13	1.82	0.42
1:A:40:ARG:HH21	1:A:200:VAL:HG12	1.84	0.42
1:C:71:ILE:HD12	1:D:152:TYR:CZ	2.55	0.42
1:D:210:ASP:OD1	1:D:210:ASP:N	2.53	0.41
1:C:313:LEU:O	1:C:374:GLY:HA2	2.20	0.41
1:A:292:GLY:N	1:A:293:PRO:CD	2.83	0.41
1:B:292:GLY:N	1:B:293:PRO:CD	2.83	0.41
1:D:292:GLY:N	1:D:293:PRO:CD	2.83	0.41
1:C:17:PHE:CZ	1:D:129:ARG:HD3	2.56	0.41
1:B:25:PRO:HD2	1:B:28:GLU:CD	2.41	0.41
1:D:238:LYS:N	1:D:238:LYS:HD3	2.36	0.41
1:C:44:LYS:H	1:C:47:GLN:HE21	1.68	0.41
1:D:88:CYS:O	1:D:378:LEU:HD22	2.21	0.41
1:B:88:CYS:HG	1:B:349:HIS:CE1	2.39	0.41
1:B:140:VAL:HG11	1:B:145:VAL:HG21	2.03	0.41
1:C:200:VAL:HG23	1:C:201:PRO:HD2	2.02	0.41
1:C:315:GLU:O	1:C:376:ALA:HA	2.20	0.41
1:A:200:VAL:HA	1:A:201:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/392 (100%)	378 (97%)	11 (3%)	1 (0%)	46	35
1	B	390/392 (100%)	379 (97%)	10 (3%)	1 (0%)	46	35
1	C	390/392 (100%)	382 (98%)	7 (2%)	1 (0%)	46	35
1	D	390/392 (100%)	381 (98%)	8 (2%)	1 (0%)	46	35
All	All	1560/1568 (100%)	1520 (97%)	36 (2%)	4 (0%)	46	35

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	VAL
1	B	87	VAL
1	C	87	VAL
1	D	87	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/284 (100%)	272 (96%)	12 (4%)	36	21
1	B	284/284 (100%)	277 (98%)	7 (2%)	55	45
1	C	284/284 (100%)	271 (95%)	13 (5%)	33	17
1	D	284/284 (100%)	268 (94%)	16 (6%)	26	11
All	All	1136/1136 (100%)	1088 (96%)	48 (4%)	36	21

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	VAL
1	A	133	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	138	LYS
1	A	155	TYR
1	A	172	ARG
1	A	208	LYS
1	A	238	LYS
1	A	289	MET
1	A	314	MET
1	A	323	GLN
1	A	333	TRP
1	A	393	LYS
1	B	87	VAL
1	B	208	LYS
1	B	289	MET
1	B	333	TRP
1	B	375	LEU
1	B	378	LEU
1	B	393	LYS
1	C	23	LYS
1	C	46	GLU
1	C	125	LEU
1	C	155	TYR
1	C	208	LYS
1	C	226	LEU
1	C	228	SER
1	C	238	LYS
1	C	289	MET
1	C	314	MET
1	C	323	GLN
1	C	333	TRP
1	C	393	LYS
1	D	2	THR
1	D	44	LYS
1	D	46	GLU
1	D	87	VAL
1	D	138	LYS
1	D	155	TYR
1	D	173	GLU
1	D	207	ARG
1	D	210	ASP
1	D	215	LYS
1	D	289	MET
1	D	314	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	323	GLN
1	D	333	TRP
1	D	378	LEU
1	D	393	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	175	GLN
1	A	251	ASN
1	A	328	HIS
1	A	329	GLN
1	B	68	GLN
1	B	154	GLN
1	B	206	GLN
1	B	251	ASN
1	B	323	GLN
1	B	328	HIS
1	B	329	GLN
1	B	349	HIS
1	C	47	GLN
1	C	115	GLN
1	C	154	GLN
1	C	364	HIS
1	D	47	GLN
1	D	115	GLN
1	D	154	GLN
1	D	184	ASN
1	D	364	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	401	-	5,5,5	0.47	0	5,5,5	1.42	1 (20%)
2	GOL	C	401	-	5,5,5	0.53	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	401	-	-	0/4/4/4	0/0/0/0
2	GOL	C	401	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	GOL	O3-C3-C2	-2.67	97.26	110.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	392/392 (100%)	-0.15	7 (1%) 71 80	12, 20, 37, 60	0
1	B	392/392 (100%)	-0.28	5 (1%) 79 86	12, 19, 34, 63	0
1	C	392/392 (100%)	-0.24	5 (1%) 79 86	12, 19, 34, 75	0
1	D	392/392 (100%)	-0.20	6 (1%) 76 84	12, 20, 37, 60	0
All	All	1568/1568 (100%)	-0.22	23 (1%) 76 84	12, 20, 36, 75	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	207	ARG	4.2
1	C	207	ARG	4.2
1	C	208	LYS	3.8
1	D	207	ARG	3.2
1	D	208	LYS	3.2
1	B	209	GLY	2.8
1	A	358	ILE	2.7
1	C	358	ILE	2.6
1	D	227	ASP	2.6
1	A	133	ARG	2.5
1	A	231	GLY	2.4
1	B	208	LYS	2.3
1	B	210	ASP	2.3
1	A	227	ASP	2.2
1	B	212	VAL	2.2
1	D	235	ALA	2.2
1	A	238	LYS	2.2
1	C	133	ARG	2.1
1	A	378	LEU	2.1
1	A	88	CYS	2.1
1	D	359	LEU	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	209	GLY	2.0
1	C	359	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	401	6/6	0.87	0.12	2.60	22,32,34,35	0
2	GOL	C	401	6/6	0.96	0.09	-0.08	24,26,28,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.