



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:17 PM GMT

PDB ID : 4OAU
Title : Complete human RNase L in complex with biological activators.
Authors : Han, Y.; Donovan, J.; Rath, S.; Whitney, G.; Chitrakar, A.; Korennykh, A.
Deposited on : 2014-01-06
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)
RNA backbone	2183	1022 (3.00-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	699	<div> <div>7%</div> <div>77%</div> <div>20%</div> <div>..</div> </div>
2	A	3	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	C	803	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5774 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-5A-dependent ribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	692	Total	C	N	O	S	0	0	0
			5539	3497	975	1045	22			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	672	ASN	HIS	ENGINEERED MUTATION	UNP Q05823

- Molecule 2 is a RNA chain called RNA (5'-R(P*A*AP*A)-2').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	3	Total	C	N	O	P	0	0	0
			67	30	15	19	3			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	2	Total	Mg	0	0
			2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	128	Total	O	0	0
			128	128		
5	A	11	Total	O	0	0
			11	11		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-5A-dependent ribonuclease



- Molecule 2: RNA (5'-R(P*A*AP*A)-2')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	60.13Å 116.60Å 162.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.44 – 2.60 81.20 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (53.44-2.60) 99.5 (81.20-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.211 , 0.251 0.213 , 0.251	Depositor DCC
R_{free} test set	1797 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	0.175	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 47.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 45325 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5774	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.22	0/5643	0.41	0/7600
2	A	9.10	29/73 (39.7%)	7.41	39/105 (37.1%)
All	All	1.05	29/5716 (0.5%)	0.96	39/7705 (0.5%)

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	A	C8-N7	27.96	1.51	1.31
2	A	2	A	N9-C4	26.26	1.53	1.37
2	A	2	A	N3-C4	22.60	1.48	1.34
2	A	2	A	C8-N7	20.43	1.45	1.31
2	A	2	A	N7-C5	20.31	1.51	1.39
2	A	1	A	N9-C4	20.05	1.49	1.37
2	A	2	A	C6-N1	19.41	1.49	1.35
2	A	1	A	C6-N1	15.86	1.46	1.35
2	A	1	A	C2-N3	14.86	1.47	1.33
2	A	3	A	C8-N7	14.57	1.41	1.31
2	A	1	A	C6-N6	13.37	1.44	1.33
2	A	1	A	N3-C4	13.24	1.42	1.34
2	A	3	A	C2-N3	12.91	1.45	1.33
2	A	3	A	C6-N6	12.63	1.44	1.33
2	A	2	A	C6-N6	12.49	1.44	1.33
2	A	3	A	C6-N1	10.55	1.43	1.35
2	A	1	A	C5-C4	9.47	1.45	1.38
2	A	3	A	N3-C4	9.29	1.40	1.34
2	A	2	A	O4'-C1'	8.10	1.52	1.41
2	A	1	A	N1-C2	7.33	1.41	1.34
2	A	1	A	O4'-C1'	6.88	1.50	1.41
2	A	3	A	O4'-C1'	6.56	1.50	1.41
2	A	2	A	N9-C8	5.89	1.42	1.37
2	A	1	A	P-O5'	-5.75	1.53	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2	A	P-OP1	5.62	1.58	1.49
2	A	1	A	N7-C5	5.56	1.42	1.39
2	A	3	A	N1-C2	5.33	1.39	1.34
2	A	1	A	OP3-P	-5.15	1.54	1.61
2	A	1	A	P-OP2	5.00	1.57	1.49

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	A	N9-C1'-C2'	26.40	148.32	114.00
2	A	3	A	O4'-C1'-N9	20.95	124.96	108.20
2	A	1	A	N1-C2-N3	-18.84	119.88	129.30
2	A	2	A	N1-C2-N3	-18.46	120.07	129.30
2	A	3	A	C2-N3-C4	18.44	119.82	110.60
2	A	3	A	N1-C2-N3	-18.42	120.09	129.30
2	A	1	A	C2-N3-C4	18.00	119.60	110.60
2	A	2	A	C2-N3-C4	17.02	119.11	110.60
2	A	2	A	N9-C1'-C2'	16.89	135.95	114.00
2	A	1	A	O4'-C1'-N9	-16.10	95.32	108.20
2	A	2	A	N7-C8-N9	-11.81	107.89	113.80
2	A	3	A	N7-C8-N9	-11.11	108.25	113.80
2	A	1	A	N7-C8-N9	-10.08	108.76	113.80
2	A	2	A	O4'-C1'-N9	-10.07	100.14	108.20
2	A	3	A	N3-C4-C5	-9.52	120.14	126.80
2	A	1	A	N3-C4-C5	-9.14	120.40	126.80
2	A	2	A	C5-N7-C8	8.85	108.32	103.90
2	A	3	A	C5-N7-C8	8.48	108.14	103.90
2	A	2	A	N3-C4-C5	-8.33	120.97	126.80
2	A	1	A	C5'-C4'-O4'	7.63	118.25	109.10
2	A	2	A	C5'-C4'-O4'	7.50	118.10	109.10
2	A	3	A	C5'-C4'-O4'	7.35	117.92	109.10
2	A	1	A	C5-N7-C8	7.15	107.47	103.90
2	A	3	A	N9-C1'-C2'	6.89	122.96	114.00
2	A	1	A	OP1-P-OP2	-6.78	109.43	119.60
2	A	2	A	N9-C4-C5	6.39	108.36	105.80
2	A	3	A	C4-C5-C6	6.25	120.12	117.00
2	A	2	A	C4-C5-N7	-6.22	107.59	110.70
2	A	3	A	C4-C5-N7	-6.03	107.69	110.70
2	A	1	A	C4-C5-C6	5.93	119.97	117.00
2	A	2	A	C4-C5-C6	5.86	119.93	117.00
2	A	3	A	N3-C4-N9	5.60	131.88	127.40
2	A	1	A	N9-C4-C5	5.53	108.01	105.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3	A	OP1-P-OP2	-5.53	111.31	119.60
2	A	2	A	OP1-P-OP2	-5.49	111.37	119.60
2	A	3	A	N9-C4-C5	5.45	107.98	105.80
2	A	3	A	C8-N9-C4	5.37	107.95	105.80
2	A	1	A	N3-C4-N9	5.23	131.58	127.40
2	A	2	A	C8-N9-C4	5.09	107.83	105.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	5539	0	5521	98	0
2	A	67	0	34	0	0
3	C	27	0	12	3	0
4	C	2	0	0	0	0
5	A	11	0	0	0	0
5	C	128	0	0	8	0
All	All	5774	0	5567	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:541:ASP:OD2	5:C:961:HOH:O	1.96	0.81
1:C:601:ASN:OD1	1:C:667:ARG:NH1	2.16	0.79
1:C:327:GLU:OE2	5:C:931:HOH:O	2.01	0.78
1:C:496:LYS:H	1:C:497:LYS:HA	1.53	0.72
1:C:595:THR:HG21	1:C:717:PRO:HD3	1.72	0.72
1:C:139:LYS:NZ	5:C:928:HOH:O	2.26	0.68
1:C:496:LYS:O	5:C:1007:HOH:O	2.13	0.66
1:C:309:ARG:NH2	1:C:327:GLU:OE2	2.22	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:344:GLY:HA2	1:C:347:LEU:HB2	1.80	0.63
1:C:551:VAL:HA	1:C:563:ILE:HD11	1.80	0.62
1:C:45:LEU:HD11	1:C:78:LEU:HD23	1.81	0.62
1:C:496:LYS:N	1:C:497:LYS:HA	2.15	0.61
1:C:232:VAL:HG11	1:C:261:GLN:HG3	1.82	0.60
1:C:473:GLN:OE1	1:C:578:SER:OG	2.17	0.60
1:C:636:ILE:HG12	1:C:637:ASN:H	1.66	0.59
1:C:523:ASP:OD1	1:C:543:LYS:NZ	2.36	0.59
1:C:349:ASP:OD1	1:C:352:ARG:NH2	2.38	0.57
1:C:355:ARG:HD2	1:C:364:PHE:HB2	1.86	0.56
1:C:371:ILE:HG12	1:C:379:ILE:HB	1.87	0.55
1:C:482:THR:HG22	1:C:484:GLN:HG2	1.88	0.55
1:C:594:ARG:NH1	1:C:597:ARG:HH21	2.04	0.54
1:C:343:TRP:CH2	1:C:371:ILE:HB	2.43	0.53
1:C:518:LYS:O	1:C:522:GLU:HG2	2.08	0.53
1:C:303:ASP:O	1:C:307:THR:OG1	2.22	0.52
1:C:263:HIS:O	5:C:959:HOH:O	2.19	0.51
1:C:583:HIS:CD2	1:C:584:PRO:HD2	2.46	0.51
1:C:374:THR:OG1	3:C:801:ADP:O2B	2.30	0.50
1:C:405:VAL:HG21	1:C:430:LEU:HD21	1.94	0.50
1:C:265:GLU:OE1	5:C:1026:HOH:O	2.20	0.49
1:C:669:LEU:HD21	1:C:685:ILE:HD13	1.94	0.49
1:C:442:GLU:HG3	1:C:488:PRO:HB2	1.94	0.49
1:C:27:HIS:HB3	1:C:31:LYS:HE2	1.95	0.48
1:C:558:GLU:HA	1:C:701:ILE:HD11	1.94	0.48
1:C:640:VAL:H	1:C:684:LYS:NZ	2.12	0.48
1:C:518:LYS:HG2	1:C:522:GLU:OE2	2.14	0.48
1:C:526:ARG:HH12	1:C:547:ASN:HB3	1.79	0.48
1:C:257:ARG:NH1	1:C:261:GLN:OE1	2.47	0.47
1:C:609:LYS:HB3	1:C:612:SER:OG	2.13	0.47
1:C:418:VAL:HG22	1:C:435:THR:HG23	1.97	0.47
1:C:355:ARG:HB2	1:C:362:LYS:HE3	1.96	0.47
1:C:639:CYS:HB3	1:C:684:LYS:HE3	1.96	0.47
1:C:275:THR:HG23	1:C:277:LEU:H	1.79	0.47
1:C:161:GLN:HB3	1:C:166:LYS:HB2	1.96	0.47
1:C:596:LEU:HD21	1:C:703:VAL:HG11	1.95	0.47
1:C:285:LEU:HD13	1:C:288:ILE:HD12	1.96	0.47
1:C:441:LEU:O	1:C:445:LEU:HG	2.15	0.46
1:C:636:ILE:HG23	1:C:638:GLU:HB3	1.96	0.46
1:C:597:ARG:HG2	1:C:666:ILE:HG22	1.98	0.46
1:C:54:GLN:HG2	1:C:61:THR:HG22	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:656:GLN:HG2	1:C:661:ASP:OD2	2.15	0.46
1:C:442:GLU:HG2	1:C:530:TYR:OH	2.16	0.45
1:C:584:PRO:HA	1:C:587:TRP:CD2	2.50	0.45
1:C:445:LEU:HA	1:C:448:HIS:HB2	1.98	0.45
1:C:357:MET:SD	1:C:362:LYS:HB2	2.57	0.45
1:C:640:VAL:N	1:C:684:LYS:HZ2	2.15	0.44
1:C:541:ASP:O	1:C:545:GLN:HG2	2.17	0.44
1:C:73:ASP:OD1	1:C:73:ASP:N	2.50	0.44
1:C:559:THR:HG22	1:C:585:PHE:CE1	2.52	0.44
1:C:494:ASP:HB2	1:C:498:ALA:H	1.83	0.44
1:C:591:SER:O	1:C:595:THR:HG23	2.18	0.44
1:C:484:GLN:OE1	1:C:519:ARG:HD2	2.18	0.44
1:C:150:ASN:HB3	1:C:153:LEU:HG	2.00	0.44
1:C:545:GLN:HG2	1:C:545:GLN:H	1.61	0.43
1:C:551:VAL:O	1:C:554:SER:OG	2.36	0.43
1:C:559:THR:HG22	1:C:585:PHE:CD1	2.53	0.43
1:C:114:LYS:NZ	5:C:966:HOH:O	2.51	0.43
1:C:79:LEU:HD23	1:C:83:ALA:HB3	1.99	0.43
1:C:695:THR:HG23	1:C:696:PHE:CD2	2.54	0.43
1:C:637:ASN:HB2	1:C:684:LYS:NZ	2.34	0.43
1:C:371:ILE:HD11	3:C:801:ADP:C4	2.54	0.42
1:C:204:ASN:ND2	1:C:234:VAL:HG11	2.33	0.42
1:C:565:ARG:HH22	1:C:708:GLN:HE22	1.67	0.42
1:C:101:ILE:HG12	1:C:131:GLU:HB3	2.01	0.42
1:C:42:GLN:HG3	1:C:77:LEU:HD21	2.02	0.42
1:C:275:THR:HG23	1:C:277:LEU:N	2.34	0.42
1:C:25:ASP:OD1	1:C:25:ASP:N	2.52	0.42
1:C:337:LYS:NZ	1:C:348:LYS:NZ	2.68	0.42
1:C:615:LEU:HA	1:C:615:LEU:HD12	1.83	0.42
1:C:612:SER:O	1:C:616:ARG:HG2	2.20	0.42
1:C:708:GLN:O	1:C:713:ARG:HD2	2.19	0.42
1:C:604:ASP:HB3	1:C:612:SER:OG	2.20	0.41
1:C:565:ARG:HH22	1:C:708:GLN:NE2	2.18	0.41
1:C:374:THR:HG22	1:C:378:GLY:N	2.35	0.41
1:C:379:ILE:HG13	3:C:801:ADP:O4'	2.19	0.41
1:C:637:ASN:HB2	1:C:684:LYS:HZ1	1.85	0.41
1:C:639:CYS:HB3	1:C:684:LYS:NZ	2.35	0.41
1:C:640:VAL:HG23	1:C:684:LYS:NZ	2.35	0.41
1:C:281:VAL:HG21	1:C:304:LEU:HD22	2.01	0.41
1:C:601:ASN:HA	1:C:667:ARG:HD2	2.03	0.41
1:C:354:TYR:O	1:C:355:ARG:HG2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:583:HIS:CG	1:C:584:PRO:HD2	2.56	0.41
1:C:256:GLN:O	1:C:260:GLU:HG3	2.21	0.41
1:C:159:GLU:O	1:C:163:ARG:HG3	2.21	0.40
1:C:674:ASP:N	1:C:674:ASP:OD1	2.54	0.40
1:C:51:VAL:O	5:C:914:HOH:O	2.22	0.40
1:C:640:VAL:H	1:C:684:LYS:HZ2	1.68	0.40
1:C:458:ASP:OD2	1:C:461:ALA:HB2	2.21	0.40
1:C:147:ARG:HA	1:C:147:ARG:HD3	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	688/699 (98%)	656 (95%)	29 (4%)	3 (0%)	39 65

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	509	LYS
1	C	503	ASP
1	C	555	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	603/607 (99%)	574 (95%)	29 (5%)	31 58

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	25	ASP
1	C	109	LYS
1	C	160	ASP
1	C	184	LEU
1	C	219	GLU
1	C	257	ARG
1	C	291	LEU
1	C	327	GLU
1	C	355	ARG
1	C	363	PHE
1	C	371	ILE
1	C	381	LEU
1	C	415	SER
1	C	419	THR
1	C	434	VAL
1	C	463	ASN
1	C	475	LEU
1	C	495	SER
1	C	501	LEU
1	C	510	TRP
1	C	524	LEU
1	C	591	SER
1	C	607	THR
1	C	638	GLU
1	C	644	MET
1	C	667	ARG
1	C	674	ASP
1	C	710	THR
1	C	719	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	A	0/3	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ADP	C	801	4	22,29,29	1.01	1 (4%)	27,45,45	1.82	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	C	801	4	-	0/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	801	ADP	C5-C4	3.14	1.47	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	C	801	ADP	N3-C2-N1	-6.75	123.72	128.89
3	C	801	ADP	C2'-C1'-N9	-3.23	109.36	114.29
3	C	801	ADP	C4-C5-N7	-3.06	106.67	109.48
3	C	801	ADP	PA-O3A-PB	-2.77	123.37	132.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	801	ADP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	692/699 (98%)	0.27	47 (6%) 20 15	42, 79, 157, 195	0
2	A	3/3 (100%)	0.33	0 100 100	56, 56, 58, 59	0
All	All	695/702 (99%)	0.27	47 (6%) 20 15	42, 79, 157, 195	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	685	ILE	6.6
1	C	675	GLU	5.5
1	C	644	MET	4.7
1	C	719	THR	4.6
1	C	676	GLU	4.3
1	C	678	HIS	3.9
1	C	647	PHE	3.6
1	C	640	VAL	3.6
1	C	528	VAL	3.5
1	C	683	LEU	3.3
1	C	648	TYR	3.3
1	C	677	LYS	3.3
1	C	593	TYR	3.2
1	C	452	ASP	3.1
1	C	625	HIS	3.0
1	C	637	ASN	3.0
1	C	616	ARG	3.0
1	C	680	LYS	3.0
1	C	702	TYR	3.0
1	C	605	ILE	2.9
1	C	672	ASN	2.9
1	C	663	LEU	2.8
1	C	576	CYS	2.8
1	C	668	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	638	GLU	2.6
1	C	686	GLY	2.6
1	C	674	ASP	2.5
1	C	468	ILE	2.5
1	C	566	LEU	2.5
1	C	690	LEU	2.5
1	C	642	LYS	2.5
1	C	532	VAL	2.4
1	C	587	TRP	2.4
1	C	682	LYS	2.3
1	C	695	THR	2.3
1	C	603	SER	2.3
1	C	692	PHE	2.3
1	C	652	GLY	2.2
1	C	694	LYS	2.1
1	C	696	PHE	2.1
1	C	666	ILE	2.1
1	C	714	LYS	2.1
1	C	617	LEU	2.1
1	C	551	VAL	2.0
1	C	597	ARG	2.0
1	C	586	PHE	2.0
1	C	589	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MG	C	803	1/1	0.98	0.22	2.70	70,70,70,70	0
3	ADP	C	801	27/27	0.97	0.17	-0.20	58,68,75,81	0
4	MG	C	802	1/1	0.96	0.12	-	61,61,61,61	0

6.5 Other polymers ⓘ

There are no such residues in this entry.