



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:16 AM GMT

PDB ID : 3ODU
Title : The 2.5 Å structure of the CXCR4 chemokine receptor in complex with small molecule antagonist IT1t
Authors : Wu, B.; Mol, C.D.; Han, G.W.; Katritch, V.; Chien, E.Y.T.; Liu, W.; Cherezov, V.; Stevens, R.C.; Accelerated Technologies Center for Gene to 3D Structure (ATCG3D); GPCR Network (GPCR)
Deposited on : 2010-08-11
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	
1	B	502	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OLC	A	1604	-	-	-	X
4	OLA	A	1610	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-X-C chemokine receptor type 4, Lysozyme Chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	466	Total	C	N	O	S	0	4	0
			3759	2464	630	649	16			
1	B	456	Total	C	N	O	S	0	2	0
			3661	2396	616	633	16			

There are 54 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	ASP	-	expression tag	UNP P61073
A	-8	TYR	-	expression tag	UNP P61073
A	-7	LYS	-	expression tag	UNP P61073
A	-6	ASP	-	expression tag	UNP P61073
A	-5	ASP	-	expression tag	UNP P61073
A	-4	ASP	-	expression tag	UNP P61073
A	-3	ASP	-	expression tag	UNP P61073
A	-2	ALA	-	expression tag	UNP P61073
A	-1	GLY	-	expression tag	UNP P61073
A	0	ALA	-	expression tag	UNP P61073
A	1	PRO	-	expression tag	UNP P61073
A	125	TRP	LEU	engineered	UNP P61073
A	900	GLY	-	linker	UNP P61073
A	901	SER	-	linker	UNP P61073
A	1200	GLY	-	linker	UNP P61073
A	1201	SER	-	linker	UNP P61073
A	1054	THR	CYS	engineered	UNP P00720
A	1097	ALA	CYS	engineered	UNP P00720
A	320	GLY	-	expression tag	UNP P61073
A	321	ARG	-	expression tag	UNP P61073
A	322	PRO	-	expression tag	UNP P61073
A	323	LEU	-	expression tag	UNP P61073
A	324	GLU	-	expression tag	UNP P61073
A	325	VAL	-	expression tag	UNP P61073
A	326	LEU	-	expression tag	UNP P61073

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	327	PHE	-	expression tag	UNP P61073
A	328	GLN	-	expression tag	UNP P61073
B	-9	ASP	-	expression tag	UNP P61073
B	-8	TYR	-	expression tag	UNP P61073
B	-7	LYS	-	expression tag	UNP P61073
B	-6	ASP	-	expression tag	UNP P61073
B	-5	ASP	-	expression tag	UNP P61073
B	-4	ASP	-	expression tag	UNP P61073
B	-3	ASP	-	expression tag	UNP P61073
B	-2	ALA	-	expression tag	UNP P61073
B	-1	GLY	-	expression tag	UNP P61073
B	0	ALA	-	expression tag	UNP P61073
B	1	PRO	-	expression tag	UNP P61073
B	125	TRP	LEU	engineered	UNP P61073
B	900	GLY	-	linker	UNP P61073
B	901	SER	-	linker	UNP P61073
B	1200	GLY	-	linker	UNP P61073
B	1201	SER	-	linker	UNP P61073
B	1054	THR	CYS	engineered	UNP P00720
B	1097	ALA	CYS	engineered	UNP P00720
B	320	GLY	-	expression tag	UNP P61073
B	321	ARG	-	expression tag	UNP P61073
B	322	PRO	-	expression tag	UNP P61073
B	323	LEU	-	expression tag	UNP P61073
B	324	GLU	-	expression tag	UNP P61073
B	325	VAL	-	expression tag	UNP P61073
B	326	LEU	-	expression tag	UNP P61073
B	327	PHE	-	expression tag	UNP P61073
B	328	GLN	-	expression tag	UNP P61073

- Molecule 2 is (6,6-DIMETHYL-5,6-DIHYDROIMIDAZO[2,1-B][1,3]THIAZOL-3-YL)METHYL N,N'-DICYCLOHEXYLIMIDOTHIOCARBAMATE (three-letter code: ITD) (formula: C₂₁H₃₄N₄S₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	S	0	0
			27	21	4	2		
2	B	1	Total	C	N	S	0	0
			27	21	4	2		

- Molecule 3 is (2R)-2,3-DIHYDROXYPROPYL (9Z)-OCTADEC-9-ENOATE (three-letter code: OLC) (formula: C₂₁H₄₀O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			18	14	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			22	18	4		
3	A	1	Total	C	O	0	0
			16	12	4		
3	B	1	Total	C	O	0	0
			15	11	4		
3	B	1	Total	C	O	0	0
			22	18	4		

- Molecule 4 is OLEIC ACID (three-letter code: OLA) (formula: $C_{18}H_{34}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	8	2		
4	B	1	Total	C	O	0	0
			20	18	2		
4	B	1	Total	C	O	0	0
			20	18	2		
4	B	1	Total	C	O	0	0
			20	18	2		
4	B	1	Total	C	O	0	0
			12	10	2		
4	B	1	Total	C	O	0	0
			13	11	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	73	Total 73	O 73	0	0
5	B	69	Total 70	O 70	0	1

GLY
ARG
PRO
LEU
GLU
VAL
LEU
PHE
GLN

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.54Å 83.69Å 120.00Å 90.00° 102.17° 90.00°	Depositor
Resolution (Å)	19.98 – 2.50 19.90 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.98-2.50) 95.7 (19.90-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.50Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.237 , 0.282 0.244 , 0.295	Depositor DCC
R_{free} test set	2079 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 66.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 41455 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7805	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OLA, ITD, OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/3863 (0.0%)	0.71	1/5242 (0.0%)
1	B	0.66	1/3758 (0.0%)	0.77	7/5102 (0.1%)
All	All	0.72	2/7621 (0.0%)	0.74	8/10344 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	206	VAL	CB-CG2	-5.97	1.40	1.52
1	B	184	TYR	CD2-CE2	-5.64	1.30	1.39

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	TYR	CB-CA-C	13.41	137.23	110.40
1	B	63	MET	CB-CA-C	11.22	132.84	110.40
1	A	1200	GLY	N-CA-C	8.84	135.19	113.10
1	B	65	TYR	N-CA-CB	-7.81	96.54	110.60
1	B	100	ALA	N-CA-CB	-6.76	100.63	110.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1200	GLY	Peptide
1	A	1201	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3759	0	3838	95	0
1	B	3661	0	3726	59	0
2	A	27	0	34	3	0
2	B	27	0	34	3	0
3	A	56	0	75	9	0
3	B	37	0	50	1	0
4	A	10	0	12	1	0
4	B	85	0	130	8	0
5	A	73	0	0	2	0
5	B	70	0	0	0	0
All	All	7805	0	7899	150	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 150 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:ASN:ND2	1:A:36:PHE:H	1.62	0.95
1:A:197:VAL:HG21	1:B:194:LEU:HD22	1.50	0.94
1:B:144:SER:O	1:B:147:PRO:HD2	1.68	0.93
1:A:35:ASN:HD22	1:A:36:PHE:H	1.15	0.92
1:A:34:ALA:HA	1:A:38:LYS:HG2	1.59	0.84

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/502 (93%)	451 (96%)	17 (4%)	0	100	100
1	B	456/502 (91%)	440 (96%)	16 (4%)	0	100	100
All	All	924/1004 (92%)	891 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/432 (94%)	378 (93%)	28 (7%)	19	35
1	B	393/432 (91%)	373 (95%)	20 (5%)	29	52
All	All	799/864 (92%)	751 (94%)	48 (6%)	23	43

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1060	LYS
1	A	286	ILE
1	B	1035	LYS
1	A	1125	ARG
1	A	230	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	B	37	ASN
1	B	143	ASN
1	B	200	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ITD	A	1500	-	25,30,30	1.95	5 (20%)	27,42,42	2.20	9 (33%)
3	OLC	A	1600	-	17,17,24	0.53	0	18,18,25	0.73	0
3	OLC	A	1603	-	21,21,24	0.48	0	22,22,25	0.60	0
3	OLC	A	1604	-	15,15,24	0.50	0	16,16,25	0.84	1 (6%)
4	OLA	A	1610	-	6,9,19	0.27	0	6,9,19	0.50	0
2	ITD	B	1500	-	25,30,30	2.00	4 (16%)	27,42,42	2.05	9 (33%)
3	OLC	B	1601	-	14,14,24	0.53	0	15,15,25	0.62	0
3	OLC	B	1602	-	21,21,24	0.44	0	22,22,25	0.57	0
4	OLA	B	1605	-	16,19,19	0.27	0	16,19,19	0.56	0
4	OLA	B	1606	-	16,19,19	0.26	0	16,19,19	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	OLA	B	1607	-	16,19,19	0.26	0	16,19,19	0.55	0
4	OLA	B	1608	-	8,11,19	0.34	0	8,11,19	0.75	0
4	OLA	B	1609	-	9,12,19	0.36	0	9,12,19	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ITD	A	1500	-	-	0/11/39/39	0/3/4/4
3	OLC	A	1600	-	-	0/17/17/24	0/0/0/0
3	OLC	A	1603	-	-	0/21/21/24	0/0/0/0
3	OLC	A	1604	-	-	0/15/15/24	0/0/0/0
4	OLA	A	1610	-	-	0/5/7/17	0/0/0/0
2	ITD	B	1500	-	-	0/11/39/39	0/3/4/4
3	OLC	B	1601	-	-	0/14/14/24	0/0/0/0
3	OLC	B	1602	-	-	0/21/21/24	0/0/0/0
4	OLA	B	1605	-	-	0/15/17/17	0/0/0/0
4	OLA	B	1606	-	-	0/15/17/17	0/0/0/0
4	OLA	B	1607	-	-	0/15/17/17	0/0/0/0
4	OLA	B	1608	-	-	0/7/9/17	0/0/0/0
4	OLA	B	1609	-	-	0/8/10/17	0/0/0/0

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1500	ITD	C9-S2	-7.27	1.67	1.75
2	A	1500	ITD	C9-S2	-7.14	1.67	1.75
2	B	1500	ITD	C2-N1	-3.12	1.46	1.49
2	A	1500	ITD	C2-N1	-2.80	1.46	1.49
2	A	1500	ITD	C4-N1	2.03	1.34	1.31

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1500	ITD	C7-C8-S2	-2.59	107.99	112.84
2	B	1500	ITD	C3-C2-C6	-2.01	109.07	112.13
2	B	1500	ITD	C21-C16-N4	2.09	112.66	109.33
2	B	1500	ITD	C11-C10-N3	2.11	114.47	110.56
3	A	1604	OLC	C21-O20-C1	2.30	123.28	116.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1500	ITD	3	0
3	A	1600	OLC	2	0
3	A	1603	OLC	3	0
3	A	1604	OLC	4	0
4	A	1610	OLA	1	0
2	B	1500	ITD	3	0
3	B	1602	OLC	1	0
4	B	1605	OLA	2	0
4	B	1606	OLA	1	0
4	B	1607	OLA	1	0
4	B	1609	OLA	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	466/502 (92%)	0.54	42 (9%)	12 12	19, 41, 69, 87	0
1	B	456/502 (90%)	0.68	53 (11%)	6 6	21, 41, 80, 112	0
All	All	922/1004 (91%)	0.61	95 (10%)	9 9	19, 41, 74, 112	0

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	317	LEU	8.0
1	B	316	ALA	6.8
1	B	28	CYS	6.7
1	A	35	ASN	6.0
1	B	1037	PRO	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	OLA	A	1610	10/20	0.80	0.31	14.66	23,39,73,117	0
3	OLC	A	1604	16/25	0.81	0.24	2.68	26,43,97,150	0
3	OLC	A	1603	22/25	0.83	0.22	1.99	23,46,79,134	0
4	OLA	B	1609	13/20	0.82	0.27	1.89	22,36,70,105	0
2	ITD	A	1500	27/27	0.94	0.21	1.79	15,40,115,153	0
3	OLC	A	1600	18/25	0.84	0.18	1.55	43,58,112,115	0
4	OLA	B	1605	20/20	0.79	0.23	1.44	32,48,83,150	0
4	OLA	B	1607	20/20	0.85	0.23	1.15	34,54,124,155	0
3	OLC	B	1602	22/25	0.81	0.27	1.14	29,52,104,134	0
3	OLC	B	1601	15/25	0.92	0.20	1.04	17,34,71,103	0
4	OLA	B	1606	20/20	0.79	0.23	0.39	30,54,125,154	0
2	ITD	B	1500	27/27	0.92	0.18	0.17	15,35,59,114	0
4	OLA	B	1608	12/20	0.90	0.14	-0.36	27,49,72,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.