



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:20 PM GMT

PDB ID : 1OI8
Title : 5'-NUCLEOTIDASE (E. COLI) WITH AN ENGINEERED DISULFIDE BRIDGE (P90C, L424C)
Authors : Schultz-Heienbrok, R.; Maier, T.; Straeter, N.
Deposited on : 2003-06-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	532	<div> <div> <div></div> <div>88%</div> <div>10%</div> <div>..</div> </div> </div>
1	B	532	<div> <div> <div>2%</div> <div>86%</div> <div>12%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CO3	A	1554	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN USHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	526	Total	C	N	O	S	0	0	1
			4097	2585	704	789	19			
1	B	526	Total	C	N	O	S	0	0	1
			4097	2585	704	789	19			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	CYS	PRO	ENGINEERED MUTATION	UNP P07024
A	424	CYS	LEU	ENGINEERED MUTATION	UNP P07024
B	90	CYS	PRO	ENGINEERED MUTATION	UNP P07024
B	424	CYS	LEU	ENGINEERED MUTATION	UNP P07024

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mn	0	0
			2	2		
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	1	3		
4	B	1	Total	C	O	0	0
			4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	475	Total 475	O 475	0	0
5	B	441	Total 441	O 441	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN USHA



• Molecule 1: PROTEIN USHA



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	97.69 Å 97.69 Å 312.05 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.96 – 2.10 19.89 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.3 (19.96-2.10) 95.5 (19.89-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.09 Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.156 , 0.205 0.175 , 0.169	Depositor DCC
R_{free} test set	4275 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 84900 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9132	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.67 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.3209e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	3/4184 (0.1%)	1.01	19/5661 (0.3%)
1	B	0.99	6/4184 (0.1%)	0.96	15/5661 (0.3%)
All	All	0.97	9/8368 (0.1%)	0.99	34/11322 (0.3%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	345	MET	SD-CE	-9.83	1.22	1.77
1	B	235	GLU	CG-CD	7.13	1.62	1.51
1	B	362	VAL	CB-CG1	-6.50	1.39	1.52
1	B	438	MET	CG-SD	6.10	1.97	1.81
1	A	362	VAL	CB-CG2	-5.84	1.40	1.52
1	B	235	GLU	CD-OE2	-5.67	1.19	1.25
1	B	235	GLU	CD-OE1	-5.34	1.19	1.25
1	A	410	ARG	CD-NE	-5.28	1.37	1.46
1	A	497	ASN	CB-CG	5.02	1.62	1.51

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ARG	NE-CZ-NH1	13.84	127.22	120.30
1	A	190	ARG	NE-CZ-NH2	-13.46	113.57	120.30
1	A	410	ARG	NE-CZ-NH2	-10.49	115.05	120.30
1	A	410	ARG	NE-CZ-NH1	9.87	125.23	120.30
1	B	235	GLU	OE1-CD-OE2	-9.43	111.99	123.30
1	B	41	ASP	CB-CG-OD2	8.29	125.76	118.30
1	B	194	ASP	CB-CG-OD2	7.50	125.05	118.30
1	A	48	ARG	NE-CZ-NH2	-7.42	116.59	120.30
1	A	162	ASP	CB-CG-OD2	7.05	124.64	118.30
1	B	411	ASP	CB-CG-OD2	7.05	124.65	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	437	ASP	CB-CG-OD2	6.83	124.45	118.30
1	A	90	CYS	CA-CB-SG	-6.79	101.77	114.00
1	A	48	ARG	NE-CZ-NH1	6.71	123.66	120.30
1	B	120	ASP	CB-CG-OD2	6.68	124.31	118.30
1	A	395	ASP	CB-CG-OD2	6.56	124.21	118.30
1	B	395	ASP	CB-CG-OD2	6.55	124.19	118.30
1	B	210	ASP	CB-CG-OD2	6.46	124.11	118.30
1	A	504	ASP	CB-CG-OD2	6.43	124.09	118.30
1	A	268	ASP	CB-CG-OD2	6.23	123.90	118.30
1	A	94	LEU	CA-CB-CG	5.87	128.81	115.30
1	A	456	ASP	CB-CG-OD2	5.87	123.58	118.30
1	B	472	ASP	CB-CG-OD2	5.77	123.50	118.30
1	B	150	ARG	NE-CZ-NH1	5.48	123.04	120.30
1	B	48	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	78	LEU	CA-CB-CG	5.44	127.80	115.30
1	B	513	PRO	C-N-CA	-5.37	111.02	122.30
1	A	374	ASP	CB-CG-OD1	5.28	123.05	118.30
1	A	445	ASP	CB-CG-OD2	5.25	123.03	118.30
1	B	90	CYS	CA-CB-SG	-5.21	104.63	114.00
1	A	100	ASP	CB-CG-OD2	5.16	122.94	118.30
1	B	510	ASP	CB-CG-OD2	5.09	122.88	118.30
1	A	194	ASP	CB-CG-OD2	5.08	122.88	118.30
1	B	255	ASP	CB-CG-OD2	5.08	122.88	118.30
1	B	360	LEU	CB-CG-CD1	5.07	119.62	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4097	0	4009	22	0
1	B	4097	0	4009	32	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	5	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	1	0
5	A	475	0	0	4	0
5	B	441	0	0	10	0
All	All	9132	0	8018	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:HIS:CD2	1:B:252:HIS:HB2	2.14	0.81
1:A:217:HIS:CD2	1:A:252:HIS:HB2	2.15	0.81
1:B:359:GLN:O	1:B:362:VAL:HG12	1.82	0.80
1:A:34:ILE:HD11	1:A:163:LEU:CD1	2.18	0.73
1:A:190:ARG:HD2	1:A:195:GLU:OE1	1.91	0.70
1:B:295:VAL:O	1:B:315:ILE:HG22	1.95	0.67
1:B:34:ILE:HD11	1:B:163:LEU:HD12	1.78	0.65
1:B:69:GLU:HG2	5:B:2049:HOH:O	1.97	0.65
1:A:410:ARG:HD3	3:A:1553:SO4:O4	2.01	0.59
1:B:62:LEU:HD21	1:B:315:ILE:HG23	1.84	0.59
1:A:489:LYS:NZ	5:A:2417:HOH:O	2.36	0.58
1:B:345:MET:HE3	1:B:348:LEU:HD23	1.87	0.56
1:B:201:GLN:NE2	5:B:2162:HOH:O	2.38	0.56
1:B:164:LYS:NZ	1:B:207:GLU:OE1	2.38	0.55
1:B:57:ALA:HA	1:B:345:MET:HG2	1.87	0.55
1:A:257:VAL:HG23	1:A:287:GLN:HB3	1.90	0.54
1:B:311:ASN:ND2	5:B:2249:HOH:O	2.41	0.54
1:A:84:ASP:CG	1:A:217:HIS:CE1	2.81	0.53
1:B:62:LEU:HD21	1:B:315:ILE:CG2	2.39	0.53
1:B:62:LEU:HD11	1:B:315:ILE:HD13	1.90	0.53
4:B:1554:CO3:O3	5:B:2441:HOH:O	2.19	0.52
1:A:90:CYS:HB3	1:A:360:LEU:HD13	1.92	0.52
1:B:355:LYS:NZ	5:B:2295:HOH:O	2.43	0.51
1:B:521:ILE:HD11	5:B:2092:HOH:O	2.11	0.50
1:B:544:LYS:NZ	5:B:2431:HOH:O	2.43	0.50
1:B:362:VAL:HG13	1:B:420:TYR:CE2	2.47	0.50
1:B:535:PRO:O	5:B:2423:HOH:O	2.19	0.49
1:B:345:MET:CE	1:B:348:LEU:HD23	2.42	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273:THR:HB	1:A:274:PRO:HD2	1.95	0.49
1:A:421:LYS:NZ	5:A:2378:HOH:O	2.07	0.49
1:B:34:ILE:HD11	1:B:163:LEU:CD1	2.43	0.49
1:A:467:SER:HB3	1:A:546:GLU:HB3	1.94	0.49
1:B:521:ILE:HD12	5:B:2352:HOH:O	2.15	0.46
1:A:336:THR:HB	1:A:337:PRO:HD2	1.97	0.45
1:B:434:VAL:HG12	1:B:517:ASN:HA	1.99	0.44
1:A:350:SER:N	1:A:351:PRO:CD	2.81	0.44
1:B:302:PHE:HA	1:B:306:GLU:O	2.18	0.43
1:A:34:ILE:HD11	1:A:163:LEU:HD13	2.00	0.43
1:B:84:ASP:CG	1:B:217:HIS:CE1	2.92	0.43
1:B:362:VAL:HG13	1:B:420:TYR:CD2	2.54	0.42
1:B:37:LEU:HA	1:B:296:GLY:O	2.19	0.42
1:B:257:VAL:HG23	1:B:287:GLN:HB3	2.02	0.42
1:A:321:LYS:HD2	5:A:2301:HOH:O	2.18	0.42
1:A:167:VAL:HA	1:A:213:ILE:O	2.19	0.42
1:A:121:ASN:ND2	5:A:2113:HOH:O	2.47	0.42
1:A:350:SER:HB2	1:A:351:PRO:HD3	2.01	0.42
1:B:167:VAL:HA	1:B:213:ILE:O	2.20	0.42
1:A:453:MET:HB2	1:A:461:PRO:HD3	2.01	0.41
1:A:84:ASP:HB2	1:A:217:HIS:HE1	1.86	0.41
1:B:151:LEU:HD11	1:B:187:ILE:HG12	2.03	0.41
1:B:401:PHE:CZ	1:B:493:MET:HB3	2.55	0.41
1:B:447:LEU:HD23	1:B:447:LEU:HA	1.86	0.41
1:A:434:VAL:HG12	1:A:517:ASN:HA	2.02	0.41
1:A:410:ARG:HD2	1:A:429:PHE:CE2	2.56	0.41
1:B:243:GLY:HA2	5:B:2232:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	524/532 (98%)	509 (97%)	13 (2%)	2 (0%)	39	37
1	B	524/532 (98%)	509 (97%)	13 (2%)	2 (0%)	39	37
All	All	1048/1064 (98%)	1018 (97%)	26 (2%)	4 (0%)	39	37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	HIS
1	A	87	THR
1	B	87	THR
1	B	289	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	433/440 (98%)	416 (96%)	17 (4%)	39	39
1	B	433/440 (98%)	423 (98%)	10 (2%)	58	62
All	All	866/880 (98%)	839 (97%)	27 (3%)	47	50

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	LEU
1	A	90	CYS
1	A	94	LEU
1	A	131	LYS
1	A	144	GLN
1	A	178	ILE
1	A	190	ARG
1	A	264	LYS
1	A	323	VAL
1	A	330	SER
1	A	355	LYS
1	A	410	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	471	LYS
1	A	472	ASP
1	A	481	LYS
1	A	527	LYS
1	A	539	SER
1	B	78	LEU
1	B	159	LYS
1	B	162	ASP
1	B	178	ILE
1	B	188	GLU
1	B	360	LEU
1	B	441	LYS
1	B	471	LYS
1	B	496	LEU
1	B	550	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	GLN
1	A	343	GLN
1	A	431	ASN
1	A	511	ASN
1	B	201	GLN
1	B	226	HIS
1	B	266	GLN
1	B	476	ASN
1	B	499	ASN
1	B	550	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1553	-	4,4,4	0.87	0	6,6,6	1.02	0
4	CO3	A	1554	2	0,3,3	0.00	-	0,3,3	0.00	-
3	SO4	B	1553	-	4,4,4	0.72	0	6,6,6	1.01	1 (16%)
4	CO3	B	1554	2	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	1553	-	-	0/0/0/0	0/0/0/0
4	CO3	A	1554	2	-	0/0/0/0	0/0/0/0
3	SO4	B	1553	-	-	0/0/0/0	0/0/0/0
4	CO3	B	1554	2	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1553	SO4	O2-S-O1	-2.14	102.71	109.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1553	SO4	1	0
4	B	1554	CO3	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	526/532 (98%)	-0.45	6 (1%) 82 86	15, 23, 37, 53	0
1	B	526/532 (98%)	-0.38	10 (1%) 70 75	14, 24, 38, 54	0
All	All	1052/1064 (98%)	-0.42	16 (1%) 76 81	14, 23, 38, 54	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	551	GLU	6.1
1	B	472	ASP	5.0
1	A	550	GLN	3.8
1	B	162	ASP	3.7
1	A	267	VAL	3.6
1	A	472	ASP	3.3
1	A	471	LYS	3.3
1	B	473	GLY	3.0
1	B	72	ALA	3.0
1	B	71	ALA	2.4
1	B	29	ASP	2.3
1	B	550	GLN	2.3
1	A	268	ASP	2.2
1	B	68	LYS	2.2
1	B	267	VAL	2.1
1	B	27	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	CO3	A	1554	4/4	0.84	0.13	2.35	26,33,34,35	0
4	CO3	B	1554	4/4	0.86	0.12	1.69	35,36,37,38	0
2	MN	B	1552	1/1	1.00	0.07	-0.50	21,21,21,21	0
2	MN	A	1551	1/1	1.00	0.06	-0.77	21,21,21,21	0
2	MN	B	1551	1/1	1.00	0.01	-2.87	32,32,32,32	0
2	MN	A	1552	1/1	1.00	0.02	-3.27	29,29,29,29	0
3	SO4	B	1553	5/5	0.89	0.24	-	51,58,64,68	0
3	SO4	A	1553	5/5	0.87	0.20	-	44,47,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.