wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 07:20 PM GMT

PDB ID	:	4OKO
Title	:	Crystal structure of Francisella tularensis REP34 (Rapid Encystment Phenotype Protein 34 KDa)
Authors	:	Feld, G.K.; Segelke, B.W.; Corzett, M.H.; Hunter, M.S.; Frank, M.; Rasley, A.
Deposited on	:	2014-01-22
Resolution	:	2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1799 (2.08-2.04)
Clashscore	102246	1910 (2.08-2.04)
Ramachandran outliers	100387	1893 (2.08-2.04)
Sidechain outliers	100360	1893 (2.08-2.04)
RSRZ outliers	91569	1802 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	11% 89% • 7%
1	B	312	11% 85% 7% 8%
1	C	312	11% 85% 6% 10%
1	D	312	10% 86% 6% 7%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 18267 atoms, of which 8817 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rapid Encystment Phenotype Protein 34 KDa.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	290	4556	1482	2234	386	444	10	0	1	0
1	B	287	4504	1471	2201	380	442	10	0	3	0
1	C	282	4410	1436	2160	372	432	10	0	1	0
1	D	289	4514	1468	2210	383	443	10	0	1	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	305	LEU	-	EXPRESSION TAG	UNP A0Q494
A	306	GLU	-	EXPRESSION TAG	UNP A0Q494
A	307	HIS	-	EXPRESSION TAG	UNP A0Q494
A	308	HIS	-	EXPRESSION TAG	UNP A0Q494
A	309	HIS	-	EXPRESSION TAG	UNP A0Q494
A	310	HIS	-	EXPRESSION TAG	UNP A0Q494
A	311	HIS	-	EXPRESSION TAG	UNP A0Q494
A	312	HIS	-	EXPRESSION TAG	UNP A0Q494
B	305	LEU	-	EXPRESSION TAG	UNP A0Q494
B	306	GLU	-	EXPRESSION TAG	UNP A0Q494
B	307	HIS	-	EXPRESSION TAG	UNP A0Q494
B	308	HIS	-	EXPRESSION TAG	UNP A0Q494
B	309	HIS	-	EXPRESSION TAG	UNP A0Q494
B	310	HIS	-	EXPRESSION TAG	UNP A0Q494
B	311	HIS	-	EXPRESSION TAG	UNP A0Q494
B	312	HIS	-	EXPRESSION TAG	UNP A0Q494
C	305	LEU	-	EXPRESSION TAG	UNP A0Q494
C	306	GLU	-	EXPRESSION TAG	UNP A0Q494
C	307	HIS	-	EXPRESSION TAG	UNP A0Q494
C	308	HIS	-	EXPRESSION TAG	UNP A0Q494
C	309	HIS	-	EXPRESSION TAG	UNP A0Q494

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	310	HIS	-	EXPRESSION TAG	UNP A0Q494
C	311	HIS	-	EXPRESSION TAG	UNP A0Q494
C	312	HIS	-	EXPRESSION TAG	UNP A0Q494
D	305	LEU	-	EXPRESSION TAG	UNP A0Q494
D	306	GLU	-	EXPRESSION TAG	UNP A0Q494
D	307	HIS	-	EXPRESSION TAG	UNP A0Q494
D	308	HIS	-	EXPRESSION TAG	UNP A0Q494
D	309	HIS	-	EXPRESSION TAG	UNP A0Q494
D	310	HIS	-	EXPRESSION TAG	UNP A0Q494
D	311	HIS	-	EXPRESSION TAG	UNP A0Q494
D	312	HIS	-	EXPRESSION TAG	UNP A0Q494

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	7	2	3	2	0	0
3	B	1	7	2	3	2	0	0
3	C	1	7	2	3	2	0	0
3	D	1	7	2	3	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	69	70	70	0	1
4	B	51	52	52	0	1
4	C	54	56	56	0	2
4	D	72	73	73	0	1

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rapid Encystment Phenotype Protein 34 KDa

Chain A:



- Molecule 1: Rapid Encystment Phenotype Protein 34 KDa

Chain B:



- Molecule 1: Rapid Encystment Phenotype Protein 34 KDa

Chain C:



- Molecule 1: Rapid Encystment Phenotype Protein 34 KDa

Chain D:




PRO
SER
ILE
ASN
ILE
LEU
GLY
ASP
ASP
I243
I244
R245
M249
E256
R257
F262
Y274
N281
P282
Q302
K303
N304
LEU
GLU
HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	62.71Å 63.31Å 81.69Å 70.36° 84.34° 82.53°	Depositor
Resolution (Å)	21.54 – 2.05 21.54 – 2.05	Depositor EDS
% Data completeness (in resolution range)	94.2 (21.54-2.05) 80.7 (21.54-2.05)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ [1]	2.61 (at 2.06Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.213 , 0.233 0.214 , 0.233	Depositor DCC
R_{free} test set	1899 reflections (2.88%)	DCC
Wilson B-factor (Å^2)	34.3	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.44 , 49.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 68959 reflections	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	18267	wwPDB-VP
Average B, all atoms (Å^2)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.15% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.25	0/2380	0.44	0/3233
1	B	0.26	0/2368	0.46	0/3215
1	C	0.25	0/2308	0.44	0/3134
1	D	0.30	1/2362 (0.0%)	0.46	0/3207
All	All	0.26	1/9418 (0.0%)	0.45	0/12789

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	218	TYR	CE1-CZ	5.89	1.46	1.38

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2322	2234	2246	5	0
1	B	2303	2201	2229	18	2
1	C	2250	2160	2173	11	0
1	D	2304	2210	2229	7	2
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	4	3	3	0	0
3	B	4	3	3	0	0
3	C	4	3	3	0	0
3	D	4	3	3	0	0
4	A	70	0	0	0	0
4	B	52	0	0	1	0
4	C	56	0	0	0	0
4	D	73	0	0	0	0
All	All	9450	8817	8889	37	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 37 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:GLU:OE2	1:C:287:LEU:HD12	1.77	0.84
1:B:182:ARG:NH1	1:B:196:TRP:HA	1.94	0.83
1:A:231:PRO:O	1:A:232:THR:HB	1.87	0.74
1:B:242:ASP:N	1:B:242:ASP:OD1	2.22	0.73
1:C:281:ASN:HB2	1:C:282:PRO:HD2	1.76	0.67

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:GLU:OE2	1:D:27:ASN:ND2[1_455]	2.10	0.10
1:B:18:GLU:OE2	1:D:27:ASN:HD22[1_455]	1.56	0.04

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/312 (92%)	281 (98%)	6 (2%)	0	100	100
1	B	284/312 (91%)	278 (98%)	6 (2%)	0	100	100
1	C	277/312 (89%)	272 (98%)	5 (2%)	0	100	100
1	D	284/312 (91%)	277 (98%)	7 (2%)	0	100	100
All	All	1132/1248 (91%)	1108 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/267 (93%)	244 (98%)	4 (2%)	70	67
1	B	247/267 (92%)	245 (99%)	2 (1%)	86	86
1	C	241/267 (90%)	239 (99%)	2 (1%)	86	86
1	D	247/267 (92%)	241 (98%)	6 (2%)	57	51
All	All	983/1068 (92%)	969 (99%)	14 (1%)	76	72

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	243	ILE
1	C	262	PHE
1	D	232	THR
1	B	262	PHE
1	D	194	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	302	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	6	HIS
1	B	33	GLN
1	D	4	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	ACT	A	402	-	1,3,3	1.14	0	0,3,3	0.00	-
3	ACT	B	402	-	1,3,3	1.13	0	0,3,3	0.00	-
3	ACT	C	402	-	1,3,3	1.09	0	0,3,3	0.00	-
3	ACT	D	402	-	1,3,3	1.06	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	402	-	-	0/0/0/0	0/0/0/0
3	ACT	B	402	-	-	0/0/0/0	0/0/0/0
3	ACT	C	402	-	-	0/0/0/0	0/0/0/0
3	ACT	D	402	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	290/312 (92%)	0.71	34 (11%)	6	6	25, 37, 59, 83	1 (0%)
1	B	287/312 (91%)	0.81	33 (11%)	6	7	26, 41, 69, 92	1 (0%)
1	C	282/312 (90%)	0.72	34 (12%)	6	6	26, 40, 64, 88	0
1	D	289/312 (92%)	0.69	30 (10%)	8	9	26, 39, 61, 93	2 (0%)
All	All	1148/1248 (91%)	0.73	131 (11%)	7	7	25, 39, 64, 93	4 (0%)

The worst 5 of 131 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	191	ILE	5.8
1	C	191	ILE	5.3
1	B	196	TRP	4.2
1	D	198	ILE	4.1
1	D	197	GLY	4.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
3	ACT	C	402	4/4	0.88	0.22	1.82	45,47,57,57	0
3	ACT	D	402	4/4	0.89	0.20	1.12	41,46,55,55	0
3	ACT	A	402	4/4	0.95	0.17	0.19	41,47,56,56	0
3	ACT	B	402	4/4	0.91	0.14	-0.80	45,45,55,55	0
2	ZN	B	401	1/1	0.99	0.13	-1.01	61,61,61,61	0
2	ZN	D	401	1/1	0.99	0.07	-1.76	48,48,48,48	0
2	ZN	C	401	1/1	0.99	0.07	-2.10	47,47,47,47	0
2	ZN	A	401	1/1	0.98	0.06	-3.14	50,50,50,50	0

6.5 Other polymers (i)

There are no such residues in this entry.