



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:16 AM GMT

PDB ID : 3OL2
Title : Receptor-ligand structure of Human Semaphorin 4D with Plexin B1.
Authors : Janssen, B.J.C.; Robinson, R.A.; Perez-Branguli, F.; Bell, C.H.; Mitchell, C.J.; Siebold, C.; Jones, E.Y.
Deposited on : 2010-08-25
Resolution : 2.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	663	 3% 80% 12% 7%
2	B	528	 2% 77% 15% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	A	9000	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Semaphorin-4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	614	Total	C	N	O	S	0	0	0
			4866	3100	840	903	23			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	678	LYS	-	EXPRESSION TAG	UNP Q92854
A	679	HIS	-	EXPRESSION TAG	UNP Q92854
A	680	HIS	-	EXPRESSION TAG	UNP Q92854
A	681	HIS	-	EXPRESSION TAG	UNP Q92854
A	682	HIS	-	EXPRESSION TAG	UNP Q92854
A	683	HIS	-	EXPRESSION TAG	UNP Q92854
A	684	HIS	-	EXPRESSION TAG	UNP Q92854

- Molecule 2 is a protein called Plexin-B1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	496	Total	C	N	O	S	0	0	0
			3747	2354	652	719	22			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	17	GLU	-	CLONING ARTIFACT	UNP O43157
B	18	THR	-	CLONING ARTIFACT	UNP O43157
B	19	GLY	-	CLONING ARTIFACT	UNP O43157
B	536	GLY	-	EXPRESSION TAG	UNP O43157
B	537	THR	-	EXPRESSION TAG	UNP O43157
B	538	LYS	-	EXPRESSION TAG	UNP O43157
B	539	HIS	-	EXPRESSION TAG	UNP O43157
B	540	HIS	-	EXPRESSION TAG	UNP O43157
B	541	HIS	-	EXPRESSION TAG	UNP O43157

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	542	HIS	-	EXPRESSION TAG	UNP O43157
B	543	HIS	-	EXPRESSION TAG	UNP O43157
B	544	HIS	-	EXPRESSION TAG	UNP O43157

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			39	22	2	15		
3	A	3	Total	C	N	O	0	0
			39	22	2	15		
3	B	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	0
			28	16	2	10		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	0
			28	16	2	10		
5	A	2	Total	C	N	O	0	0
			28	16	2	10		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Semaphorin-4D



• Molecule 2: Plexin-B1



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	83.06Å 173.44Å 482.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.49 – 2.99 49.49 – 2.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.49-2.99) 99.3 (49.49-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, R_{free}	0.190 , 0.237 0.194 , 0.241	Depositor DCC
R_{free} test set	1760 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	71.6	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 35192 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8828	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/4984	0.76	0/6757
2	B	0.55	0/3842	0.81	2/5247 (0.0%)
All	All	0.51	0/8826	0.78	2/12004 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	350	GLY	C-N-CA	6.25	137.34	121.70
2	B	351	THR	N-CA-CB	5.10	119.99	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4866	0	4783	38	0
2	B	3747	0	3586	46	0
3	A	78	0	68	1	0
3	B	39	0	34	1	0
4	A	14	0	13	2	0
5	A	84	0	75	1	0
All	All	8828	0	8559	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:350:GLY:HA2	2:B:351:THR:HG23	1.40	1.00
2:B:350:GLY:HA2	2:B:351:THR:CG2	2.17	0.72
2:B:232:PHE:HE2	2:B:249:SER:HB2	1.60	0.66
1:A:278:ARG:NH2	1:A:401:ASP:O	2.28	0.66
2:B:279:ALA:HB1	2:B:293:LEU:HD11	1.77	0.66
2:B:531:LEU:HB3	2:B:532:GLY:HA2	1.77	0.65
2:B:148:PRO:O	2:B:150:VAL:N	2.30	0.65
1:A:320:VAL:N	1:A:321:GLY:HA2	2.13	0.63
2:B:398:LEU:HA	2:B:439:GLY:HA3	1.81	0.62
1:A:440:PHE:HB3	1:A:448:LEU:HD11	1.83	0.59
1:A:255:ARG:HH11	1:A:255:ARG:HG3	1.67	0.59
2:B:481:SER:O	2:B:484:GLN:HG2	2.02	0.58
1:A:93:LYS:HE3	1:A:132:GLN:HE22	1.69	0.57
2:B:148:PRO:C	2:B:150:VAL:H	2.07	0.57
2:B:432:TYR:HB2	2:B:497:PRO:HG3	1.88	0.56
1:A:172:GLU:HG2	1:A:192:SER:HA	1.86	0.56
1:A:604:MET:HB2	1:A:610:LEU:HD13	1.86	0.56
1:A:193:SER:HB2	5:A:9000:NAG:H4	1.88	0.56
1:A:80:GLU:HG2	4:A:3000:NAG:H82	1.87	0.56
2:B:356:ILE:HD11	2:B:383:PRO:HB2	1.88	0.55
2:B:147:ASP:C	2:B:148:PRO:O	2.42	0.55
2:B:55:LEU:HD11	2:B:110:VAL:HG11	1.89	0.54
2:B:330:ASP:HB2	3:B:2000:NAG:H82	1.89	0.53
1:A:524:PRO:HD2	1:A:525:PRO:HD2	1.91	0.53
2:B:301:ALA:HB2	2:B:312:ALA:HA	1.91	0.52
2:B:201:GLU:C	2:B:203:THR:H	2.14	0.51
2:B:531:LEU:HB3	2:B:532:GLY:CA	2.42	0.50
1:A:539:SER:HA	1:A:542:LEU:HD12	1.92	0.50
2:B:346:ARG:HA	2:B:352:GLU:H	1.76	0.50
1:A:255:ARG:NH1	1:A:255:ARG:HG3	2.27	0.49
1:A:278:ARG:HD2	1:A:280:ILE:HD11	1.94	0.49
1:A:206:LEU:HB3	1:A:209:PRO:HG3	1.94	0.48
2:B:226:ALA:HB1	2:B:254:ARG:HH12	1.78	0.48
1:A:111:TYR:CE1	3:A:1000:NAG:H82	2.49	0.48
1:A:594:LEU:HD11	1:A:601:TYR:CZ	2.48	0.48
2:B:49:LEU:HD13	2:B:58:LEU:HD13	1.95	0.48
2:B:224:ARG:HD3	2:B:326:LEU:HD13	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:102:LEU:HD11	2:B:154:GLY:HA3	1.96	0.47
2:B:35:LEU:HD22	2:B:49:LEU:HD23	1.96	0.47
2:B:487:ASP:OD1	2:B:490:SER:HB2	2.15	0.47
2:B:183:PRO:HD3	2:B:211:LEU:HD11	1.97	0.47
2:B:409:ALA:HB2	2:B:457:LEU:HB3	1.94	0.47
2:B:346:ARG:O	2:B:351:THR:HA	2.14	0.47
1:A:320:VAL:H	1:A:321:GLY:HA2	1.80	0.47
1:A:589:PHE:HB3	1:A:594:LEU:HD13	1.97	0.47
2:B:331:ARG:HG2	2:B:335:ARG:HH12	1.80	0.46
1:A:32:TRP:CZ3	1:A:579:LYS:HD2	2.51	0.46
1:A:626:SER:O	1:A:636:PHE:HA	2.16	0.45
2:B:356:ILE:HG13	2:B:362:SER:HB3	1.99	0.45
2:B:492:LEU:HD13	2:B:529:PRO:HB3	1.98	0.45
2:B:331:ARG:O	2:B:335:ARG:HG2	2.17	0.45
2:B:168:VAL:O	2:B:184:ILE:HA	2.16	0.45
1:A:114:VAL:O	1:A:125:VAL:HA	2.16	0.45
1:A:444:ASP:O	1:A:472:PRO:HG3	2.17	0.45
1:A:587:TRP:CE2	1:A:609:LEU:HB2	2.52	0.45
2:B:230:PHE:HB2	2:B:249:SER:HB3	1.99	0.44
2:B:321:LEU:HD23	2:B:397:ILE:HD11	1.99	0.44
1:A:445:ARG:O	1:A:470:PHE:HA	2.17	0.44
2:B:249:SER:OG	2:B:261:TYR:CZ	2.69	0.44
1:A:614:LEU:HB3	1:A:646:VAL:HG21	1.99	0.44
2:B:245:ARG:HB3	2:B:247:TYR:CE2	2.52	0.44
1:A:251:PRO:HB2	1:A:279:LEU:HB3	1.99	0.44
1:A:279:LEU:HD12	1:A:279:LEU:HA	1.85	0.44
1:A:561:TYR:CD1	1:A:641:LYS:HB3	2.52	0.44
2:B:531:LEU:CB	2:B:532:GLY:HA2	2.43	0.44
1:A:192:SER:O	1:A:193:SER:HB3	2.18	0.44
2:B:350:GLY:HA2	2:B:351:THR:CB	2.48	0.43
2:B:343:ARG:HE	2:B:347:ALA:HA	1.83	0.43
2:B:77:ARG:NH1	2:B:146:ASN:O	2.51	0.43
1:A:442:SER:HB2	1:A:473:VAL:HB	2.00	0.43
2:B:262:VAL:HG23	2:B:386:MET:HB2	2.01	0.42
1:A:80:GLU:HA	4:A:3000:NAG:H82	2.00	0.42
2:B:279:ALA:HA	2:B:294:PHE:O	2.19	0.42
2:B:97:PRO:HD2	2:B:113:SER:HB2	2.02	0.42
1:A:113:ARG:HB2	1:A:156:CYS:SG	2.59	0.41
1:A:133:PRO:HB3	1:A:156:CYS:HB3	2.01	0.41
1:A:250:ILE:HG12	1:A:278:ARG:HG2	2.02	0.41
1:A:209:PRO:HA	1:A:236:THR:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:525:TRP:CE2	2:B:527:PHE:HB2	2.56	0.41
2:B:229:TYR:HA	2:B:249:SER:O	2.21	0.41
2:B:185:THR:HG22	2:B:205:LYS:HB3	2.02	0.41
1:A:110:ASN:OD1	1:A:127:GLY:HA3	2.20	0.40
1:A:74:ASN:HB3	1:A:80:GLU:HB2	2.03	0.40
1:A:629:ARG:N	1:A:629:ARG:HD2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	608/663 (92%)	565 (93%)	42 (7%)	1 (0%)	52	88
2	B	486/528 (92%)	449 (92%)	30 (6%)	7 (1%)	14	51
All	All	1094/1191 (92%)	1014 (93%)	72 (7%)	8 (1%)	26	70

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	351	THR
2	B	149	ALA
2	B	529	PRO
2	B	148	PRO
2	B	347	ALA
1	A	616	GLU
2	B	456	ASP
2	B	439	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	539/582 (93%)	508 (94%)	31 (6%)	25	63
2	B	399/425 (94%)	382 (96%)	17 (4%)	35	75
All	All	938/1007 (93%)	890 (95%)	48 (5%)	29	69

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ARG
1	A	36	GLU
1	A	37	VAL
1	A	55	LEU
1	A	59	LYS
1	A	90	SER
1	A	94	LYS
1	A	118	LEU
1	A	143	PHE
1	A	165	THR
1	A	175	SER
1	A	191	ASN
1	A	195	SER
1	A	210	SER
1	A	255	ARG
1	A	301	SER
1	A	313	PHE
1	A	320	VAL
1	A	350	GLN
1	A	366	ARG
1	A	417	ASP
1	A	418	VAL
1	A	422	GLN
1	A	431	LEU
1	A	435	VAL
1	A	453	SER
1	A	478	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	487	VAL
1	A	539	SER
1	A	549	ASP
1	A	644	LEU
2	B	92	GLN
2	B	104	SER
2	B	114	VAL
2	B	151	SER
2	B	220	SER
2	B	256	GLN
2	B	344	GLU
2	B	348	GLU
2	B	362	SER
2	B	366	GLN
2	B	370	ASP
2	B	433	LEU
2	B	440	HIS
2	B	449	GLN
2	B	472	SER
2	B	504	LEU
2	B	512	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

15 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1000	1,3	14,14,15	0.55	0	15,19,21	1.12	1 (6%)
3	NAG	A	1001	3	14,14,15	0.73	0	15,19,21	1.22	1 (6%)
3	BMA	A	1002	3	11,11,12	0.74	0	14,15,17	0.70	0
5	NAG	A	4000	1,5	14,14,15	0.53	0	15,19,21	1.32	3 (20%)
5	NAG	A	4001	5	14,14,15	0.50	0	15,19,21	1.22	1 (6%)
3	NAG	A	6000	1,3	14,14,15	0.42	0	15,19,21	1.31	3 (20%)
3	NAG	A	6001	3	14,14,15	0.69	0	15,19,21	1.06	1 (6%)
3	BMA	A	6002	3	11,11,12	0.57	0	14,15,17	0.82	0
5	NAG	A	8000	1,5	14,14,15	0.45	0	15,19,21	1.20	2 (13%)
5	NAG	A	8001	5	14,14,15	0.56	0	15,19,21	1.06	1 (6%)
5	NAG	A	9000	1,5	14,14,15	0.96	1 (7%)	15,19,21	1.03	0
5	NAG	A	9001	5	14,14,15	0.62	0	15,19,21	1.14	1 (6%)
3	NAG	B	2000	3,2	14,14,15	0.42	0	15,19,21	1.25	2 (13%)
3	NAG	B	2001	3	14,14,15	0.59	0	15,19,21	1.18	2 (13%)
3	BMA	B	2002	3	11,11,12	0.60	0	14,15,17	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1000	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1001	3	-	0/6/23/26	0/1/1/1
3	BMA	A	1002	3	-	0/2/19/22	0/1/1/1
5	NAG	A	4000	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	4001	5	-	0/6/23/26	0/1/1/1
3	NAG	A	6000	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	6001	3	-	0/6/23/26	0/1/1/1
3	BMA	A	6002	3	-	0/2/19/22	0/1/1/1
5	NAG	A	8000	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	8001	5	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	9000	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	9001	5	-	0/6/23/26	0/1/1/1
3	NAG	B	2000	3,2	-	0/6/23/26	0/1/1/1
3	NAG	B	2001	3	-	0/6/23/26	0/1/1/1
3	BMA	B	2002	3	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	9000	NAG	C1-C2	2.15	1.55	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1000	NAG	C4-C3-C2	-2.88	106.75	111.23
3	B	2001	NAG	C4-C3-C2	-2.76	106.93	111.23
5	A	4000	NAG	C2-N2-C7	-2.76	119.49	123.04
3	A	6000	NAG	C4-C3-C2	-2.73	106.99	111.23
5	A	4000	NAG	C4-C3-C2	-2.65	107.11	111.23
3	A	6000	NAG	C2-N2-C7	-2.53	119.79	123.04
5	A	8001	NAG	C4-C3-C2	-2.41	107.48	111.23
5	A	9001	NAG	C4-C3-C2	-2.37	107.54	111.23
3	A	6001	NAG	C4-C3-C2	-2.35	107.57	111.23
5	A	8000	NAG	C3-C4-C5	-2.28	106.23	110.20
3	B	2000	NAG	C4-C3-C2	-2.26	107.71	111.23
5	A	4001	NAG	C4-C3-C2	-2.26	107.72	111.23
3	B	2000	NAG	C6-C5-C4	-2.21	107.56	113.02
5	A	8000	NAG	C1-O5-C5	-2.16	109.51	112.25
3	A	1001	NAG	C4-C3-C2	-2.11	107.94	111.23
3	B	2001	NAG	C8-C7-N2	2.11	120.14	116.11
3	A	6000	NAG	C8-C7-N2	2.12	120.15	116.11
5	A	4000	NAG	C8-C7-N2	2.17	120.26	116.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1000	NAG	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	9000	NAG	1	0
3	B	2000	NAG	1	0

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	3000	1	14,14,15	0.52	0	15,19,21	1.29	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	3000	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	3000	NAG	C2-N2-C7	-2.42	119.93	123.04
4	A	3000	NAG	C4-C3-C2	-2.26	107.72	111.23
4	A	3000	NAG	C8-C7-N2	2.12	120.16	116.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3000	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	614/663 (92%)	0.30	20 (3%)	50 22	47, 78, 116, 144	0
2	B	496/528 (93%)	0.11	8 (1%)	74 47	43, 61, 100, 129	0
All	All	1110/1191 (93%)	0.22	28 (2%)	61 30	43, 70, 112, 144	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	535	THR	6.0
2	B	531	LEU	4.4
2	B	530	GLU	3.8
1	A	377	ALA	3.5
2	B	504	LEU	3.3
1	A	543	ILE	3.2
1	A	536	GLU	3.2
1	A	540	ARG	3.1
1	A	541	GLY	2.9
1	A	38	HIS	2.8
2	B	529	PRO	2.8
2	B	349	ASP	2.6
1	A	532	LEU	2.6
2	B	515	SER	2.6
2	B	136	ARG	2.5
1	A	565	PHE	2.5
1	A	380	TYR	2.5
1	A	375	ALA	2.5
1	A	612	PHE	2.4
1	A	514	LEU	2.3
2	B	372	LEU	2.3
1	A	631	LYS	2.3
1	A	378	ALA	2.2
1	A	371	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	530	VAL	2.2
1	A	632	ASN	2.1
1	A	304	LEU	2.1
1	A	542	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NAG	A	9000	14/15	0.75	0.29	2.90	111,124,129,132	0
3	NAG	A	6000	14/15	0.90	0.28	0.45	112,118,126,134	0
3	NAG	A	1000	14/15	0.94	0.16	-0.61	65,82,91,103	0
3	NAG	B	2000	14/15	0.95	0.18	-0.65	61,78,89,99	0
5	NAG	A	8000	14/15	0.92	0.17	-0.84	88,93,101,109	0
3	BMA	A	6002	11/12	0.65	0.41	-	151,153,158,159	0
3	NAG	A	6001	14/15	0.79	0.47	-	138,143,149,150	0
5	NAG	A	4001	14/15	0.78	0.32	-	124,130,140,141	0
3	NAG	B	2001	14/15	0.87	0.25	-	94,110,118,124	0
5	NAG	A	9001	14/15	0.83	0.23	-	126,133,139,141	0
3	BMA	A	1002	11/12	0.65	0.23	-	123,128,131,132	0
3	BMA	B	2002	11/12	0.79	0.20	-	125,128,131,131	0
5	NAG	A	4000	14/15	0.89	0.38	-	114,118,122,125	0
5	NAG	A	8001	14/15	0.85	0.23	-	112,119,121,123	0
3	NAG	A	1001	14/15	0.89	0.26	-	107,116,121,125	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	A	3000	14/15	0.86	0.23	-	131,133,141,142	0

6.5 Other polymers [i](#)

There are no such residues in this entry.