



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:18 AM GMT

PDB ID : 3ONW
Title : Structure of a G-alpha-i1 mutant with enhanced affinity for the RGS14 GoLoco motif.
Authors : Bosch, D.; Kimple, A.J.; Sammond, D.W.; Miley, M.J.; Machius, M.; Kuhlman, B.; Willard, F.S.; Siderovski, D.P.
Deposited on : 2010-08-30
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	<div> <div>11%</div> <div>88%</div> <div>8%</div> <div>•</div> </div>
1	B	328	<div> <div>17%</div> <div>90%</div> <div>7%</div> <div>•</div> </div>
2	C	36	<div> <div>22%</div> <div>72%</div> <div>25%</div> <div>•</div> </div>
2	D	36	<div> <div>33%</div> <div>83%</div> <div>11%</div> <div>6%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5811 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(i) subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2543	1617	428	483	15			
1	B	317	Total	C	N	O	S	0	0	0
			2551	1623	429	484	15			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	SER	-	EXPRESSION TAG	UNP P63096
A	28	ASN	-	EXPRESSION TAG	UNP P63096
A	29	ALA	-	EXPRESSION TAG	UNP P63096
A	30	GLY	-	EXPRESSION TAG	UNP P63096
A	147	LEU	GLN	ENGINEERED MUTATION	UNP P63096
B	27	SER	-	EXPRESSION TAG	UNP P63096
B	28	ASN	-	EXPRESSION TAG	UNP P63096
B	29	ALA	-	EXPRESSION TAG	UNP P63096
B	30	GLY	-	EXPRESSION TAG	UNP P63096
B	147	LEU	GLN	ENGINEERED MUTATION	UNP P63096

- Molecule 2 is a protein called Regulator of G-protein signaling 14.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	35	Total	C	N	O	0	0	0
			279	175	51	53			
2	D	34	Total	C	N	O	0	0	0
			270	168	49	53			

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 28	C 10	N 5	O 11	P 2	0	0
3	B	1	Total 28	C 10	N 5	O 11	P 2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	52	Total	O	0	0
			52	52		
5	B	40	Total	O	0	0
			40	40		
5	C	2	Total	O	0	0
			2	2		
5	D	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Guanine nucleotide-binding protein G(i) subunit alpha-1




- Molecule 1: Guanine nucleotide-binding protein G(i) subunit alpha-1



- Molecule 2: Regulator of G-protein signaling 14



- Molecule 2: Regulator of G-protein signaling 14



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	70.39 Å 83.68 Å 190.15 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	95.07 – 2.38 28.88 – 2.38	Depositor EDS
% Data completeness (in resolution range)	98.5 (95.07-2.38) 98.6 (28.88-2.38)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.39 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.228 , 0.265 0.220 , 0.255	Depositor DCC
R_{free} test set	1528 reflections (3.49%)	DCC
Wilson B-factor (Å ²)	53.7	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 45283 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5811	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2589	0.55	0/3488
1	B	0.46	0/2597	0.53	0/3499
2	C	0.43	0/281	0.52	0/377
2	D	0.42	0/272	0.56	0/365
All	All	0.47	0/5739	0.54	0/7729

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2543	0	2522	12	0
1	B	2551	0	2533	11	0
2	C	279	0	286	5	0
2	D	270	0	271	1	0
3	A	28	0	12	0	0
3	B	28	0	12	1	0
4	A	5	0	0	0	0
4	B	10	0	0	0	0
5	A	52	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	40	0	0	0	0
5	C	2	0	0	0	0
5	D	3	0	0	0	0
All	All	5811	0	5636	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:LEU:HB2	1:B:218:VAL:HG11	1.76	0.67
1:A:81:ILE:O	1:A:85:ILE:HG12	1.99	0.61
1:B:61:TYR:H	1:B:171:GLN:HE22	1.53	0.55
1:A:277:LYS:HA	1:A:280:LYS:HE2	1.89	0.55
1:B:251:ASP:OD1	1:B:255:ASN:ND2	2.41	0.53
1:A:251:ASP:OD1	1:A:255:ASN:ND2	2.40	0.53
1:A:248:LYS:HD3	2:C:503:LEU:HD21	1.92	0.51
1:B:143:SER:HA	1:B:146:TYR:CE1	2.48	0.48
1:B:186:GLU:HG2	1:B:187:THR:N	2.29	0.48
1:A:208:ARG:O	1:A:212:ILE:HG12	2.15	0.47
1:A:230:TYR:O	1:A:286:CYS:HB2	2.16	0.44
1:B:254:CYS:SG	1:B:319:ILE:HD11	2.57	0.44
1:A:36:LEU:CD2	1:A:220:ALA:HB3	2.48	0.44
1:B:255:ASN:O	1:B:313:ARG:NH2	2.51	0.43
1:A:36:LEU:HD21	1:A:220:ALA:HB3	2.00	0.43
2:C:522:GLU:O	2:C:525:VAL:HG12	2.17	0.43
1:A:254:CYS:SG	1:A:319:ILE:HD11	2.58	0.43
2:C:497:ILE:HG13	2:C:499:GLY:H	1.83	0.43
1:A:42:GLY:HA3	2:C:510:SER:O	2.18	0.42
1:B:270:LYS:HG2	3:B:401:GDP:C6	2.55	0.42
2:C:513:HIS:C	2:C:515:GLN:H	2.23	0.41
1:A:240:MET:HG3	1:A:245:GLU:HG3	2.03	0.41
1:A:115:GLU:HG3	1:A:118:PHE:HB3	2.02	0.41
1:B:86:ARG:NH2	2:D:525:VAL:O	2.52	0.40
1:B:230:TYR:O	1:B:286:CYS:HB2	2.21	0.40
1:B:52:GLN:HG2	1:B:175:LEU:HD21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/328 (96%)	308 (98%)	5 (2%)	1 (0%)	46	61
1	B	315/328 (96%)	306 (97%)	9 (3%)	0	100	100
2	C	33/36 (92%)	32 (97%)	1 (3%)	0	100	100
2	D	32/36 (89%)	30 (94%)	2 (6%)	0	100	100
All	All	694/728 (95%)	676 (97%)	17 (2%)	1 (0%)	56	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	114	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/283 (97%)	269 (98%)	6 (2%)	60	78
1	B	276/283 (98%)	272 (99%)	4 (1%)	74	88
2	C	31/32 (97%)	30 (97%)	1 (3%)	46	66
2	D	30/32 (94%)	27 (90%)	3 (10%)	9	12
All	All	612/630 (97%)	598 (98%)	14 (2%)	58	76

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	ASP
1	A	106	GLN
1	A	181	THR
1	A	209	LYS
1	A	246	SER
1	A	258	TRP
1	B	94	ASP
1	B	100	ARG
1	B	214	CYS
1	B	318	GLU
2	C	498	GLU
2	D	496	ASP
2	D	513	HIS
2	D	526	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	GLN
1	B	171	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SO4	A	3	-	4,4,4	0.20	0	6,6,6	0.06	0
3	GDP	A	401	-	23,30,30	1.23	3 (13%)	30,47,47	1.91	7 (23%)
4	SO4	B	1	-	4,4,4	0.20	0	6,6,6	0.18	0
4	SO4	B	2	-	4,4,4	0.15	0	6,6,6	0.06	0
3	GDP	B	401	-	23,30,30	1.06	2 (8%)	30,47,47	1.94	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	3	-	-	0/0/0/0	0/0/0/0
3	GDP	A	401	-	-	0/12/32/32	0/3/3/3
4	SO4	B	1	-	-	0/0/0/0	0/0/0/0
4	SO4	B	2	-	-	0/0/0/0	0/0/0/0
3	GDP	B	401	-	-	0/12/32/32	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	GDP	O4'-C1'	2.31	1.44	1.41
3	B	401	GDP	C6-C5	2.67	1.46	1.41
3	B	401	GDP	C5-C4	2.82	1.46	1.40
3	A	401	GDP	C5-C4	2.87	1.47	1.40
3	A	401	GDP	C6-C5	3.48	1.48	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	GDP	C5-C6-N1	-4.29	117.73	123.59
3	A	401	GDP	C5-C6-N1	-4.04	118.06	123.59
3	A	401	GDP	C6-C5-C4	-3.76	116.41	120.90
3	B	401	GDP	C6-C5-C4	-3.70	116.48	120.90
3	A	401	GDP	PA-O3A-PB	-3.48	120.99	132.67
3	B	401	GDP	C1'-N9-C4	-3.46	121.72	126.94
3	A	401	GDP	N3-C2-N1	-3.37	122.31	127.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	GDP	N3-C2-N1	-3.24	122.51	127.44
3	A	401	GDP	C1'-N9-C4	-3.04	122.36	126.94
3	B	401	GDP	PA-O3A-PB	-2.71	123.57	132.67
3	B	401	GDP	C4-C5-N7	-2.59	107.10	109.48
3	A	401	GDP	C4-C5-N7	-2.56	107.13	109.48
3	B	401	GDP	O4'-C1'-N9	-2.14	103.63	108.10
3	B	401	GDP	C6-N1-C2	5.04	122.93	115.94
3	A	401	GDP	C6-N1-C2	5.04	122.94	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	GDP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/328 (96%)	0.66	37 (11%) 6 7	33, 59, 85, 97	0
1	B	317/328 (96%)	0.89	56 (17%) 2 2	31, 60, 90, 102	0
2	C	35/36 (97%)	1.24	8 (22%) 1 1	60, 78, 91, 101	0
2	D	34/36 (94%)	1.33	12 (35%) 0 0	51, 80, 102, 108	0
All	All	702/728 (96%)	0.82	113 (16%) 3 3	31, 61, 90, 108	0

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	348	LEU	8.1
1	B	113	ALA	6.3
1	B	182	THR	6.2
1	B	315	ASP	5.8
1	A	182	THR	5.6
1	A	315	ASP	5.5
1	B	118	PHE	4.9
2	C	512	ALA	4.8
1	B	59	ALA	4.8
2	D	525	VAL	4.7
1	B	190	THR	4.7
1	A	193	ASP	4.3
1	A	222	ILE	4.2
1	B	195	HIS	4.1
1	B	112	GLY	4.0
1	A	168	ILE	4.0
1	B	183	GLY	4.0
1	A	205	ARG	3.8
1	B	203	GLY	3.6
1	B	347	ASN	3.6
1	B	344	ILE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	109	VAL	3.6
1	A	113	ALA	3.4
1	B	92	LYS	3.4
1	B	209	LYS	3.4
2	D	529	PHE	3.3
1	B	32	ARG	3.3
2	D	496	ASP	3.2
2	D	523	ASP	3.2
2	D	522	GLU	3.2
1	B	63	GLU	3.1
1	B	94	ASP	3.1
1	A	207	GLU	3.1
1	B	345	LYS	3.0
2	C	525	VAL	3.0
1	B	193	ASP	2.9
1	A	90	ARG	2.9
1	B	224	CYS	2.9
2	D	524	LEU	2.9
1	A	330	LYS	2.8
1	B	267	PHE	2.8
2	C	522	GLU	2.8
1	B	67	LYS	2.8
1	B	181	THR	2.8
1	B	314	LYS	2.8
2	C	502	GLU	2.8
1	B	115	GLU	2.7
1	A	218	VAL	2.7
1	A	39	LEU	2.7
1	A	186	GLU	2.7
1	B	180	LYS	2.7
2	D	528	GLU	2.6
1	B	205	ARG	2.6
2	C	514	ASP	2.5
1	A	69	TYR	2.5
2	C	520	ARG	2.5
1	B	90	ARG	2.5
1	A	225	VAL	2.5
1	A	32	ARG	2.5
2	D	518	LEU	2.4
1	B	225	VAL	2.4
1	A	210	LYS	2.4
1	B	64	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	313	ARG	2.4
1	B	97	ASP	2.4
1	A	61	TYR	2.4
1	B	75	SER	2.4
1	A	316	THR	2.4
1	A	166	ASN	2.3
1	B	127	ILE	2.3
1	A	190	THR	2.3
1	B	320	TYR	2.3
1	A	237	ASP	2.3
1	B	196	PHE	2.3
1	B	199	PHE	2.3
2	D	521	LYS	2.3
1	A	266	LEU	2.3
1	B	222	ILE	2.3
1	B	69	TYR	2.3
1	A	38	LEU	2.3
1	B	194	LEU	2.3
1	B	207	GLU	2.3
1	A	41	ALA	2.3
1	A	124	ALA	2.3
1	B	96	GLY	2.3
1	B	124	ALA	2.3
1	B	41	ALA	2.2
1	B	316	THR	2.2
1	B	220	ALA	2.2
1	A	68	GLN	2.2
1	B	108	PHE	2.2
1	A	192	LYS	2.2
1	A	59	ALA	2.2
2	D	498	GLU	2.2
1	A	44	SER	2.1
1	A	224	CYS	2.1
2	C	521	LYS	2.1
1	B	47	SER	2.1
1	A	239	GLU	2.1
1	A	209	LYS	2.1
2	D	520	ARG	2.1
1	B	188	HIS	2.1
1	B	266	LEU	2.1
1	A	123	LEU	2.1
1	B	73	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	166	ASN	2.0
2	C	505	ASN	2.0
1	A	97	ASP	2.0
1	A	204	GLN	2.0
2	D	514	ASP	2.0
1	B	62	SER	2.0
1	B	123	LEU	2.0
1	A	195	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	B	2	5/5	0.93	0.24	1.94	99,100,100,100	0
3	GDP	B	401	28/28	0.96	0.23	1.00	41,46,49,49	0
3	GDP	A	401	28/28	0.98	0.16	-0.21	40,44,48,49	0
4	SO4	B	1	5/5	0.91	0.26	-	84,85,85,86	0
4	SO4	A	3	5/5	0.91	0.29	-	120,120,121,121	0

6.5 Other polymers [i](#)

There are no such residues in this entry.