



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 04:13 PM BST

PDB ID : 1OQ6
Title : solution structure of Copper-S46V CopA from Bacillus subtilis
Authors : Banci, L.; Bertini, I.; Ciofi-Baffoni, S.; Gonnelli, I.; Su, X.C.; Structural Proteomics in Europe (SPINE)
Deposited on : 2003-03-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 52%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	76	

2 Ensemble composition and analysis i

This entry contains 30 models. Model 22 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:73 (71)	0.26	22

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 5, 6, 8, 10, 11, 15, 19, 20, 21, 22, 24, 25, 26, 27
2	3, 4, 7, 16, 18, 28, 29, 30
3	2, 12, 17
Single-model clusters	9; 13; 14; 23

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1148 atoms, of which 580 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Potential copper-transporting ATPase.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	76	1147	350	580	98	113	6	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	VAL	SER	ENGINEERED	UNP O32220
A	73	ILE	THR	CLONING ARTIFACT	UNP O32220
A	75	GLY	LYS	CLONING ARTIFACT	UNP O32220
A	76	ARG	ALA	CLONING ARTIFACT	UNP O32220

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	
			Total	Cu
2	A	1	1	1

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Potential copper-transporting ATPase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Potential copper-transporting ATPase




4.2.2 Score per residue for model 2

- Molecule 1: Potential copper-transporting ATPase




4.2.3 Score per residue for model 3

- Molecule 1: Potential copper-transporting ATPase




4.2.4 Score per residue for model 4

- Molecule 1: Potential copper-transporting ATPase




4.2.5 Score per residue for model 5

- Molecule 1: Potential copper-transporting ATPase




4.2.6 Score per residue for model 6

- Molecule 1: Potential copper-transporting ATPase




4.2.7 Score per residue for model 7

- Molecule 1: Potential copper-transporting ATPase




4.2.8 Score per residue for model 8

- Molecule 1: Potential copper-transporting ATPase




4.2.9 Score per residue for model 9

- Molecule 1: Potential copper-transporting ATPase




4.2.10 Score per residue for model 10

- Molecule 1: Potential copper-transporting ATPase




4.2.11 Score per residue for model 11

- Molecule 1: Potential copper-transporting ATPase




4.2.12 Score per residue for model 12

- Molecule 1: Potential copper-transporting ATPase




4.2.13 Score per residue for model 13

- Molecule 1: Potential copper-transporting ATPase




4.2.14 Score per residue for model 14

- Molecule 1: Potential copper-transporting ATPase




4.2.15 Score per residue for model 15

- Molecule 1: Potential copper-transporting ATPase




4.2.16 Score per residue for model 16

- Molecule 1: Potential copper-transporting ATPase




4.2.17 Score per residue for model 17

- Molecule 1: Potential copper-transporting ATPase




4.2.18 Score per residue for model 18

- Molecule 1: Potential copper-transporting ATPase




4.2.19 Score per residue for model 19

- Molecule 1: Potential copper-transporting ATPase




4.2.20 Score per residue for model 20

- Molecule 1: Potential copper-transporting ATPase




4.2.21 Score per residue for model 21

- Molecule 1: Potential copper-transporting ATPase




4.2.22 Score per residue for model 22 (medoid)

- Molecule 1: Potential copper-transporting ATPase




4.2.23 Score per residue for model 23

- Molecule 1: Potential copper-transporting ATPase




4.2.24 Score per residue for model 24

- Molecule 1: Potential copper-transporting ATPase




4.2.25 Score per residue for model 25

- Molecule 1: Potential copper-transporting ATPase



4.2.26 Score per residue for model 26

- Molecule 1: Potential copper-transporting ATPase




4.2.27 Score per residue for model 27

- Molecule 1: Potential copper-transporting ATPase




4.2.28 Score per residue for model 28

- Molecule 1: Potential copper-transporting ATPase




4.2.29 Score per residue for model 29

- Molecule 1: Potential copper-transporting ATPase



4.2.30 Score per residue for model 30

- Molecule 1: Potential copper-transporting ATPase



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry, simulated annealing, molecular dynamics, torsion angle dynamics,*.

Of the 300 calculated structures, 30 were deposited, based on the following criterion: *structures with lowest energy violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Xeasy	structure solution	1.3
DYANA	structure solution	1.5
AMBER	refinement	5.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5768
Number of chemical shift lists	1
Total number of shifts	544
Number of shifts mapped to atoms	544
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	52%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.48±0.01	0±0/531 (0.0±0.0%)	0.94±0.02	1±0/719 (0.1±0.1%)
All	All	0.48	0/15930 (0.0%)	0.94	17/21570 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.0±1.0
All	All	0	60

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	56	GLY	C-N-CA	6.86	138.84	121.70	4	17

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	50	TYR	Sidechain	27
1	A	56	GLY	Peptide	14
1	A	69	TYR	Sidechain	11
1	A	55	THR	Peptide	8

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	526	536	536	2±2
All	All	15810	16080	16080	52

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:39:VAL:HG22	1:A:41:LEU:HD23	0.73	1.57	4	7
1:A:55:THR:HG22	1:A:59:ALA:HB3	0.72	1.60	17	4
1:A:21:ALA:HB1	1:A:39:VAL:HG21	0.66	1.67	20	12
1:A:55:THR:CG2	1:A:59:ALA:HB3	0.54	2.31	2	2
1:A:31:MET:CE	1:A:59:ALA:HB1	0.52	2.35	2	2
1:A:31:MET:SD	1:A:59:ALA:HB1	0.50	2.47	14	1
1:A:12:VAL:HG21	1:A:41:LEU:HD21	0.47	1.85	23	2
1:A:8:ILE:N	1:A:8:ILE:HD12	0.46	2.26	10	2
1:A:8:ILE:C	1:A:8:ILE:HD12	0.46	2.31	26	1
1:A:8:ILE:HD12	1:A:8:ILE:N	0.46	2.26	3	3
1:A:8:ILE:HD11	1:A:48:VAL:CG1	0.45	2.41	26	1
1:A:39:VAL:CG2	1:A:41:LEU:HD23	0.45	2.42	25	2
1:A:20:CYS:SG	1:A:23:ARG:NH2	0.44	2.90	27	1
1:A:25:GLU:HG3	1:A:37:ALA:HB1	0.44	1.88	23	1
1:A:6:LYS:NZ	1:A:7:GLU:OE1	0.43	2.51	19	1
1:A:29:LYS:HE2	1:A:36:ASP:HA	0.43	1.90	17	1
1:A:8:ILE:HD11	1:A:48:VAL:HG13	0.43	1.90	26	1
1:A:8:ILE:HD13	1:A:50:TYR:CE1	0.42	2.49	10	1
1:A:41:LEU:H	1:A:41:LEU:HD22	0.42	1.74	17	1
1:A:24:ILE:HG21	1:A:69:TYR:CD1	0.41	2.50	12	3
1:A:22:ALA:HA	1:A:39:VAL:HG11	0.41	1.91	9	1
1:A:62:GLU:OE2	1:A:66:LYS:NZ	0.41	2.53	23	2

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	71/76 (93%)	62±2 (87±3%)	8±2 (11±3%)	1±1 (1±1%)	18	63
All	All	2130/2280 (93%)	1858 (87%)	241 (11%)	31 (1%)	18	63

All 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	57	THR	17
1	A	3	SER	5
1	A	44	GLU	3
1	A	56	GLY	3
1	A	16	THR	2
1	A	18	ALA	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	56/60 (93%)	42±2 (76±4%)	14±2 (24±4%)	3	27
All	All	1680/1800 (93%)	1269 (76%)	411 (24%)	3	27

All 41 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	38	ASN	30
1	A	61	GLN	28
1	A	41	LEU	28
1	A	69	TYR	28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	15	MET	27
1	A	16	THR	25
1	A	36	ASP	23
1	A	25	GLU	21
1	A	55	THR	20
1	A	5	GLN	19
1	A	35	THR	17
1	A	64	ILE	14
1	A	7	GLU	13
1	A	30	ARG	12
1	A	54	GLU	11
1	A	10	MET	10
1	A	11	GLN	10
1	A	63	LYS	10
1	A	29	LYS	8
1	A	66	LYS	7
1	A	62	GLU	6
1	A	60	ILE	5
1	A	31	MET	5
1	A	45	THR	4
1	A	6	LYS	4
1	A	23	ARG	3
1	A	70	HIS	3
1	A	8	ILE	2
1	A	12	VAL	2
1	A	3	SER	2
1	A	28	LEU	2
1	A	48	VAL	2
1	A	46	VAL	2
1	A	57	THR	1
1	A	40	ASN	1
1	A	71	VAL	1
1	A	50	TYR	1
1	A	26	LYS	1
1	A	4	GLU	1
1	A	44	GLU	1
1	A	65	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 52% for the well-defined parts and 50% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5768

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	544
Number of shifts mapped to atoms	544
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	—
$^{13}\text{C}_\beta$	0	—	—
$^{13}\text{C}'$	0	—	—
^{15}N	71	0.72 ± 0.40	None needed (imprecise)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 52%, i.e. 423 atoms were assigned a chemical shift out of a possible 813. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	205/351 (58%)	137/140 (98%)	0/142 (0%)	68/69 (99%)
Sidechain	214/439 (49%)	208/253 (82%)	0/169 (0%)	6/17 (35%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	4/23 (17%)	4/12 (33%)	0/10 (0%)	0/1 (0%)
Overall	423/813 (52%)	349/405 (86%)	0/321 (0%)	74/87 (85%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 50%, i.e. 439 atoms were assigned a chemical shift out of a possible 880. 0 out of 11 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	214/376 (57%)	143/150 (95%)	0/152 (0%)	71/74 (96%)
Sidechain	221/481 (46%)	215/278 (77%)	0/183 (0%)	6/20 (30%)
Aromatic	4/23 (17%)	4/12 (33%)	0/10 (0%)	0/1 (0%)
Overall	439/880 (50%)	362/440 (82%)	0/345 (0%)	77/95 (81%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

