



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:19 AM GMT

PDB ID : 3ORC  
Title : CRYSTAL STRUCTURE OF AN ENGINEERED CRO MONOMER BOUND  
NONSPECIFICALLY TO DNA  
Authors : Albright, R.A.; Mossing, M.C.; Matthews, B.W.  
Deposited on : 1998-04-23  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	8	<div> <div>25%</div> <div>13% 38% 50%</div> </div>
1	S	8	<div> <div>25%</div> <div>13% 38% 50%</div> </div>
2	A	65	<div> <div>18% 54% 25%</div> <div>.</div> </div>



## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*TP\*AP\*TP\*CP\*GP\*AP\*TP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	R	8	Total	C	N	O	P	0	0	0
			161	79	29	46	7			
1	S	8	Total	C	N	O	P	0	0	0
			161	79	29	46	7			

- Molecule 2 is a protein called PROTEIN (CRO REPRESSOR).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	65	Total	C	N	O	S	0	0	0
			501	317	88	95	1			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ASP	ASP	SEE REMARK 999	UNP P03040
A	58	GLY	GLY	SEE REMARK 999	UNP P03040
A	59	GLU	GLU	SEE REMARK 999	UNP P03040
A	60	VAL	VAL	SEE REMARK 999	UNP P03040
A	61	LYS	LYS	SEE REMARK 999	UNP P03040
A	62	PRO	PRO	SEE REMARK 999	UNP P03040
A	63	PHE	PHE	SEE REMARK 999	UNP P03040
A	64	PRO	PRO	SEE REMARK 999	UNP P03040
A	65	SER	SER	SEE REMARK 999	UNP P03040
A	66	ASN	ASN	SEE REMARK 999	UNP P03040

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		



### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(\*TP\*AP\*TP\*CP\*GP\*AP\*TP\*A)-3')



- Molecule 1: DNA (5'-D(\*TP\*AP\*TP\*CP\*GP\*AP\*TP\*A)-3')



- Molecule 2: PROTEIN (CRO REPRESSOR)





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.72Å 60.66Å 45.72Å 90.00° 112.70° 90.00°	Depositor
Resolution (Å)	20.00 – 3.00 24.62 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-3.00) 92.3 (24.62-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.28 (at 2.39Å)	Xtriage
Refinement program	TNT V. 5-EB	Depositor
R, $R_{free}$	(Not available) , (Not available) 0.212 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	43.2	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.16 , 91.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 4237 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	826	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	R	1.67	4/180 (2.2%)	2.07	7/276 (2.5%)
1	S	1.67	4/180 (2.2%)	2.07	7/276 (2.5%)
2	A	1.06	3/509 (0.6%)	1.62	10/685 (1.5%)
All	All	1.34	11/869 (1.3%)	1.84	24/1237 (1.9%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	59	GLU	CD-OE2	6.51	1.32	1.25
2	A	54	GLU	CD-OE2	6.27	1.32	1.25
1	R	4	DC	C3'-O3'	6.21	1.52	1.44
1	S	4	DC	C3'-O3'	6.21	1.52	1.44
1	R	2	DA	C3'-O3'	6.16	1.51	1.44
1	S	2	DA	C3'-O3'	6.16	1.51	1.44
1	R	3	DT	N1-C2	5.25	1.42	1.38
1	S	3	DT	N1-C2	5.25	1.42	1.38
1	R	7	DT	C1'-N1	-5.22	1.40	1.47
1	S	7	DT	C1'-N1	-5.22	1.40	1.47
2	A	53	GLU	CD-OE1	5.08	1.31	1.25

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	2	DA	P-O3'-C3'	10.45	132.24	119.70
1	S	2	DA	P-O3'-C3'	10.45	132.24	119.70
1	R	7	DT	O4'-C4'-C3'	-8.64	100.82	106.00
1	S	7	DT	O4'-C4'-C3'	-8.64	100.82	106.00
1	R	4	DC	O4'-C1'-N1	-8.14	102.30	108.00
1	S	4	DC	O4'-C1'-N1	-8.14	102.30	108.00
2	A	22	ASP	CB-CG-OD2	-7.49	111.56	118.30
2	A	22	ASP	CB-CG-OD1	7.13	124.72	118.30
2	A	4	ARG	NE-CZ-NH1	6.64	123.62	120.30
1	R	6	DA	P-O5'-C5'	-6.33	110.77	120.90

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	6	DA	P-O5'-C5'	-6.33	110.77	120.90
2	A	9	ASP	CB-CG-OD2	-6.23	112.69	118.30
2	A	5	ILE	CB-CA-C	-6.15	99.30	111.60
1	R	7	DT	P-O3'-C3'	5.66	126.49	119.70
1	S	7	DT	P-O3'-C3'	5.66	126.49	119.70
2	A	57	ASP	CB-CG-OD2	-5.66	113.21	118.30
1	S	7	DT	C6-C5-C7	-5.61	119.54	122.90
1	R	7	DT	C6-C5-C7	-5.55	119.57	122.90
2	A	35	HIS	CA-CB-CG	-5.42	104.38	113.60
2	A	61	LYS	C-N-CD	-5.38	108.77	120.60
2	A	9	ASP	CB-CG-OD1	5.30	123.07	118.30
2	A	57	ASP	CB-CG-OD1	5.17	122.95	118.30
1	R	7	DT	C1'-O4'-C4'	-5.03	105.07	110.10
1	S	7	DT	C1'-O4'-C4'	-5.03	105.07	110.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	161	0	93	14	0
1	S	161	0	93	15	0
2	A	501	0	497	73	0
3	A	3	0	0	0	0
All	All	826	0	683	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 71.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:16:GLN:HE22	2:A:31:ASN:HB2	1.20	1.00
2:A:31:ASN:HA	2:A:34:ILE:HD12	1.45	0.97

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:7:DT:H2''	1:S:8:DA:H5'	1.47	0.96
1:R:7:DT:H2''	1:R:8:DA:H5'	1.47	0.94
2:A:45:ASN:HB2	2:A:49:SER:OG	1.84	0.78
2:A:59:GLU:HB2	2:A:61:LYS:HE2	1.67	0.77
1:R:6:DA:H2'	1:R:7:DT:H71	1.68	0.74
2:A:56:LYS:HB3	2:A:61:LYS:HE3	1.70	0.74
1:S:6:DA:H2'	1:S:7:DT:H71	1.68	0.72
2:A:59:GLU:HB2	2:A:61:LYS:CE	2.19	0.72
2:A:10:TYR:CD2	2:A:19:THR:HG23	2.24	0.72
1:R:7:DT:H2'	1:R:8:DA:C8	2.26	0.71
2:A:8:LYS:O	2:A:11:ALA:HB3	1.90	0.71
1:S:7:DT:H2'	1:S:8:DA:C8	2.26	0.71
2:A:30:ILE:HG22	2:A:31:ASN:N	2.08	0.68
2:A:42:LEU:HA	2:A:51:TYR:O	1.95	0.67
2:A:59:GLU:HB2	2:A:61:LYS:NZ	2.08	0.67
2:A:41:PHE:C	2:A:42:LEU:HD12	2.15	0.67
2:A:43:THR:C	2:A:44:ILE:HD12	2.16	0.66
2:A:44:ILE:N	2:A:44:ILE:HD12	2.11	0.66
2:A:30:ILE:CG2	2:A:34:ILE:HD11	2.26	0.65
2:A:5:ILE:HG22	2:A:6:THR:H	1.62	0.65
1:R:6:DA:H2''	1:R:7:DT:O5'	1.98	0.64
1:S:7:DT:H4'	1:S:8:DA:OP1	1.98	0.64
2:A:17:THR:OG1	2:A:27:GLN:NE2	2.31	0.64
2:A:26:TYR:O	2:A:29:ALA:N	2.29	0.64
2:A:16:GLN:OE1	2:A:31:ASN:ND2	2.30	0.64
1:R:7:DT:H4'	1:R:8:DA:OP1	1.98	0.63
1:S:6:DA:H2''	1:S:7:DT:O5'	1.98	0.63
2:A:9:ASP:HB3	2:A:13:ARG:NH1	2.14	0.62
2:A:50:VAL:HG12	2:A:51:TYR:N	2.15	0.62
2:A:16:GLN:NE2	2:A:31:ASN:HB2	2.05	0.61
2:A:17:THR:HA	2:A:27:GLN:HE21	1.67	0.59
2:A:25:VAL:HG23	2:A:26:TYR:N	2.17	0.58
2:A:10:TYR:HD2	2:A:19:THR:HG23	1.66	0.58
1:S:6:DA:OP2	2:A:29:ALA:HB2	2.05	0.57
1:R:7:DT:H2''	1:R:8:DA:C5'	2.28	0.57
2:A:25:VAL:HG23	2:A:29:ALA:HB3	1.87	0.56
2:A:5:ILE:N	2:A:5:ILE:HD12	2.21	0.56
2:A:30:ILE:O	2:A:33:ALA:HB3	2.07	0.55
2:A:14:PHE:CD2	2:A:18:LYS:HD3	2.42	0.55
2:A:5:ILE:HG21	2:A:13:ARG:HH12	1.72	0.54
1:S:7:DT:H2''	1:S:8:DA:C5'	2.28	0.53

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:23:LEU:HB2	2:A:25:VAL:HG13	1.91	0.53
2:A:45:ASN:O	2:A:48:GLY:N	2.42	0.52
2:A:40:ILE:HG13	2:A:54:GLU:HB2	1.90	0.52
2:A:5:ILE:H	2:A:5:ILE:HD12	1.73	0.51
2:A:56:LYS:HB3	2:A:61:LYS:CE	2.38	0.51
2:A:10:TYR:CE2	2:A:19:THR:HG23	2.44	0.51
2:A:14:PHE:HB3	2:A:18:LYS:HD3	1.92	0.51
1:R:7:DT:C4	1:S:2:DA:N1	2.79	0.50
2:A:5:ILE:HD11	2:A:44:ILE:HD11	1.93	0.50
1:R:2:DA:N1	1:S:7:DT:C4	2.79	0.50
2:A:63:PHE:HA	2:A:64:PRO:HA	1.60	0.49
2:A:57:ASP:C	2:A:59:GLU:H	2.15	0.49
2:A:65:SER:O	2:A:65:SER:OG	2.30	0.49
2:A:55:VAL:O	2:A:55:VAL:HG12	2.13	0.48
2:A:17:THR:HA	2:A:27:GLN:NE2	2.30	0.46
2:A:30:ILE:HG23	2:A:34:ILE:HD11	1.96	0.46
2:A:31:ASN:CA	2:A:34:ILE:HD12	2.32	0.46
2:A:60:VAL:O	2:A:60:VAL:HG22	2.16	0.46
1:R:7:DT:O4	1:S:2:DA:N1	2.49	0.46
1:R:2:DA:N1	1:S:7:DT:O4	2.49	0.46
2:A:40:ILE:HG22	2:A:41:PHE:N	2.31	0.46
2:A:47:ASP:OD1	2:A:47:ASP:N	2.34	0.45
2:A:9:ASP:CB	2:A:13:ARG:NH1	2.80	0.45
2:A:46:ALA:C	2:A:48:GLY:H	2.20	0.45
2:A:53:GLU:OE1	2:A:60:VAL:HG21	2.17	0.45
2:A:40:ILE:CG2	2:A:41:PHE:N	2.80	0.44
2:A:5:ILE:HG22	2:A:6:THR:N	2.29	0.44
2:A:42:LEU:HD23	2:A:50:VAL:HG11	1.99	0.44
2:A:50:VAL:CG1	2:A:51:TYR:N	2.80	0.44
2:A:25:VAL:CG2	2:A:29:ALA:HB3	2.48	0.44
2:A:44:ILE:N	2:A:44:ILE:CD1	2.79	0.43
2:A:30:ILE:O	2:A:33:ALA:N	2.52	0.43
1:R:8:DA:N1	1:S:1:DT:C4	2.87	0.43
2:A:17:THR:CA	2:A:27:GLN:HE21	2.32	0.43
2:A:30:ILE:HG22	2:A:34:ILE:CD1	2.49	0.43
1:S:2:DA:C2'	1:S:3:DT:H72	2.49	0.43
1:R:2:DA:C2'	1:R:3:DT:H72	2.49	0.43
1:R:1:DT:C4	1:S:8:DA:N1	2.87	0.42
2:A:50:VAL:HG12	2:A:51:TYR:H	1.81	0.42
2:A:14:PHE:HD2	2:A:18:LYS:HD3	1.81	0.42
2:A:40:ILE:HG22	2:A:42:LEU:HD12	2.02	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:S:1:DT:H2"	1:S:2:DA:OP2	2.20	0.42
2:A:30:ILE:HG22	2:A:34:ILE:HD11	1.99	0.41
2:A:59:GLU:HB2	2:A:61:LYS:HZ1	1.85	0.41
2:A:16:GLN:HE21	2:A:16:GLN:HB3	1.27	0.41
2:A:59:GLU:CB	2:A:61:LYS:NZ	2.80	0.41
2:A:17:THR:HG22	2:A:18:LYS:N	2.35	0.41
1:R:1:DT:H2"	1:R:2:DA:OP2	2.20	0.41
2:A:9:ASP:HB3	2:A:13:ARG:CZ	2.50	0.41
2:A:38:ARG:O	2:A:40:ILE:HD12	2.21	0.40
2:A:20:ALA:HA	2:A:30:ILE:HD11	2.03	0.40
2:A:57:ASP:C	2:A:59:GLU:N	2.74	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	63/65 (97%)	42 (67%)	18 (29%)	3 (5%)	<b>3</b> <b>17</b>

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	15	GLY
2	A	46	ALA
2	A	56	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	51/53 (96%)	37 (72%)	14 (28%)	0	2

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	3	GLN
2	A	4	ARG
2	A	5	ILE
2	A	6	THR
2	A	9	ASP
2	A	13	ARG
2	A	27	GLN
2	A	28	SER
2	A	43	THR
2	A	49	SER
2	A	51	TYR
2	A	60	VAL
2	A	65	SER
2	A	66	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	16	GLN
2	A	27	GLN
2	A	31	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	R	8/8 (100%)	0.93	2 (25%) 1 1	52, 64, 77, 79	8 (100%)
1	S	8/8 (100%)	0.93	2 (25%) 1 1	52, 64, 77, 79	8 (100%)
2	A	65/65 (100%)	-0.87	0 100 100	27, 49, 83, 88	0
All	All	81/81 (100%)	-0.51	4 (4%) 33 13	27, 54, 80, 88	16 (19%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	8	DA	4.7
1	S	8	DA	4.7
1	R	1	DT	2.6
1	S	1	DT	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.