



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:20 AM GMT

PDB ID : 3ORQ  
Title : Crystal Structure of N5-Carboxyaminoimidazole synthetase from Staphylococcus aureus complexed with ADP  
Authors : Brugarolas, P.; Duguid, E.M.; Zhang, W.; Poor, C.B.; He, C.  
Deposited on : 2010-09-07  
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1611 (2.26-2.22)
Clashscore	102246	1764 (2.26-2.22)
Ramachandran outliers	100387	1724 (2.26-2.22)
Sidechain outliers	100360	1724 (2.26-2.22)
RSRZ outliers	91569	1616 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	<div> <div>4%</div> <div>79%</div> <div>18%</div> <div>..</div> </div>
1	B	377	<div> <div>6%</div> <div>81%</div> <div>14%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ADP	B	375	-	-	-	X
4	POP	A	377	-	-	-	X
5	GOL	B	377	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N5-Carboxyaminoimidazole ribonucleotide synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	Se	0	1	0
			2990	1896	503	575	6	10			
1	B	366	Total	C	N	O	S	Se	0	0	0
			2922	1853	491	564	6	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP A6QFS4
A	-1	ASN	-	EXPRESSION TAG	UNP A6QFS4
A	0	ALA	-	EXPRESSION TAG	UNP A6QFS4
B	-2	SER	-	EXPRESSION TAG	UNP A6QFS4
B	-1	ASN	-	EXPRESSION TAG	UNP A6QFS4
B	0	ALA	-	EXPRESSION TAG	UNP A6QFS4

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 4 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: H<sub>2</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			9	7	2		
4	B	1	Total	O	P	0	0
			9	7	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	77	Total	O	0	0
			77	77		
6	B	105	Total	O	0	0
			105	105		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N5-Carboxyaminoimidazole ribonucleotide synthetase



- Molecule 1: N5-Carboxyaminoimidazole ribonucleotide synthetase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.41Å 53.45Å 110.39Å 90.00° 94.70° 90.00°	Depositor
Resolution (Å)	44.10 – 2.23 39.58 – 2.23	Depositor EDS
% Data completeness (in resolution range)	(Not available) (44.10-2.23) 98.1 (39.58-2.23)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.22Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.171 , 0.216 0.190 , 0.232	Depositor DCC
$R_{free}$ test set	1891 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.6	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 24.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 37291 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6174	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, POP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	2/3036 (0.1%)	0.72	1/4079 (0.0%)
1	B	0.75	0/2967	0.70	0/3988
All	All	0.76	2/6003 (0.0%)	0.71	1/8067 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	153	GLY	C-N	-5.90	1.20	1.34
1	A	90	CYS	CB-SG	-5.43	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	ASP	CB-CG-OD2	5.22	123.00	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2990	0	2970	52	0
1	B	2922	0	2896	38	0
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
3	A	27	0	12	0	0
3	B	27	0	12	1	0
4	A	9	0	0	0	0
4	B	9	0	0	0	0
5	B	6	0	8	0	0
6	A	77	0	0	1	0
6	B	105	0	0	1	0
All	All	6174	0	5898	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:ASP:HB3	6:B:463:HOH:O	1.68	0.93
1:B:134:ILE:HD11	1:B:175:ILE:HG13	1.55	0.86
1:A:274:ASN:HD21	1:A:314:ASN:HD21	1.23	0.86
1:B:101:ALA:O	1:B:105:LEU:HG	1.76	0.86
1:A:258:ASP:HB3	1:A:260:ASN:H	1.43	0.82
1:A:78:PHE:HB3	1:A:80:ASN:HD21	1.47	0.80
1:A:78:PHE:HB3	1:A:80:ASN:ND2	1.98	0.79
1:A:164:ASN:ND2	1:A:166:LYS:HB2	2.03	0.73
1:A:274:ASN:ND2	1:A:314:ASN:HD21	1.85	0.73
1:A:257:ILE:HD12	1:A:261:ASN:HA	1.72	0.72
1:B:335:LEU:HD21	1:B:352:MSE:HE2	1.71	0.71
1:B:115:LEU:HB3	1:B:122:VAL:HG21	1.73	0.69
1:A:21:GLY:N	1:A:77:GLU:HG3	2.08	0.68
1:B:177:THR:O	1:B:178:SER:HB2	1.92	0.68
1:B:373:SER:O	1:B:374:ASN:HB2	1.93	0.68
1:A:242:ILE:HD11	1:A:251:PHE:CZ	2.30	0.67
1:A:164:ASN:HD21	1:A:166:LYS:HB2	1.62	0.64
1:A:258:ASP:HB2	1:A:262:GLN:H	1.63	0.64
1:A:68:GLY:HA2	1:A:74:ILE:HD11	1.80	0.63
1:B:68:GLY:HA2	1:B:74:ILE:HD11	1.81	0.63
1:A:335:LEU:HD21	1:A:352:MSE:HE2	1.80	0.62
3:B:375:ADP:O2B	3:B:375:ADP:O2A	2.17	0.62
1:A:128:VAL:O	1:A:128:VAL:HG22	2.02	0.60
1:A:1:MSE:SE	1:A:202:ASN:HB2	2.52	0.59
1:A:328:ASN:HB2	1:B:298:GLN:OE1	2.03	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:335:LEU:HD21	1:A:352:MSE:CE	2.33	0.58
1:A:164:ASN:HD22	1:A:166:LYS:H	1.52	0.57
1:B:274:ASN:HD22	1:B:348:LYS:NZ	2.02	0.56
1:A:257:ILE:CD1	1:A:261:ASN:HA	2.33	0.56
1:A:257:ILE:HG13	1:A:257:ILE:O	2.05	0.56
1:B:134:ILE:O	1:B:138:ILE:HG22	2.06	0.56
1:A:242:ILE:HD11	1:A:251:PHE:HZ	1.71	0.56
1:B:115:LEU:HB3	1:B:122:VAL:CG2	2.37	0.53
1:B:315:LEU:HD21	1:B:352:MSE:HE3	1.91	0.51
1:B:150:ARG:NH1	1:B:179:GLU:OE1	2.44	0.51
1:B:3:PHE:CD1	1:B:3:PHE:C	2.85	0.50
1:A:258:ASP:HB3	1:A:260:ASN:N	2.20	0.50
1:A:36:VAL:HG22	1:A:52:GLU:HB3	1.94	0.50
1:A:34:LYS:HA	1:A:51:HIS:CE1	2.47	0.50
1:B:82:SER:HB2	1:B:84:GLN:HE21	1.77	0.49
1:B:274:ASN:ND2	1:B:314:ASN:HD21	2.11	0.49
1:A:189:LYS:HG2	1:A:258:ASP:O	2.12	0.49
1:B:3:PHE:HD1	1:B:3:PHE:C	2.15	0.49
1:B:165:GLU:O	1:B:168:LEU:HB2	2.13	0.48
1:B:274:ASN:HD21	1:B:314:ASN:HD21	1.60	0.48
1:B:82:SER:HB2	1:B:84:GLN:NE2	2.29	0.48
1:A:155:ASP:OD2	1:A:348:LYS:HE3	2.13	0.48
1:A:64:LEU:HB3	1:A:89:LEU:HD11	1.96	0.48
1:A:333:TRP:CE3	1:A:352:MSE:HE1	2.49	0.48
1:B:87:LYS:HG3	1:B:99:TYR:CE2	2.48	0.48
1:A:329:GLU:HG3	1:A:330:HIS:CD2	2.48	0.47
1:A:274:ASN:HD21	1:A:314:ASN:ND2	2.01	0.47
1:B:112:LYS:HB3	1:B:125:PHE:CZ	2.49	0.47
1:A:315:LEU:HD11	1:A:352:MSE:HE3	1.96	0.47
1:B:335:LEU:CD2	1:B:352:MSE:HE2	2.42	0.47
1:B:74:ILE:HB	1:B:95:ILE:HG12	1.97	0.47
1:B:161:LEU:HD12	1:B:162:ILE:H	1.79	0.47
1:A:144:PRO:HB2	1:A:185:TYR:HB3	1.97	0.47
1:A:211:GLU:O	1:A:222:THR:HA	2.16	0.46
1:A:20:LEU:HB2	1:A:77:GLU:HG2	1.97	0.46
1:A:122:VAL:HG12	1:A:123:VAL:O	2.16	0.45
1:A:111:GLU:O	1:A:115:LEU:HG	2.17	0.45
1:A:273:HIS:H	1:A:287:GLN:HE22	1.64	0.45
1:A:223:ILE:CG2	1:A:359:VAL:HG11	2.48	0.44
1:A:316:LEU:HA	1:A:347:ARG:O	2.17	0.44
1:B:291:HIS:O	1:B:295:VAL:HB	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:335:LEU:HD21	1:B:352:MSE:CE	2.43	0.44
1:B:102:ILE:O	1:B:106:GLN:HG3	2.17	0.44
1:B:126:ILE:HG22	1:B:182:ALA:HB3	1.98	0.43
1:A:27:SER:O	1:A:31[A]:MSE:HG3	2.19	0.43
1:A:78:PHE:CB	1:A:80:ASN:HD21	2.26	0.43
1:A:194:LEU:HD23	1:A:238:VAL:HG21	2.01	0.43
1:A:237:GLN:O	1:A:241:ILE:HG13	2.19	0.43
1:A:205:THR:HG22	1:A:304:ILE:HD12	2.01	0.43
1:B:58:TYR:CD2	1:B:80:ASN:HB2	2.54	0.43
1:B:6:LEU:O	1:B:298:GLN:NE2	2.52	0.42
1:B:104:LEU:CD1	1:B:110:THR:HG22	2.50	0.42
1:A:128:VAL:CG2	1:A:128:VAL:O	2.68	0.42
1:B:104:LEU:HD21	1:B:114:THR:OG1	2.20	0.42
1:A:104:LEU:C	1:A:104:LEU:HD23	2.40	0.42
1:B:123:VAL:HG22	1:B:186:LEU:HD11	2.01	0.41
1:A:150:ARG:HG2	1:A:179:GLU:O	2.19	0.41
1:A:196:VAL:O	1:A:250:THR:HA	2.20	0.41
1:A:44:CYS:HA	1:A:45:PRO:HD3	1.88	0.41
1:A:136:LYS:HD3	1:A:136:LYS:HA	1.79	0.41
1:A:155:ASP:OD1	6:A:407:HOH:O	2.22	0.41
1:A:241:ILE:HD13	1:A:268:ILE:HD11	2.03	0.41
1:B:196:VAL:O	1:B:250:THR:HA	2.21	0.41
1:B:260:ASN:O	1:B:261:ASN:HB2	2.22	0.40
1:B:174:LEU:HA	1:B:174:LEU:HD12	1.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	372/377 (99%)	360 (97%)	11 (3%)	1 (0%)	46 51

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	362/377 (96%)	347 (96%)	14 (4%)	1 (0%)	46	51
All	All	734/754 (97%)	707 (96%)	25 (3%)	2 (0%)	46	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	178	SER
1	A	258	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/324 (102%)	317 (96%)	14 (4%)	36	41
1	B	324/324 (100%)	315 (97%)	9 (3%)	51	61
All	All	655/648 (101%)	632 (96%)	23 (4%)	43	51

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	80	ASN
1	A	84	GLN
1	A	128	VAL
1	A	136	LYS
1	A	154	TYR
1	A	177	THR
1	A	198	ARG
1	A	210	GLN
1	A	257	ILE
1	A	261	ASN
1	A	295	VAL
1	A	312	MSE
1	A	328	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	3	PHE
1	B	26	GLN
1	B	85	GLN
1	B	115	LEU
1	B	128	VAL
1	B	138	ILE
1	B	149	THR
1	B	167	ASP
1	B	295	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	80	ASN
1	A	106	GLN
1	A	164	ASN
1	A	187	ASN
1	A	274	ASN
1	A	287	GLN
1	A	302	ASN
1	A	330	HIS
1	B	26	GLN
1	B	84	GLN
1	B	106	GLN
1	B	210	GLN
1	B	212	ASN
1	B	274	ASN
1	B	330	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	ADP	A	376	2	22,29,29	1.11	3 (13%)	27,45,45	2.17	5 (18%)
4	POP	A	377	-	8,8,8	0.61	0	13,13,13	1.34	1 (7%)
3	ADP	B	375	2	22,29,29	1.11	2 (9%)	27,45,45	1.93	6 (22%)
5	GOL	B	377	-	5,5,5	0.19	0	5,5,5	0.59	0
4	POP	B	378	-	8,8,8	0.98	0	13,13,13	1.10	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	376	2	-	0/12/32/32	0/3/3/3
4	POP	A	377	-	-	0/6/6/6	0/0/0/0
3	ADP	B	375	2	-	0/12/32/32	0/3/3/3
5	GOL	B	377	-	-	0/4/4/4	0/0/0/0
4	POP	B	378	-	-	0/6/6/6	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	376	ADP	C2-N3	2.14	1.36	1.32
3	A	376	ADP	O4'-C1'	2.30	1.44	1.41
3	B	375	ADP	O4'-C1'	2.65	1.44	1.41
3	B	375	ADP	C5-C4	3.01	1.47	1.40
3	A	376	ADP	C5-C4	3.08	1.47	1.40

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	376	ADP	N3-C2-N1	-7.07	123.48	128.89
3	B	375	ADP	N3-C2-N1	-7.01	123.53	128.89
3	A	376	ADP	C2'-C1'-N9	-6.57	104.25	114.29
4	A	377	POP	P2-O-P1	-3.69	122.35	132.73
3	B	375	ADP	PA-O3A-PB	-3.22	121.88	132.67
3	A	376	ADP	PA-O3A-PB	-2.69	123.66	132.67
3	B	375	ADP	C4-C5-N7	-2.23	107.43	109.48
4	B	378	POP	P2-O-P1	-2.20	126.56	132.73
3	B	375	ADP	C4'-O4'-C1'	2.01	111.93	109.72
4	B	378	POP	O5-P2-O	2.06	114.14	104.65
3	A	376	ADP	C2-N1-C6	2.14	122.59	118.77
3	B	375	ADP	O4'-C1'-N9	2.40	113.11	108.10
3	B	375	ADP	C2'-C3'-C4'	2.41	107.57	102.61
3	A	376	ADP	O2B-PB-O1B	2.57	118.84	110.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	375	ADP	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	364/377 (96%)	0.14	14 (3%) 44 46	2, 16, 51, 61	0
1	B	358/377 (94%)	0.05	21 (5%) 26 27	2, 9, 61, 72	0
All	All	722/754 (95%)	0.09	35 (4%) 34 36	2, 13, 56, 72	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	177	THR	9.8
1	B	175	ILE	5.4
1	B	171	GLY	4.1
1	B	178	SER	3.9
1	B	176	GLU	3.9
1	A	174	LEU	3.6
1	A	161	LEU	3.3
1	A	173	LYS	3.2
1	B	172	PHE	3.2
1	A	143	TYR	3.1
1	A	177	THR	3.0
1	A	176	GLU	2.8
1	B	132	THR	2.8
1	A	165	GLU	2.8
1	A	137	ALA	2.8
1	B	374	ASN	2.7
1	B	153	GLY	2.6
1	B	135	ASP	2.5
1	B	173	LYS	2.5
1	B	130	GLU	2.5
1	A	130	GLU	2.4
1	A	135	ASP	2.4
1	B	145	PHE	2.4
1	B	164	ASN	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	373	SER	2.2
1	B	181	VAL	2.2
1	B	165	GLU	2.2
1	A	145	PHE	2.2
1	A	167	ASP	2.1
1	B	185	TYR	2.1
1	B	136	LYS	2.1
1	A	160	VAL	2.1
1	B	84	GLN	2.1
1	A	184	LYS	2.0
1	B	139	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	B	377	6/6	0.84	0.39	7.44	41,58,63,67	0
3	ADP	B	375	27/27	0.62	0.45	6.18	50,66,76,77	27
4	POP	A	377	9/9	0.83	0.21	2.06	76,79,88,96	0
3	ADP	A	376	27/27	0.91	0.17	0.15	48,59,66,69	0
2	MG	A	375	1/1	0.67	0.38	-	46,46,46,46	0
4	POP	B	378	9/9	0.89	0.22	-	51,55,67,73	0
2	MG	B	376	1/1	0.87	0.27	-	69,69,69,69	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.