



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:58 PM GMT

PDB ID : 4ORI
Title : Rat dihydroorotate dehydrogenase bound with DSM338 (N-[3,5-difluoro-4-(trifluoromethyl)phenyl]-5-methyl-2-(trifluoromethyl)[1,2,4]triazolo[1,5-a]pyrimidin-7-amine)
Authors : Deng, X.; Phillips, M.A.
Deposited on : 2014-02-11
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dihydroorotate dehydrogenase (quinone), mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	355	2707	1701	495	508	3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	397	LEU	-	EXPRESSION TAG	UNP Q63707
A	398	GLU	-	EXPRESSION TAG	UNP Q63707
A	399	HIS	-	EXPRESSION TAG	UNP Q63707
A	400	HIS	-	EXPRESSION TAG	UNP Q63707
A	401	HIS	-	EXPRESSION TAG	UNP Q63707
A	402	HIS	-	EXPRESSION TAG	UNP Q63707
A	403	HIS	-	EXPRESSION TAG	UNP Q63707
A	404	HIS	-	EXPRESSION TAG	UNP Q63707

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is OROTIC ACID (three-letter code: ORO) (formula: $C_5H_4N_2O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			11	5	2	4		

- Molecule 4 is N-[3,5-DIFLUORO-4-(TRIFLUOROMETHYL)PHENYL]-5-METHYL-2-(TRIFLUOROMETHYL)[1,2,4]TRIAZOLO[1,5-A]PYRIMIDIN-7-AMINE (three-letter code: 2V6) (formula: $C_{14}H_7F_8N_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	0	0
			27	14	8	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	111	Total	O	0	0
			111	111		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (quinone), mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	124.36 Å 43.84 Å 63.30 Å 90.00° 99.96° 90.00°	Depositor
Resolution (Å)	20.00 – 1.50 23.08 – 1.46	Depositor EDS
% Data completeness (in resolution range)	95.4 (20.00-1.50) 95.5 (23.08-1.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 1.46 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.180 , 0.234 0.187 , 0.235	Depositor DCC
R_{free} test set	2659 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 68.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 56300 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2887	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ORO, 2V6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.33	11/2749 (0.4%)	0.94	12/3721 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	364	PRO	N-CD	5.79	1.55	1.47
1	A	216	PRO	N-CD	5.74	1.55	1.47
1	A	125	PRO	N-CD	5.61	1.55	1.47
1	A	138	PRO	N-CD	5.43	1.55	1.47
1	A	132	PRO	N-CD	5.34	1.55	1.47
1	A	177	PRO	N-CD	5.28	1.55	1.47
1	A	197	GLU	CD-OE1	-5.19	1.20	1.25
1	A	258	PRO	N-CD	5.16	1.55	1.47
1	A	52	PRO	N-CD	5.10	1.54	1.47
1	A	69	PRO	N-CD	5.05	1.54	1.47
1	A	44	PRO	N-CD	5.01	1.54	1.47

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	289	ARG	NE-CZ-NH1	-6.33	117.14	120.30
1	A	289	ARG	C-N-CD	5.89	140.78	128.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	328	ILE	C-N-CD	5.84	140.67	128.40
1	A	249	LYS	C-N-CD	5.84	140.66	128.40
1	A	68	LEU	C-N-CD	5.81	140.61	128.40
1	A	43	MET	C-N-CD	5.45	139.84	128.40
1	A	51	ASP	C-N-CD	5.33	139.59	128.40
1	A	215	SER	C-N-CD	5.32	139.56	128.40
1	A	137	LEU	C-N-CD	5.25	139.44	128.40
1	A	257	ALA	C-N-CD	5.21	139.34	128.40
1	A	364	PRO	C-N-CD	5.19	139.29	128.40
1	A	124	GLN	C-N-CD	5.07	139.04	128.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	244	LEU	Peptide
1	A	246	GLY	Peptide
1	A	248	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2707	0	2778	75	0
2	A	31	0	19	0	0
3	A	11	0	3	0	0
4	A	27	0	6	3	0
5	A	111	0	0	3	0
All	All	2887	0	2806	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ASP:OD2	1:A:293:LEU:HA	1.66	0.94
1:A:315:GLN:HE22	1:A:318:ARG:HH22	1.14	0.93
1:A:229:GLU:OE1	1:A:229:GLU:N	2.01	0.92
1:A:234:LEU:HD13	1:A:277:ILE:HD11	1.52	0.92
1:A:271:VAL:HA	1:A:274:GLU:HG2	1.61	0.83
1:A:315:GLN:NE2	1:A:318:ARG:HH22	1.79	0.80
1:A:57:ARG:HH22	1:A:131:ARG:HH22	1.28	0.79
1:A:247:THR:O	1:A:248:ARG:NH2	2.16	0.79
1:A:246:GLY:HA2	1:A:248:ARG:HH22	1.53	0.74
1:A:246:GLY:HA2	1:A:248:ARG:NH2	2.03	0.73
1:A:246:GLY:C	1:A:248:ARG:HH22	1.91	0.73
1:A:246:GLY:CA	1:A:248:ARG:HH22	2.02	0.72
1:A:36:HIS:N	5:A:702:HOH:O	2.21	0.72
1:A:89:ARG:HD3	1:A:176:LEU:HD21	1.71	0.71
1:A:273:ARG:HG2	1:A:273:ARG:NH1	2.05	0.69
1:A:244:LEU:CD2	1:A:245:LYS:HE2	2.21	0.69
1:A:244:LEU:HD22	1:A:245:LYS:HE2	1.77	0.66
1:A:139:GLU:HB3	1:A:294:GLN:HB2	1.77	0.65
1:A:53:GLU:HG2	1:A:57:ARG:NH2	2.12	0.65
1:A:158:GLU:OE2	1:A:162:ARG:HG3	1.97	0.65
1:A:234:LEU:HD13	1:A:277:ILE:CD1	2.27	0.63
1:A:37:PHE:HA	1:A:41:TYR:HD2	1.63	0.62
1:A:278:ASP:O	1:A:329:PRO:HD2	2.00	0.62
1:A:273:ARG:HG2	1:A:273:ARG:HH11	1.63	0.60
1:A:271:VAL:HA	1:A:274:GLU:CG	2.29	0.60
1:A:53:GLU:HG2	1:A:57:ARG:HH22	1.67	0.59
1:A:388:ASP:O	1:A:395:ARG:NH1	2.34	0.59
1:A:261:THR:O	1:A:265:LYS:HD3	2.03	0.59
1:A:261:THR:HB	1:A:264:ASP:OD1	2.02	0.59
1:A:261:THR:HG21	5:A:711:HOH:O	2.02	0.58
1:A:80:GLU:OE2	1:A:87:LYS:HD2	2.03	0.57
1:A:37:PHE:HA	1:A:41:TYR:CD2	2.39	0.57
1:A:264:ASP:OD1	1:A:264:ASP:N	2.38	0.56
1:A:273:ARG:HG3	1:A:328:ILE:HD11	1.88	0.55
1:A:123:PRO:HA	1:A:154:LEU:HD12	1.88	0.55
1:A:136:ARG:C	1:A:137:LEU:HD12	2.26	0.55
1:A:273:ARG:HH11	1:A:273:ARG:CG	2.21	0.54
1:A:265:LYS:HE3	1:A:319:GLU:OE2	2.08	0.53
1:A:100:LYS:HE2	5:A:710:HOH:O	2.09	0.52
1:A:260:LEU:HB2	1:A:265:LYS:HD2	1.92	0.51
1:A:36:HIS:ND1	1:A:36:HIS:N	2.59	0.50
1:A:139:GLU:CB	1:A:294:GLN:HB2	2.40	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ARG:HA	1:A:217:ASN:ND2	2.26	0.50
1:A:57:ARG:NH2	1:A:131:ARG:HH22	2.04	0.49
1:A:137:LEU:CD1	1:A:137:LEU:N	2.75	0.49
1:A:44:PRO:O	1:A:47:GLN:HB3	2.12	0.48
1:A:137:LEU:HD12	1:A:137:LEU:N	2.29	0.48
1:A:271:VAL:CA	1:A:274:GLU:HG2	2.38	0.48
1:A:36:HIS:HD1	1:A:36:HIS:N	2.12	0.48
1:A:57:ARG:HH22	1:A:131:ARG:NH2	2.05	0.47
1:A:356:TYR:CD1	1:A:356:TYR:C	2.89	0.47
1:A:40:GLU:OE1	1:A:40:GLU:N	2.48	0.47
1:A:261:THR:HB	1:A:264:ASP:CG	2.36	0.46
1:A:324:THR:HB	1:A:328:ILE:HG13	1.97	0.45
1:A:315:GLN:NE2	1:A:315:GLN:HA	2.32	0.45
1:A:315:GLN:HE22	1:A:318:ARG:NH2	1.96	0.45
1:A:203:GLY:N	1:A:204:PRO:CD	2.80	0.45
1:A:57:ARG:HH12	1:A:131:ARG:HH12	1.63	0.45
1:A:37:PHE:O	1:A:41:TYR:HB2	2.17	0.45
1:A:393:ASP:O	1:A:396:ARG:HD2	2.17	0.44
1:A:307:LYS:N	1:A:308:PRO:CD	2.80	0.44
1:A:56:HIS:HE1	1:A:98:PHE:O	2.00	0.44
1:A:270:SER:O	1:A:274:GLU:HG2	2.18	0.44
1:A:53:GLU:OE1	1:A:133:ARG:NH1	2.44	0.43
1:A:150:ASN:OD1	1:A:150:ASN:N	2.52	0.43
1:A:261:THR:HG22	1:A:263:GLN:N	2.34	0.42
4:A:502:2V6:H4	4:A:502:2V6:C2	2.47	0.42
1:A:282:VAL:HG12	1:A:283:THR:CG2	2.48	0.42
1:A:103:GLU:HA	1:A:160:ARG:HD3	2.02	0.41
1:A:282:VAL:HG12	1:A:283:THR:HG23	2.02	0.41
4:A:502:2V6:C9	4:A:502:2V6:C2	2.97	0.41
1:A:122:THR:O	1:A:153:GLY:HA2	2.20	0.41
1:A:261:THR:HG22	1:A:263:GLN:H	1.86	0.41
1:A:123:PRO:HA	1:A:154:LEU:CD1	2.51	0.40
1:A:59:ALA:HB2	4:A:502:2V6:C2	2.52	0.40
1:A:62:VAL:HG13	1:A:67:LEU:HB2	2.03	0.40
1:A:315:GLN:NE2	1:A:318:ARG:NH2	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/372 (94%)	344 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/302 (95%)	273 (95%)	15 (5%)	29	4

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	HIS
1	A	40	GLU
1	A	74	GLN
1	A	123	PRO
1	A	132	PRO
1	A	228	THR
1	A	235	SER
1	A	245	LYS
1	A	248	ARG
1	A	264	ASP
1	A	265	LYS
1	A	273	ARG
1	A	299	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	356	TYR
1	A	396	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS
1	A	315	GLN
1	A	347	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FMN	A	500	-	32,33,33	1.75	10 (31%)	34,50,50	2.61	11 (32%)
3	ORO	A	501	-	6,11,11	2.20	3 (50%)	6,15,15	3.84	6 (100%)
4	2V6	A	502	-	23,29,29	1.93	4 (17%)	32,46,46	4.51	11 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	500	-	-	0/18/18/18	0/3/3/3
3	ORO	A	501	-	-	0/0/4/4	0/1/1/1
4	2V6	A	502	-	-	0/16/16/16	0/3/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	2V6	C8-N1	-5.80	1.28	1.39
3	A	501	ORO	C2-N3	-3.56	1.30	1.38
2	A	500	FMN	C6-C5A	-3.49	1.36	1.41
3	A	501	ORO	C2-N1	-3.05	1.31	1.38
2	A	500	FMN	C1'-N10	-2.92	1.45	1.48
2	A	500	FMN	C10-N1	-2.79	1.30	1.35
2	A	500	FMN	P-O2P	-2.71	1.45	1.54
2	A	500	FMN	C2-N1	-2.67	1.32	1.38
2	A	500	FMN	P-O3P	-2.37	1.46	1.54
3	A	501	ORO	O4-C4	-2.17	1.19	1.24
2	A	500	FMN	C2-N3	-2.17	1.33	1.38
4	A	502	2V6	C1-N1	-2.06	1.36	1.40
4	A	502	2V6	C11-N4	-2.03	1.32	1.35
2	A	500	FMN	C9A-C5A	2.13	1.47	1.42
2	A	500	FMN	C4A-C10	2.31	1.45	1.40
2	A	500	FMN	C4-C4A	2.46	1.46	1.41
4	A	502	2V6	C4-C5	5.00	1.47	1.39

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	502	2V6	C9-C8-N1	-10.92	108.95	123.57
2	A	500	FMN	C4-C4A-C10	-7.36	115.23	119.94
4	A	502	2V6	C2-C3-C4	-4.03	121.17	124.27
3	A	501	ORO	N3-C2-N1	-3.86	121.19	127.69
2	A	500	FMN	C4A-C4-N3	-3.75	118.63	123.52
3	A	501	ORO	C5-C4-N3	-3.67	120.14	124.02
4	A	502	2V6	C6-C5-C4	-3.30	121.74	124.27
2	A	500	FMN	N3-C2-N1	-3.29	122.15	127.69
4	A	502	2V6	C9-C10-N3	-3.13	119.85	122.76
2	A	500	FMN	C4A-C10-N10	-3.00	118.34	120.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	ORO	C5-C6-N1	-2.82	118.93	122.40
4	A	502	2V6	F6-C14-C12	-2.25	106.13	112.80
4	A	502	2V6	C1-C6-C5	-2.02	117.31	118.83
4	A	502	2V6	C6-C1-N1	-2.01	114.26	120.44
2	A	500	FMN	O3P-P-O2P	2.19	115.49	107.44
2	A	500	FMN	C6-C5A-N5	2.19	121.65	118.92
4	A	502	2V6	C1-C2-C3	2.20	120.49	118.83
4	A	502	2V6	F7-C3-C2	2.29	123.00	118.58
3	A	501	ORO	C2-N1-C6	2.64	120.48	117.03
2	A	500	FMN	C1'-N10-C9A	2.78	122.05	118.83
2	A	500	FMN	C5A-C9A-N10	3.15	119.94	117.58
3	A	501	ORO	C4-C5-C6	3.25	118.83	116.73
2	A	500	FMN	C4-C4A-N5	3.44	122.88	118.70
2	A	500	FMN	C4A-N5-C5A	4.00	121.44	116.72
4	A	502	2V6	C8-C9-C10	4.72	120.23	116.92
3	A	501	ORO	C4-N3-C2	5.89	120.42	114.21
2	A	500	FMN	C4-N3-C2	8.84	122.53	115.16
4	A	502	2V6	N1-C8-N2	20.68	131.98	113.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	2V6	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/372 (95%)	1.00	72 (20%) ⓘ ⓘ	18, 38, 66, 84	4 (1%)

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	41	TYR	9.5
1	A	37	PHE	8.1
1	A	247	THR	6.7
1	A	296	ALA	4.8
1	A	232	HIS	4.8
1	A	248	ARG	4.7
1	A	246	GLY	4.6
1	A	36	HIS	4.5
1	A	263	GLN	4.5
1	A	40	GLU	4.3
1	A	128	GLY	4.2
1	A	130	PRO	4.2
1	A	327	ARG	4.1
1	A	272	ALA	3.8
1	A	322	ALA	3.7
1	A	270	SER	3.7
1	A	297	LEU	3.6
1	A	228	THR	3.5
1	A	274	GLU	3.5
1	A	187	THR	3.4
1	A	261	THR	3.4
1	A	132	PRO	3.4
1	A	273	ARG	3.2
1	A	209	LEU	3.2
1	A	293	LEU	3.2
1	A	118	VAL	3.1
1	A	242	ASP	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	149	PHE	3.1
1	A	291	VAL	3.1
1	A	395	ARG	3.1
1	A	268	ILE	3.1
1	A	218	THR	3.0
1	A	317	ILE	2.9
1	A	239	GLN	2.9
1	A	189	GLU	2.8
1	A	360	ILE	2.8
1	A	260	LEU	2.8
1	A	325	GLN	2.7
1	A	69	PRO	2.6
1	A	396	ARG	2.6
1	A	186	LYS	2.6
1	A	180	ILE	2.6
1	A	245	LYS	2.5
1	A	67	LEU	2.5
1	A	226	GLY	2.5
1	A	127	GLU	2.5
1	A	294	GLN	2.5
1	A	182	LEU	2.4
1	A	253	LEU	2.4
1	A	231	ARG	2.4
1	A	384	THR	2.4
1	A	299	SER	2.3
1	A	143	VAL	2.3
1	A	288	SER	2.3
1	A	298	ARG	2.3
1	A	210	VAL	2.3
1	A	287	VAL	2.3
1	A	295	GLY	2.3
1	A	70	ARG	2.2
1	A	202	LEU	2.2
1	A	348	ALA	2.2
1	A	281	ILE	2.2
1	A	323	LEU	2.2
1	A	331	ILE	2.2
1	A	148	GLY	2.1
1	A	359	LEU	2.1
1	A	271	VAL	2.1
1	A	355	LEU	2.1
1	A	188	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	211	VAL	2.1
1	A	131	ARG	2.1
1	A	350	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	2V6	A	502	27/27	0.95	0.10	-0.45	24,28,38,47	0
2	FMN	A	500	31/31	0.97	0.11	-0.67	16,20,25,27	0
3	ORO	A	501	11/11	0.90	0.09	-0.82	26,29,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.