



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:30 PM GMT

PDB ID : 4OX9
Title : Crystal structure of the aminoglycoside resistance methyltransferase NpmA bound to the 30S ribosomal subunit
Authors : Dunkle, J.A.; Conn, G.L.; Dunham, C.M.
Deposited on : 2014-02-04
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





















| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 91344 | 1317 (4.10-3.50) |
| Clashscore | 102246 | 1458 (4.10-3.50) |
| Ramachandran outliers | 100387 | 1397 (4.10-3.50) |
| Sidechain outliers | 100360 | 1392 (4.10-3.50) |
| RSRZ outliers | 91569 | 1325 (4.10-3.50) |
| RNA backbone | 2183 | 1070 (4.76-2.80) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|---|
| 1 | A | 1513 | <div> <div>2%</div> <div>32% 52% 13% .</div> </div> |
| 2 | B | 256 | <div> <div>3%</div> <div>25% 52% 12% . 9%</div> </div> |
| 3 | C | 239 | <div> <div>4%</div> <div>26% 46% 14% . 14%</div> </div> |
| 4 | D | 208 | <div> <div>%</div> <div>43% 52% .</div> </div> |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 5 | E | 161 |  |
| 6 | F | 101 |  |
| 7 | G | 155 |  |
| 8 | H | 138 |  |
| 9 | I | 128 |  |
| 10 | J | 104 |  |
| 11 | K | 129 |  |
| 12 | L | 131 |  |
| 13 | M | 126 |  |
| 14 | N | 60 |  |
| 15 | O | 88 |  |
| 16 | P | 88 |  |
| 17 | Q | 104 |  |
| 18 | R | 88 |  |
| 19 | S | 92 |  |
| 20 | T | 106 |  |
| 21 | V | 26 |  |
| 22 | Y | 222 |  |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|------|-----------|----------|---------|------------------|
| 23 | MG | A | 1601 | - | - | - | X |
| 23 | MG | A | 1607 | - | - | - | X |
| 23 | MG | A | 1609 | - | - | - | X |
| 23 | MG | A | 1611 | - | - | - | X |
| 23 | MG | A | 1616 | - | - | - | X |
| 23 | MG | A | 1624 | - | - | - | X |
| 23 | MG | A | 1629 | - | - | - | X |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|------|-----------|----------|---------|------------------|
| 23 | MG | A | 1634 | - | - | - | X |
| 23 | MG | A | 1637 | - | - | - | X |
| 23 | MG | A | 1639 | - | - | - | X |
| 23 | MG | A | 1645 | - | - | - | X |
| 23 | MG | A | 1651 | - | - | - | X |
| 23 | MG | A | 1655 | - | - | - | X |
| 23 | MG | A | 1661 | - | - | - | X |
| 23 | MG | A | 1663 | - | - | - | X |
| 23 | MG | A | 1669 | - | - | - | X |
| 23 | MG | A | 1670 | - | - | - | X |
| 23 | MG | A | 1681 | - | - | - | X |
| 23 | MG | A | 1700 | - | - | - | X |
| 23 | MG | A | 1701 | - | - | - | X |
| 23 | MG | A | 1702 | - | - | - | X |
| 23 | MG | A | 1703 | - | - | - | X |
| 23 | MG | A | 1704 | - | - | - | X |
| 23 | MG | A | 1713 | - | - | - | X |
| 23 | MG | A | 1715 | - | - | - | X |
| 23 | MG | N | 102 | - | - | - | X |

2 Entry composition

There are 25 unique types of molecules in this entry. The entry contains 53527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S rRNA.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-------|------|-------|------|---------|---------|-------|
| 1 | A | 1507 | Total | C | N | O | P | 0 | 0 | 0 |
| | | | 32394 | 14418 | 6002 | 10467 | 1507 | | | |

- Molecule 2 is a protein called 30S ribosomal protein S2.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 2 | B | 234 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1900 | 1213 | 341 | 341 | 5 | | | |

- Molecule 3 is a protein called 30S ribosomal protein S3.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 3 | C | 206 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1612 | 1016 | 314 | 281 | 1 | | | |

- Molecule 4 is a protein called 30S ribosomal protein S4.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 4 | D | 208 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1703 | 1066 | 339 | 291 | 7 | | | |

- Molecule 5 is a protein called 30S ribosomal protein S5.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 5 | E | 150 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1146 | 724 | 217 | 201 | 4 | | | |

- Molecule 6 is a protein called 30S ribosomal protein S6.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 6 | F | 101 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 843 | 531 | 155 | 154 | 3 | | | |

- Molecule 7 is a protein called 30S ribosomal protein S7.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 7 | G | 155 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1257 | 781 | 252 | 218 | 6 | | | |

- Molecule 8 is a protein called 30S ribosomal protein S8.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 8 | H | 138 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1116 | 705 | 215 | 193 | 3 | | | |

- Molecule 9 is a protein called 30S ribosomal protein S9.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 9 | I | 127 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1010 | 639 | 197 | 174 | | | | |

- Molecule 10 is a protein called 30S ribosomal protein S10.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 10 | J | 98 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 785 | 494 | 153 | 137 | 1 | | | |

- Molecule 11 is a protein called 30S ribosomal protein S11.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 11 | K | 119 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 885 | 549 | 168 | 165 | 3 | | | |

- Molecule 12 is a protein called 30S ribosomal protein S12.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 12 | L | 124 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 970 | 611 | 195 | 163 | 1 | | | |

- Molecule 13 is a protein called 30S ribosomal protein S13.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 13 | M | 125 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 997 | 617 | 207 | 171 | 2 | | | |

- Molecule 14 is a protein called 30S ribosomal protein S14 type Z.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|----|---|---------|---------|-------|
| 14 | N | 60 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 492 | 312 | 104 | 72 | 4 | | | |

- Molecule 15 is a protein called 30S ribosomal protein S15.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 15 | O | 88 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 734 | 459 | 147 | 126 | 2 | | | |

- Molecule 16 is a protein called 30S ribosomal protein S16.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 16 | P | 83 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 700 | 443 | 139 | 117 | 1 | | | |

- Molecule 17 is a protein called 30S ribosomal protein S17.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 17 | Q | 104 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 857 | 547 | 160 | 148 | 2 | | | |

- Molecule 18 is a protein called 30S ribosomal protein S18.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|----|---------|---------|-------|
| 18 | R | 73 | Total | C | N | O | 0 | 0 | 0 |
| | | | 598 | 381 | 118 | 99 | | | |

- Molecule 19 is a protein called 30S ribosomal protein S19.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 19 | S | 80 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 647 | 414 | 119 | 112 | 2 | | | |

- Molecule 20 is a protein called 30S ribosomal protein S20.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| 20 | T | 99 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 763 | 470 | 162 | 129 | 2 | | | |

- Molecule 21 is a protein called 30S ribosomal protein Thx.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---------|---------|-------|
| 21 | V | 24 | Total | C | N | O | 0 | 0 | 0 |
| | | | 208 | 128 | 50 | 30 | | | |

- Molecule 22 is a protein called 16S rRNA (adenine(1408)-N(1))-methyltransferase.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 22 | Y | 219 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1761 | 1138 | 294 | 328 | 1 | | | |


- Molecule 23 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 23 | B | 1 | Total | Mg | 0 | 0 |
| | | | 1 | 1 | | |
| 23 | A | 117 | Total | Mg | 0 | 0 |
| | | | 117 | 117 | | |
| 23 | N | 1 | Total | Mg | 0 | 0 |
| | | | 1 | 1 | | |
| 23 | E | 1 | Total | Mg | 0 | 0 |
| | | | 1 | 1 | | |

- Molecule 24 is ZINC ION (three-letter code: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 24 | D | 1 | Total | Zn | 0 | 0 |
| | | | 1 | 1 | | |
| 24 | N | 1 | Total | Zn | 0 | 0 |
| | | | 1 | 1 | | |

- Molecule 25 is SINEFUNGIN (three-letter code: SFG) (formula: C₁₅H₂₃N₇O₅).





| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---|---------|---------|
| 25 | Y | 1 | Total | C | N | O | 0 | 0 |
| | | | 27 | 15 | 7 | 5 | | |

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 16S rRNA







• Molecule 2: 30S ribosomal protein S2




- Molecule 3: 30S ribosomal protein S3




- Molecule 4: 30S ribosomal protein S4




- Molecule 5: 30S ribosomal protein S5





- Molecule 6: 30S ribosomal protein S6








• Molecule 7: 30S ribosomal protein S7






• Molecule 8: 30S ribosomal protein S8



• Molecule 9: 30S ribosomal protein S9



• Molecule 10: 30S ribosomal protein S10




• Molecule 11: 30S ribosomal protein S11






• Molecule 12: 30S ribosomal protein S12




• Molecule 13: 30S ribosomal protein S13



• Molecule 14: 30S ribosomal protein S14 type Z



• Molecule 15: 30S ribosomal protein S15





• Molecule 16: 30S ribosomal protein S16







• Molecule 17: 30S ribosomal protein S17




• Molecule 18: 30S ribosomal protein S18



• Molecule 19: 30S ribosomal protein S19



• Molecule 20: 30S ribosomal protein S20




• Molecule 21: 30S ribosomal protein Thx





● Molecule 22: 16S rRNA (adenine(1408)-N(1))-methyltransferase



4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 41 21 2 | Depositor |
| Cell constants a, b, c, α , β , γ | 403.52Å 403.52Å 176.61Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 49.72 – 3.80 49.71 – 3.80 | Depositor EDS |
| % Data completeness (in resolution range) | 99.7 (49.72-3.80) 94.1 (49.71-3.80) | Depositor EDS |
| R_{merge} | 0.12 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 2.16 (at 3.77Å) | Xtriage |
| Refinement program | PHENIX | Depositor |
| R, R_{free} | 0.233 , 0.257 0.238 , 0.262 | Depositor DCC |
| R_{free} test set | 7034 reflections (5.53%) | DCC |
| Wilson B-factor (Å ²) | 135.6 | Xtriage |
| Anisotropy | 0.316 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.26 , 81.2 | EDS |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| L-test for twinning ² | $\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$ | Xtriage |
| Outliers | 1 of 141757 reflections (0.001%) | Xtriage |
| F_o, F_c correlation | 0.92 | EDS |
| Total number of atoms | 53527 | wwPDB-VP |
| Average B, all atoms (Å ²) | 85.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SFG, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.49 | 4/36262 (0.0%) | 0.77 | 54/56597 (0.1%) |
| 2 | B | 0.42 | 0/1935 | 0.69 | 1/2609 (0.0%) |
| 3 | C | 0.43 | 0/1636 | 0.67 | 0/2205 |
| 4 | D | 0.44 | 0/1733 | 0.65 | 0/2318 |
| 5 | E | 0.69 | 2/1162 (0.2%) | 0.83 | 2/1564 (0.1%) |
| 6 | F | 0.43 | 0/856 | 0.64 | 0/1154 |
| 7 | G | 0.38 | 0/1276 | 0.62 | 0/1709 |
| 8 | H | 0.44 | 0/1136 | 0.74 | 0/1527 |
| 9 | I | 0.36 | 0/1029 | 0.62 | 0/1379 |
| 10 | J | 0.47 | 0/798 | 0.73 | 0/1073 |
| 11 | K | 0.39 | 0/900 | 0.70 | 0/1213 |
| 12 | L | 0.42 | 0/986 | 0.72 | 0/1320 |
| 13 | M | 0.40 | 0/1008 | 0.67 | 0/1347 |
| 14 | N | 0.40 | 0/501 | 0.78 | 0/664 |
| 15 | O | 0.36 | 0/745 | 0.63 | 1/992 (0.1%) |
| 16 | P | 0.43 | 0/716 | 0.76 | 0/963 |
| 17 | Q | 0.48 | 0/870 | 0.75 | 0/1159 |
| 18 | R | 0.43 | 0/604 | 0.64 | 0/801 |
| 19 | S | 0.35 | 0/661 | 0.72 | 1/890 (0.1%) |
| 20 | T | 0.45 | 0/765 | 0.74 | 0/1007 |
| 21 | V | 0.43 | 0/212 | 0.64 | 0/277 |
| 22 | Y | 0.49 | 0/1796 | 0.83 | 3/2418 (0.1%) |
| All | All | 0.47 | 6/57587 (0.0%) | 0.75 | 62/85186 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | A | 0 | 41 |

Continued on next page...

Continued from previous page...

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 22 | Y | 0 | 2 |
| All | All | 0 | 43 |

The worst 5 of 6 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|------|------|-------|-------|-------------|----------|
| 5 | E | 69 | VAL | C-N | 11.28 | 1.55 | 1.34 |
| 5 | E | 101 | ILE | C-N | 10.62 | 1.58 | 1.34 |
| 1 | A | 669 | U | O3'-P | 6.03 | 1.68 | 1.61 |
| 1 | A | 604 | G | O3'-P | -5.28 | 1.54 | 1.61 |
| 1 | A | 1405 | G | O3'-P | 5.17 | 1.67 | 1.61 |

The worst 5 of 62 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|------|------|-------------|--------|-------------|----------|
| 1 | A | 1406 | U | O5'-P-OP2 | -11.47 | 95.38 | 105.70 |
| 1 | A | 1498 | U | C2'-C3'-O3' | 9.91 | 131.29 | 109.50 |
| 1 | A | 1405 | G | P-O3'-C3' | -9.56 | 108.22 | 119.70 |
| 1 | A | 243 | A | C2'-C3'-O3' | 9.46 | 130.32 | 109.50 |
| 1 | A | 559 | A | C2'-C3'-O3' | 9.15 | 129.63 | 109.50 |

There are no chirality outliers.

5 of 43 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 1 | A | 12 | U | Sidechain |
| 1 | A | 197 | A | Sidechain |
| 1 | A | 203 | U | Sidechain |
| 1 | A | 220 | G | Sidechain |
| 1 | A | 231 | G | Sidechain |

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 32394 | 0 | 16349 | 1169 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 2 | B | 1900 | 0 | 1951 | 206 | 0 |
| 3 | C | 1612 | 0 | 1677 | 212 | 0 |
| 4 | D | 1703 | 0 | 1764 | 134 | 0 |
| 5 | E | 1146 | 0 | 1207 | 101 | 0 |
| 6 | F | 843 | 0 | 857 | 65 | 0 |
| 7 | G | 1257 | 0 | 1296 | 91 | 0 |
| 8 | H | 1116 | 0 | 1177 | 66 | 0 |
| 9 | I | 1010 | 0 | 1037 | 114 | 0 |
| 10 | J | 785 | 0 | 823 | 123 | 0 |
| 11 | K | 885 | 0 | 904 | 82 | 0 |
| 12 | L | 970 | 0 | 1057 | 106 | 0 |
| 13 | M | 997 | 0 | 1072 | 119 | 0 |
| 14 | N | 492 | 0 | 529 | 74 | 0 |
| 15 | O | 734 | 0 | 771 | 58 | 0 |
| 16 | P | 700 | 0 | 720 | 56 | 0 |
| 17 | Q | 857 | 0 | 928 | 85 | 0 |
| 18 | R | 598 | 0 | 670 | 61 | 0 |
| 19 | S | 647 | 0 | 673 | 83 | 0 |
| 20 | T | 763 | 0 | 860 | 88 | 0 |
| 21 | V | 208 | 0 | 221 | 26 | 0 |
| 22 | Y | 1761 | 0 | 1792 | 47 | 0 |
| 23 | A | 117 | 0 | 0 | 0 | 0 |
| 23 | B | 1 | 0 | 0 | 0 | 0 |
| 23 | E | 1 | 0 | 0 | 0 | 0 |
| 23 | N | 1 | 0 | 0 | 0 | 0 |
| 24 | D | 1 | 0 | 0 | 0 | 0 |
| 24 | N | 1 | 0 | 0 | 0 | 0 |
| 25 | Y | 27 | 0 | 22 | 3 | 0 |
| All | All | 53527 | 0 | 38357 | 2882 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 2882 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 9:I:57:GLY:O | 9:I:58:HIS:HD2 | 1.22 | 1.18 |
| 20:T:54:LYS:HG3 | 20:T:100:ILE:CD1 | 1.73 | 1.18 |
| 12:L:41:ARG:HG2 | 12:L:42:THR:H | 1.03 | 1.15 |
| 9:I:57:GLY:O | 9:I:58:HIS:CD2 | 2.00 | 1.14 |
| 2:B:77:ALA:HB2 | 2:B:211:ILE:HD13 | 1.22 | 1.14 |

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|----------|-------------|----|
| 2 | B | 232/256 (91%) | 174 (75%) | 34 (15%) | 24 (10%) | 1 | 12 |
| 3 | C | 204/239 (85%) | 135 (66%) | 41 (20%) | 28 (14%) | 0 | 6 |
| 4 | D | 206/208 (99%) | 166 (81%) | 31 (15%) | 9 (4%) | 3 | 35 |
| 5 | E | 148/161 (92%) | 130 (88%) | 13 (9%) | 5 (3%) | 5 | 43 |
| 6 | F | 99/101 (98%) | 79 (80%) | 19 (19%) | 1 (1%) | 19 | 66 |
| 7 | G | 153/155 (99%) | 127 (83%) | 16 (10%) | 10 (6%) | 1 | 25 |
| 8 | H | 136/138 (99%) | 125 (92%) | 7 (5%) | 4 (3%) | 6 | 46 |
| 9 | I | 125/128 (98%) | 88 (70%) | 27 (22%) | 10 (8%) | 1 | 19 |
| 10 | J | 96/104 (92%) | 59 (62%) | 20 (21%) | 17 (18%) | 0 | 3 |
| 11 | K | 117/129 (91%) | 88 (75%) | 20 (17%) | 9 (8%) | 1 | 20 |
| 12 | L | 122/131 (93%) | 97 (80%) | 16 (13%) | 9 (7%) | 1 | 21 |
| 13 | M | 123/126 (98%) | 88 (72%) | 27 (22%) | 8 (6%) | 1 | 25 |
| 14 | N | 58/60 (97%) | 40 (69%) | 11 (19%) | 7 (12%) | 0 | 8 |
| 15 | O | 86/88 (98%) | 70 (81%) | 11 (13%) | 5 (6%) | 2 | 28 |
| 16 | P | 81/88 (92%) | 65 (80%) | 15 (18%) | 1 (1%) | 16 | 63 |
| 17 | Q | 102/104 (98%) | 84 (82%) | 10 (10%) | 8 (8%) | 1 | 20 |
| 18 | R | 71/88 (81%) | 62 (87%) | 7 (10%) | 2 (3%) | 6 | 47 |
| 19 | S | 78/92 (85%) | 49 (63%) | 18 (23%) | 11 (14%) | 0 | 6 |
| 20 | T | 97/106 (92%) | 63 (65%) | 21 (22%) | 13 (13%) | 0 | 6 |
| 21 | V | 22/26 (85%) | 19 (86%) | 2 (9%) | 1 (4%) | 3 | 34 |
| 22 | Y | 217/222 (98%) | 206 (95%) | 6 (3%) | 5 (2%) | 8 | 51 |
| All | All | 2573/2750 (94%) | 2014 (78%) | 372 (14%) | 187 (7%) | 1 | 22 |

5 of 187 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | B | 9 | GLU |
| 2 | B | 15 | VAL |
| 2 | B | 16 | HIS |
| 2 | B | 17 | PHE |
| 2 | B | 21 | ARG |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|----------------|-----------|----------|-------------|----|
| 2 | B | 202/220 (92%) | 182 (90%) | 20 (10%) | 10 | 44 |
| 3 | C | 160/188 (85%) | 142 (89%) | 18 (11%) | 7 | 38 |
| 4 | D | 180/180 (100%) | 173 (96%) | 7 (4%) | 39 | 75 |
| 5 | E | 115/122 (94%) | 101 (88%) | 14 (12%) | 6 | 34 |
| 6 | F | 90/90 (100%) | 87 (97%) | 3 (3%) | 45 | 79 |
| 7 | G | 126/126 (100%) | 123 (98%) | 3 (2%) | 57 | 84 |
| 8 | H | 119/119 (100%) | 109 (92%) | 10 (8%) | 14 | 52 |
| 9 | I | 98/99 (99%) | 92 (94%) | 6 (6%) | 23 | 65 |
| 10 | J | 86/91 (94%) | 77 (90%) | 9 (10%) | 8 | 41 |
| 11 | K | 90/99 (91%) | 84 (93%) | 6 (7%) | 20 | 62 |
| 12 | L | 104/108 (96%) | 96 (92%) | 8 (8%) | 16 | 56 |
| 13 | M | 100/101 (99%) | 90 (90%) | 10 (10%) | 9 | 43 |
| 14 | N | 49/49 (100%) | 47 (96%) | 2 (4%) | 37 | 74 |
| 15 | O | 79/79 (100%) | 72 (91%) | 7 (9%) | 12 | 50 |
| 16 | P | 72/74 (97%) | 67 (93%) | 5 (7%) | 19 | 60 |
| 17 | Q | 96/96 (100%) | 90 (94%) | 6 (6%) | 22 | 63 |
| 18 | R | 64/77 (83%) | 62 (97%) | 2 (3%) | 47 | 80 |
| 19 | S | 71/79 (90%) | 68 (96%) | 3 (4%) | 36 | 74 |
| 20 | T | 76/82 (93%) | 70 (92%) | 6 (8%) | 15 | 55 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|-----|
| 21 | V | 19/21 (90%) | 19 (100%) | 0 | 100 | 100 |
| 22 | Y | 193/195 (99%) | 185 (96%) | 8 (4%) | 37 | 74 |
| All | All | 2189/2295 (95%) | 2036 (93%) | 153 (7%) | 19 | 60 |

5 of 153 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 8 | H | 85 | ARG |
| 10 | J | 64 | GLU |
| 20 | T | 73 | HIS |
| 8 | H | 92 | ARG |
| 9 | I | 53 | VAL |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 9 | I | 58 | HIS |

5.3.3 RNA ⓘ

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|-----------------|-------------------|-----------------|
| 1 | A | 1507/1513 (99%) | 229 (15%) | 90 (5%) |

5 of 229 RNA backbone outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 6 | G |
| 1 | A | 8 | A |
| 1 | A | 9 | G |
| 1 | A | 31 | G |
| 1 | A | 32 | A |

5 of 90 RNA pucker outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 1 | A | 687 | A |
| 1 | A | 976 | G |
| 1 | A | 1409 | C |
| 1 | A | 701 | C |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 840 | C |

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 123 ligands modelled in this entry, 122 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | # $ Z > 2$ | Counts | RMSZ | # $ Z > 2$ |
| 25 | SFG | Y | 301 | - | 21,29,29 | 1.01 | 2 (9%) | 19,42,42 | 2.30 | 4 (21%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 25 | SFG | Y | 301 | - | - | 0/9/33/33 | 0/3/3/3 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 25 | Y | 301 | SFG | C8-N7 | -2.05 | 1.30 | 1.34 |
| 25 | Y | 301 | SFG | O4'-C1' | 2.97 | 1.45 | 1.41 |

All (4) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 25 | Y | 301 | SFG | N3-C2-N1 | -8.43 | 122.44 | 128.89 |
| 25 | Y | 301 | SFG | C4-C5-N7 | -2.58 | 107.11 | 109.48 |
| 25 | Y | 301 | SFG | C4'-O4'-C1' | -2.52 | 106.95 | 109.72 |
| 25 | Y | 301 | SFG | C2'-C1'-N9 | -2.30 | 110.77 | 114.29 |

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 25 | Y | 301 | SFG | 3 | 0 |

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|----------------|-----------------------|-------|
| 1 | A | 1507/1513 (99%) | 0.08 | 24 (1%) 74 60 | 18, 71, 170, 260 | 0 |
| 2 | B | 234/256 (91%) | 0.16 | 8 (3%) 49 34 | 29, 94, 152, 190 | 0 |
| 3 | C | 206/239 (86%) | 0.08 | 10 (4%) 33 22 | 44, 104, 145, 171 | 0 |
| 4 | D | 208/208 (100%) | 0.05 | 2 (0%) 84 72 | 25, 79, 127, 182 | 0 |
| 5 | E | 150/161 (93%) | -0.15 | 0 100 100 | 18, 56, 106, 148 | 0 |
| 6 | F | 101/101 (100%) | -0.08 | 1 (0%) 84 72 | 48, 103, 142, 156 | 0 |
| 7 | G | 155/155 (100%) | -0.15 | 3 (1%) 70 54 | 40, 96, 148, 172 | 0 |
| 8 | H | 138/138 (100%) | 0.01 | 1 (0%) 89 80 | 11, 41, 81, 116 | 0 |
| 9 | I | 127/128 (99%) | 0.61 | 17 (13%) 4 4 | 40, 105, 144, 165 | 0 |
| 10 | J | 98/104 (94%) | 0.83 | 13 (13%) 4 4 | 55, 123, 165, 188 | 0 |
| 11 | K | 119/129 (92%) | 0.16 | 3 (2%) 61 44 | 36, 81, 129, 171 | 0 |
| 12 | L | 124/131 (94%) | 0.30 | 3 (2%) 62 46 | 18, 74, 120, 165 | 0 |
| 13 | M | 125/126 (99%) | 1.10 | 22 (17%) 2 2 | 56, 97, 137, 185 | 0 |
| 14 | N | 60/60 (100%) | 0.35 | 2 (3%) 50 34 | 44, 102, 136, 160 | 0 |
| 15 | O | 88/88 (100%) | 0.16 | 2 (2%) 64 48 | 17, 63, 122, 167 | 0 |
| 16 | P | 83/88 (94%) | 0.40 | 2 (2%) 62 46 | 21, 56, 95, 122 | 0 |
| 17 | Q | 104/104 (100%) | 0.42 | 5 (4%) 34 22 | 13, 56, 119, 198 | 0 |
| 18 | R | 73/88 (82%) | 0.27 | 4 (5%) 29 19 | 35, 82, 143, 191 | 0 |
| 19 | S | 80/92 (86%) | 0.51 | 8 (10%) 9 6 | 73, 124, 155, 197 | 0 |
| 20 | T | 99/106 (93%) | 0.14 | 2 (2%) 68 53 | 38, 80, 118, 156 | 0 |
| 21 | V | 24/26 (92%) | 0.68 | 4 (16%) 2 2 | 51, 88, 137, 141 | 0 |
| 22 | Y | 219/222 (98%) | 0.35 | 19 (8%) 13 8 | 64, 119, 160, 196 | 0 |
| All | All | 4122/4263 (96%) | 0.18 | 155 (3%) 44 30 | 11, 82, 153, 260 | 0 |

The worst 5 of 155 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 13 | M | 124 | PRO | 17.0 |
| 13 | M | 123 | ALA | 13.0 |
| 15 | O | 89 | GLY | 10.2 |
| 10 | J | 64 | GLU | 7.8 |
| 13 | M | 125 | ARG | 7.0 |

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | LLDF | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|------|-------|------|------|-------|----------------------------|-------|
| 23 | MG | A | 1651 | 1/1 | 0.86 | 1.11 | 47.91 | 8,8,8,8 | 0 |
| 23 | MG | A | 1703 | 1/1 | 0.92 | 1.06 | 44.92 | 70,70,70,70 | 0 |
| 23 | MG | A | 1713 | 1/1 | 0.72 | 0.81 | 39.84 | 39,39,39,39 | 0 |
| 23 | MG | A | 1637 | 1/1 | 0.87 | 0.97 | 38.25 | 70,70,70,70 | 0 |
| 23 | MG | A | 1629 | 1/1 | 0.85 | 0.94 | 30.64 | 70,70,70,70 | 0 |
| 23 | MG | A | 1700 | 1/1 | 0.85 | 0.94 | 25.44 | 10,10,10,10 | 0 |
| 23 | MG | A | 1607 | 1/1 | 0.76 | 0.57 | 24.79 | 61,61,61,61 | 0 |
| 23 | MG | A | 1669 | 1/1 | 0.40 | 0.81 | 20.22 | 33,33,33,33 | 0 |
| 23 | MG | A | 1616 | 1/1 | 0.78 | 0.71 | 20.20 | 19,19,19,19 | 0 |
| 23 | MG | A | 1624 | 1/1 | 0.65 | 0.88 | 16.98 | 70,70,70,70 | 0 |
| 23 | MG | A | 1634 | 1/1 | 0.96 | 0.56 | 15.44 | 3,3,3,3 | 0 |
| 23 | MG | A | 1702 | 1/1 | 0.91 | 0.46 | 12.62 | 32,32,32,32 | 0 |
| 23 | MG | A | 1609 | 1/1 | 0.89 | 0.53 | 12.22 | 33,33,33,33 | 0 |
| 23 | MG | A | 1663 | 1/1 | 0.95 | 0.42 | 11.49 | 16,16,16,16 | 0 |
| 23 | MG | A | 1601 | 1/1 | 0.24 | 0.59 | 10.95 | 42,42,42,42 | 0 |
| 23 | MG | N | 102 | 1/1 | 0.56 | 1.19 | 10.30 | 24,24,24,24 | 0 |
| 23 | MG | A | 1670 | 1/1 | 0.91 | 0.40 | 7.96 | 34,34,34,34 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | LLDF | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|------|-------|------|------|-------|-----------------------------|-------|
| 23 | MG | A | 1681 | 1/1 | 0.95 | 0.38 | 7.26 | 41,41,41,41 | 0 |
| 23 | MG | A | 1645 | 1/1 | 0.66 | 0.35 | 6.82 | 12,12,12,12 | 0 |
| 23 | MG | A | 1655 | 1/1 | 0.91 | 0.20 | 5.25 | 47,47,47,47 | 0 |
| 23 | MG | A | 1611 | 1/1 | 0.76 | 0.38 | 4.91 | 57,57,57,57 | 0 |
| 23 | MG | A | 1704 | 1/1 | 0.89 | 0.37 | 3.76 | 29,29,29,29 | 0 |
| 23 | MG | A | 1639 | 1/1 | 0.94 | 0.28 | 3.28 | 15,15,15,15 | 0 |
| 23 | MG | A | 1701 | 1/1 | 0.97 | 0.28 | 3.14 | 6,6,6,6 | 0 |
| 23 | MG | A | 1661 | 1/1 | 0.59 | 0.40 | 2.93 | 93,93,93,93 | 0 |
| 23 | MG | A | 1715 | 1/1 | 0.93 | 0.45 | 2.47 | 9,9,9,9 | 0 |
| 23 | MG | A | 1710 | 1/1 | 0.92 | 0.36 | 1.93 | 20,20,20,20 | 0 |
| 23 | MG | A | 1689 | 1/1 | 0.68 | 0.34 | 1.66 | 58,58,58,58 | 0 |
| 25 | SFG | Y | 301 | 27/27 | 0.87 | 0.36 | 0.45 | 110,110,110,110 | 0 |
| 23 | MG | A | 1659 | 1/1 | 0.98 | 0.23 | 0.25 | 12,12,12,12 | 0 |
| 23 | MG | B | 301 | 1/1 | 0.66 | 0.23 | -0.30 | 59,59,59,59 | 0 |
| 24 | ZN | N | 101 | 1/1 | 0.97 | 0.16 | -0.75 | 108,108,108,108 | 0 |
| 23 | MG | A | 1638 | 1/1 | 0.97 | 0.16 | -1.75 | 47,47,47,47 | 0 |
| 23 | MG | A | 1641 | 1/1 | 0.91 | 0.30 | - | 0,0,0,0 | 0 |
| 23 | MG | A | 1652 | 1/1 | 0.48 | 1.17 | - | 42,42,42,42 | 0 |
| 23 | MG | A | 1630 | 1/1 | 0.74 | 0.78 | - | 7,7,7,7 | 0 |
| 23 | MG | A | 1696 | 1/1 | 0.62 | 0.73 | - | 27,27,27,27 | 0 |
| 23 | MG | A | 1642 | 1/1 | 0.97 | 0.38 | - | 33,33,33,33 | 0 |
| 23 | MG | A | 1605 | 1/1 | 0.83 | 1.19 | - | 8,8,8,8 | 0 |
| 23 | MG | A | 1690 | 1/1 | 0.86 | 1.17 | - | 33,33,33,33 | 0 |
| 23 | MG | A | 1677 | 1/1 | 0.67 | 0.71 | - | 63,63,63,63 | 0 |
| 23 | MG | A | 1644 | 1/1 | 0.84 | 0.36 | - | 54,54,54,54 | 0 |
| 23 | MG | A | 1678 | 1/1 | 0.83 | 0.98 | - | 27,27,27,27 | 0 |
| 23 | MG | A | 1625 | 1/1 | 0.96 | 0.78 | - | 0,0,0,0 | 0 |
| 23 | MG | A | 1654 | 1/1 | 0.77 | 0.83 | - | 16,16,16,16 | 0 |
| 23 | MG | A | 1693 | 1/1 | 0.70 | 1.39 | - | 18,18,18,18 | 0 |
| 23 | MG | A | 1646 | 1/1 | 0.38 | 0.68 | - | 18,18,18,18 | 0 |
| 23 | MG | A | 1687 | 1/1 | 0.98 | 0.69 | - | 17,17,17,17 | 0 |
| 23 | MG | A | 1648 | 1/1 | 0.96 | 0.52 | - | 4,4,4,4 | 0 |
| 23 | MG | A | 1708 | 1/1 | 0.68 | 0.75 | - | 21,21,21,21 | 0 |
| 23 | MG | A | 1658 | 1/1 | 0.59 | 0.59 | - | 40,40,40,40 | 0 |
| 23 | MG | A | 1627 | 1/1 | 0.43 | 0.46 | - | 52,52,52,52 | 0 |
| 23 | MG | A | 1674 | 1/1 | 0.77 | 0.77 | - | 8,8,8,8 | 0 |
| 23 | MG | A | 1695 | 1/1 | 0.71 | 1.55 | - | 52,52,52,52 | 0 |
| 23 | MG | A | 1667 | 1/1 | 0.73 | 0.46 | - | 76,76,76,76 | 0 |
| 23 | MG | A | 1628 | 1/1 | 0.97 | 0.30 | - | 54,54,54,54 | 0 |
| 23 | MG | A | 1619 | 1/1 | 0.93 | 0.35 | - | 40,40,40,40 | 0 |
| 23 | MG | A | 1643 | 1/1 | 0.81 | 0.45 | - | 10,10,10,10 | 0 |
| 23 | MG | A | 1640 | 1/1 | 0.80 | 0.31 | - | 33,33,33,33 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | LLDF | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|------|-------|------|------|------|-----------------------------|-------|
| 23 | MG | A | 1686 | 1/1 | 0.89 | 0.30 | - | 62,62,62,62 | 0 |
| 23 | MG | A | 1603 | 1/1 | 0.49 | 0.79 | - | 36,36,36,36 | 0 |
| 23 | MG | A | 1712 | 1/1 | 0.91 | 0.51 | - | 36,36,36,36 | 0 |
| 23 | MG | A | 1632 | 1/1 | 0.82 | 0.79 | - | 27,27,27,27 | 0 |
| 23 | MG | A | 1675 | 1/1 | 0.52 | 0.36 | - | 43,43,43,43 | 0 |
| 23 | MG | A | 1613 | 1/1 | 0.79 | 0.91 | - | 63,63,63,63 | 0 |
| 23 | MG | A | 1673 | 1/1 | 0.91 | 0.34 | - | 3,3,3,3 | 0 |
| 23 | MG | A | 1610 | 1/1 | 0.98 | 0.24 | - | 63,63,63,63 | 0 |
| 23 | MG | A | 1623 | 1/1 | 0.92 | 1.39 | - | 30,30,30,30 | 0 |
| 23 | MG | A | 1662 | 1/1 | 0.51 | 0.49 | - | 67,67,67,67 | 0 |
| 23 | MG | A | 1709 | 1/1 | 0.86 | 0.35 | - | 34,34,34,34 | 0 |
| 23 | MG | A | 1660 | 1/1 | 0.94 | 0.29 | - | 4,4,4,4 | 0 |
| 23 | MG | A | 1617 | 1/1 | 0.90 | 0.20 | - | 76,76,76,76 | 0 |
| 23 | MG | A | 1664 | 1/1 | 0.98 | 0.26 | - | 7,7,7,7 | 0 |
| 23 | MG | A | 1716 | 1/1 | 0.72 | 0.32 | - | 42,42,42,42 | 0 |
| 23 | MG | A | 1691 | 1/1 | 0.96 | 0.22 | - | 2,2,2,2 | 0 |
| 23 | MG | A | 1614 | 1/1 | 0.84 | 0.78 | - | 59,59,59,59 | 0 |
| 23 | MG | A | 1622 | 1/1 | 0.90 | 0.35 | - | 29,29,29,29 | 0 |
| 23 | MG | A | 1647 | 1/1 | 0.72 | 0.73 | - | 70,70,70,70 | 0 |
| 23 | MG | A | 1688 | 1/1 | 0.53 | 0.85 | - | 28,28,28,28 | 0 |
| 23 | MG | A | 1679 | 1/1 | 0.96 | 0.19 | - | 27,27,27,27 | 0 |
| 23 | MG | A | 1705 | 1/1 | 0.77 | 0.81 | - | 16,16,16,16 | 0 |
| 23 | MG | A | 1620 | 1/1 | 0.65 | 1.90 | - | 15,15,15,15 | 0 |
| 23 | MG | A | 1676 | 1/1 | 0.91 | 0.41 | - | 5,5,5,5 | 0 |
| 23 | MG | E | 201 | 1/1 | 0.96 | 1.23 | - | 21,21,21,21 | 0 |
| 23 | MG | A | 1671 | 1/1 | 0.81 | 1.29 | - | 10,10,10,10 | 0 |
| 23 | MG | A | 1606 | 1/1 | 0.70 | 0.61 | - | 37,37,37,37 | 0 |
| 24 | ZN | D | 301 | 1/1 | 0.99 | 0.35 | - | 34,34,34,34 | 0 |
| 23 | MG | A | 1602 | 1/1 | 0.75 | 1.14 | - | 22,22,22,22 | 0 |
| 23 | MG | A | 1698 | 1/1 | 0.74 | 0.49 | - | 23,23,23,23 | 0 |
| 23 | MG | A | 1626 | 1/1 | 0.21 | 1.00 | - | 70,70,70,70 | 0 |
| 23 | MG | A | 1668 | 1/1 | 0.86 | 0.38 | - | 15,15,15,15 | 0 |
| 23 | MG | A | 1631 | 1/1 | 0.58 | 0.22 | - | 46,46,46,46 | 0 |
| 23 | MG | A | 1656 | 1/1 | 0.66 | 0.80 | - | 33,33,33,33 | 0 |
| 23 | MG | A | 1653 | 1/1 | 0.91 | 1.26 | - | 11,11,11,11 | 0 |
| 23 | MG | A | 1680 | 1/1 | 0.88 | 0.69 | - | 22,22,22,22 | 0 |
| 23 | MG | A | 1618 | 1/1 | 0.91 | 0.37 | - | 69,69,69,69 | 0 |
| 23 | MG | A | 1682 | 1/1 | 0.97 | 0.87 | - | 36,36,36,36 | 0 |
| 23 | MG | A | 1711 | 1/1 | 0.57 | 0.62 | - | 9,9,9,9 | 0 |
| 23 | MG | A | 1685 | 1/1 | 0.96 | 0.90 | - | 36,36,36,36 | 0 |
| 23 | MG | A | 1615 | 1/1 | 0.98 | 0.57 | - | 27,27,27,27 | 0 |
| 23 | MG | A | 1684 | 1/1 | 0.53 | 0.52 | - | 17,17,17,17 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | LLDF | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|------|-------|------|------|------|-----------------------------|-------|
| 23 | MG | A | 1649 | 1/1 | 0.75 | 0.81 | - | 15,15,15,15 | 0 |
| 23 | MG | A | 1635 | 1/1 | 0.81 | 0.55 | - | 25,25,25,25 | 0 |
| 23 | MG | A | 1608 | 1/1 | 0.96 | 0.09 | - | 56,56,56,56 | 0 |
| 23 | MG | A | 1706 | 1/1 | 0.91 | 0.46 | - | 30,30,30,30 | 0 |
| 23 | MG | A | 1621 | 1/1 | 0.56 | 0.58 | - | 59,59,59,59 | 0 |
| 23 | MG | A | 1666 | 1/1 | 0.70 | 0.24 | - | 26,26,26,26 | 0 |
| 23 | MG | A | 1665 | 1/1 | 0.24 | 0.84 | - | 107,107,107,107 | 0 |
| 23 | MG | A | 1697 | 1/1 | 0.79 | 0.73 | - | 38,38,38,38 | 0 |
| 23 | MG | A | 1604 | 1/1 | 0.60 | 1.96 | - | 70,70,70,70 | 0 |
| 23 | MG | A | 1694 | 1/1 | 0.68 | 0.94 | - | 14,14,14,14 | 0 |
| 23 | MG | A | 1650 | 1/1 | 0.79 | 0.91 | - | 57,57,57,57 | 0 |
| 23 | MG | A | 1633 | 1/1 | 0.99 | 0.52 | - | 27,27,27,27 | 0 |
| 23 | MG | A | 1707 | 1/1 | 0.83 | 0.83 | - | 42,42,42,42 | 0 |
| 23 | MG | A | 1657 | 1/1 | 0.98 | 0.73 | - | 0,0,0,0 | 0 |
| 23 | MG | A | 1692 | 1/1 | 0.88 | 0.41 | - | 23,23,23,23 | 0 |
| 23 | MG | A | 1672 | 1/1 | 0.97 | 0.08 | - | 15,15,15,15 | 0 |
| 23 | MG | A | 1714 | 1/1 | 0.91 | 0.88 | - | 19,19,19,19 | 0 |
| 23 | MG | A | 1612 | 1/1 | 0.36 | 0.33 | - | 46,46,46,46 | 0 |
| 23 | MG | A | 1699 | 1/1 | 0.72 | 0.58 | - | 43,43,43,43 | 0 |
| 23 | MG | A | 1600 | 1/1 | 0.87 | 0.11 | - | 63,63,63,63 | 0 |
| 23 | MG | A | 1683 | 1/1 | 0.95 | 0.39 | - | 38,38,38,38 | 0 |
| 23 | MG | A | 1636 | 1/1 | 0.98 | 0.47 | - | 1,1,1,1 | 0 |

6.5 Other polymers [i](#)

There are no such residues in this entry.