



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:01 AM GMT

PDB ID : 2OYI
Title : Crystal Structure of Fragment D of gammaD298,301A Fibrinogen with the Peptide Ligand Gly-Pro-Arg-Pro-Amide
Authors : Kostelansky, M.S.; Gorkun, O.V.; Lord, S.T.
Deposited on : 2007-02-22
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	66	<div> <div>8%</div> <div>45%</div> <div>48%</div> <div>5%</div> </div>
1	D	66	<div> <div>15%</div> <div>35%</div> <div>45%</div> <div>5%</div> <div>15%</div> </div>
2	B	313	<div> <div>3%</div> <div>61%</div> <div>33%</div> <div>• •</div> </div>
2	E	313	<div> <div>2%</div> <div>64%</div> <div>27%</div> <div>• 6%</div> </div>
3	C	311	<div> <div>4%</div> <div>66%</div> <div>27%</div> <div>• •</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	311	
4	G	4	
4	H	4	
4	I	4	
4	J	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	E	3	-	-	X	-
5	FUC	E	5	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10784 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibrinogen alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	65	Total	C	N	O	S	0	0	0
			530	327	100	100	3			
1	D	56	Total	C	N	O	S	0	0	0
			458	280	87	88	3			

- Molecule 2 is a protein called Fibrinogen beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	301	Total	C	N	O	S	0	0	0
			2414	1505	427	460	22			
2	E	294	Total	C	N	O	S	0	0	0
			2362	1475	417	448	22			

- Molecule 3 is a protein called Fibrinogen gamma chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	299	Total	C	N	O	S	0	0	0
			2393	1521	403	458	11			
3	F	285	Total	C	N	O	S	0	0	0
			2277	1448	384	434	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	298	ALA	ASP	ENGINEERED	UNP P02679
C	301	ALA	ASP	ENGINEERED	UNP P02679
F	298	ALA	ASP	ENGINEERED	UNP P02679
F	301	ALA	ASP	ENGINEERED	UNP P02679

- Molecule 4 is a protein called GPRP Peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	4	Total	C	N	O	0	0	0
			30	18	7	5			
4	H	4	Total	C	N	O	0	0	0
			30	18	7	5			
4	I	4	Total	C	N	O	0	0	0
			30	18	7	5			
4	J	4	Total	C	N	O	0	0	0
			30	18	7	5			

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	3	Total	C	N	O	0	0
			38	22	2	14		
5	E	3	Total	C	N	O	0	0
			38	22	2	14		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ca	0	0
			1	1		
6	C	1	Total	Ca	0	0
			1	1		
6	F	1	Total	Ca	0	0
			1	1		
6	E	1	Total	Ca	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total	O	0	0
			3	3		
7	B	27	Total	O	0	0
			27	27		
7	C	18	Total	O	0	0
			18	18		
7	D	5	Total	O	0	0
			5	5		
7	E	63	Total	O	0	0
			63	63		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	F	33	Total	O	0	0
			33	33		
7	I	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fibrinogen alpha chain




- Molecule 1: Fibrinogen alpha chain




- Molecule 2: Fibrinogen beta chain




- Molecule 2: Fibrinogen beta chain






• Molecule 3: Fibrinogen gamma chain



• Molecule 3: Fibrinogen gamma chain




• Molecule 4: GPRP Peptide



• Molecule 4: GPRP Peptide



• Molecule 4: GPRP Peptide



- Molecule 4: GPRP Peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.99Å 94.04Å 226.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.97 – 2.70 17.97 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.5 (17.97-2.70) 98.0 (17.97-2.69)	Depositor EDS
R_{merge}	0.00	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 2.70Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.216 , 0.256 0.218 , 0.257	Depositor DCC
R_{free} test set	2634 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 52264 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10784	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/531	0.51	0/709
1	D	0.30	0/458	0.53	0/610
2	B	0.35	0/2475	0.60	0/3342
2	E	0.38	0/2422	0.65	1/3268 (0.0%)
3	C	0.36	0/2459	0.57	0/3327
3	F	0.39	0/2340	0.63	0/3165
4	G	0.46	0/31	0.63	0/40
4	H	0.48	0/31	0.66	0/40
4	I	0.58	0/31	0.68	0/40
4	J	0.60	0/31	0.68	0/40
All	All	0.37	0/10809	0.60	1/14581 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	399	GLY	N-CA-C	6.25	128.73	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	530	0	556	50	0
1	D	458	0	477	43	0
2	B	2414	0	2279	115	0
2	E	2362	0	2231	92	0
3	C	2393	0	2248	84	0
3	F	2277	0	2140	105	0
4	G	30	0	32	5	0
4	H	30	0	32	6	0
4	I	30	0	32	4	0
4	J	30	0	32	4	0
5	B	38	0	34	5	0
5	E	38	0	34	11	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	E	1	0	0	0	0
6	F	1	0	0	0	0
7	A	3	0	0	0	0
7	B	27	0	0	4	0
7	C	18	0	0	2	0
7	D	5	0	0	0	0
7	E	63	0	0	1	0
7	F	33	0	0	2	0
7	I	1	0	0	0	0
All	All	10784	0	10127	448	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 448 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:3:NAG:H62	5:E:4:NAG:H2	1.30	1.11
2:B:397:GLU:HB3	2:B:431:THR:HG21	1.40	1.04
1:A:188:VAL:HG11	2:B:167:VAL:HG21	1.35	1.03
2:E:165:LEU:HD11	2:E:168:LEU:HD22	1.49	0.94
4:G:3:ARG:HG3	4:G:4:PRO:HD2	1.51	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	63/66 (96%)	59 (94%)	4 (6%)	0	100	100
1	D	54/66 (82%)	44 (82%)	9 (17%)	1 (2%)	10	25
2	B	299/313 (96%)	266 (89%)	30 (10%)	3 (1%)	19	45
2	E	292/313 (93%)	268 (92%)	21 (7%)	3 (1%)	19	45
3	C	297/311 (96%)	269 (91%)	23 (8%)	5 (2%)	11	29
3	F	283/311 (91%)	262 (93%)	17 (6%)	4 (1%)	14	35
4	G	2/4 (50%)	1 (50%)	1 (50%)	0	100	100
4	H	2/4 (50%)	1 (50%)	1 (50%)	0	100	100
4	I	2/4 (50%)	1 (50%)	1 (50%)	0	100	100
4	J	2/4 (50%)	2 (100%)	0	0	100	100
All	All	1296/1396 (93%)	1173 (90%)	107 (8%)	16 (1%)	16	39

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	435	VAL
3	C	97	GLU
3	C	301	ALA
1	D	137	GLN
2	E	281	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/61 (98%)	57 (95%)	3 (5%)	30	60
1	D	52/61 (85%)	49 (94%)	3 (6%)	25	52
2	B	259/271 (96%)	252 (97%)	7 (3%)	52	82
2	E	252/271 (93%)	244 (97%)	8 (3%)	46	77
3	C	250/257 (97%)	238 (95%)	12 (5%)	31	62
3	F	237/257 (92%)	227 (96%)	10 (4%)	36	68
4	G	3/3 (100%)	2 (67%)	1 (33%)	0	0
4	H	3/3 (100%)	3 (100%)	0	100	100
4	I	3/3 (100%)	2 (67%)	1 (33%)	0	0
4	J	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	1122/1190 (94%)	1076 (96%)	46 (4%)	37	69

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	382	THR
2	E	190	MET
3	F	383	THR
3	C	383	THR
1	D	172	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	307	HIS
1	D	187	GLN
3	F	307	HIS
3	C	317	ASN
3	C	325	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	3	2,5	14,14,15	0.68	0	15,19,21	0.79	1 (6%)
5	NAG	B	4	5	14,14,15	0.66	0	15,19,21	0.69	0
5	FUC	B	5	5	10,10,11	0.59	0	14,14,16	0.58	0
5	NAG	E	3	2,5	14,14,15	0.61	0	15,19,21	1.27	2 (13%)
5	NAG	E	4	5	14,14,15	0.73	0	15,19,21	0.79	0
5	FUC	E	5	5	10,10,11	0.60	0	14,14,16	0.95	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	3	2,5	-	0/6/23/26	0/1/1/1
5	NAG	B	4	5	-	0/6/23/26	0/1/1/1
5	FUC	B	5	5	-	0/0/17/20	0/1/1/1
5	NAG	E	3	2,5	-	0/6/23/26	0/1/1/1
5	NAG	E	4	5	-	0/6/23/26	0/1/1/1
5	FUC	E	5	5	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	3	NAG	C2-N2-C7	-2.85	119.37	123.04
5	B	3	NAG	C2-N2-C7	-2.50	119.83	123.04
5	E	5	FUC	C1-C2-C3	2.24	112.19	109.54
5	E	3	NAG	C3-C4-C5	2.66	114.84	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	3	NAG	5	0
5	B	5	FUC	1	0
5	E	3	NAG	9	0
5	E	4	NAG	5	0
5	E	5	FUC	6	0

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	65/66 (98%)	0.31	5 (7%) 16 14	24, 57, 85, 99	0
1	D	56/66 (84%)	0.77	10 (17%) 2 1	24, 68, 110, 115	0
2	B	301/313 (96%)	-0.06	8 (2%) 58 58	22, 41, 69, 96	0
2	E	294/313 (93%)	-0.37	5 (1%) 73 74	13, 26, 64, 105	0
3	C	299/311 (96%)	-0.03	11 (3%) 45 45	25, 44, 70, 86	0
3	F	285/311 (91%)	-0.20	12 (4%) 40 39	16, 30, 66, 113	0
4	G	4/4 (100%)	0.70	0 100 100	63, 64, 65, 70	0
4	H	4/4 (100%)	0.61	0 100 100	62, 62, 65, 66	0
4	I	4/4 (100%)	0.02	0 100 100	36, 38, 40, 43	0
4	J	4/4 (100%)	0.01	0 100 100	34, 37, 38, 41	0
All	All	1316/1396 (94%)	-0.10	51 (3%) 43 43	13, 37, 80, 115	0

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	126	VAL	6.8
3	F	394	ILE	5.9
3	C	394	ILE	5.4
1	D	133	ILE	4.8
1	D	138	LYS	4.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	FUC	E	5	10/11	0.93	0.28	-	62,63,63,64	0
5	NAG	E	3	14/15	0.86	0.19	-	48,56,61,67	0
5	NAG	E	4	14/15	0.74	0.41	-	69,72,77,78	0
5	FUC	B	5	10/11	0.75	0.40	-	101,103,103,103	0
5	NAG	B	4	14/15	0.73	0.55	-	105,106,108,109	0
5	NAG	B	3	14/15	0.76	0.32	-	91,95,100,102	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	CA	C	1	1/1	0.98	0.04	-1.81	44,44,44,44	0
6	CA	F	1	1/1	0.96	0.05	-2.51	29,29,29,29	0
6	CA	B	2	1/1	0.94	0.05	-	54,54,54,54	0
6	CA	E	2	1/1	0.91	0.10	-	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.