



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:25 PM GMT

PDB ID : 1OZR
Title : Crystal Structures of the Ferric, Ferrous and Ferrous-NO Forms of the Asp140Ala Mutant of Human Heme Oxygenase-1: Catalytic Implications
Authors : Lad, L.; Wang, J.; Li, H.; Friedman, J.; Ortiz de Montellano, P.R.; Poulos, T.L.
Deposited on : 2003-04-09
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2570 (1.76-1.72)
Ramachandran outliers	100387	2544 (1.76-1.72)
Sidechain outliers	100360	2544 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	233	 72% 20% 8%
1	B	233	 80% 12% 8%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heme oxygenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	28	0	0
			1745	1121	299	320	5			
1	B	214	Total	C	N	O	S	20	0	0
			1745	1121	299	320	5			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	1	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	1	0
			43	34	1	4	4		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	187	Total 187	O 187	0	0
3	B	163	Total 163	O 163	0	0

Note EDS was not executed.

- Chain A: 72% 20% 8%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.85Å 54.65Å 72.14Å 90.00° 99.28° 90.00°	Depositor
Resolution (Å)	50.00 – 1.74	Depositor
% Data completeness (in resolution range)	99.8 (50.00-1.74)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.06	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.209 , 0.232	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3926	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1786	0.53	0/2414
1	B	0.33	0/1786	0.54	0/2414
All	All	0.33	0/3572	0.54	0/4828

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1738	37	0
1	B	1745	0	1738	18	0
2	A	43	0	30	1	0
2	B	43	0	30	1	0
3	A	187	0	0	9	0
3	B	163	0	0	4	0
All	All	3926	0	3536	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195:VAL:HG22	1:A:198:ARG:HH22	1.27	0.99
1:A:142:SER:O	1:A:145:GLN:HG2	1.82	0.80
1:A:142:SER:HB3	1:A:179:LYS:NZ	2.06	0.70
1:A:41:GLN:HG3	3:A:462:HOH:O	1.90	0.69
1:B:44:ARG:O	1:B:48:LYS:HG3	1.93	0.69
1:A:149:LYS:HA	1:A:152:GLN:HE21	1.57	0.69
1:A:195:VAL:HG22	1:A:198:ARG:NH2	2.06	0.66
1:A:91:GLN:HE22	1:B:85:ARG:HH22	1.44	0.65
1:B:84:HIS:HD2	3:B:384:HOH:O	1.79	0.65
1:A:179:LYS:O	1:A:183:ARG:HG3	1.96	0.64
1:A:102:GLN:HG3	3:A:420:HOH:O	1.96	0.64
1:A:142:SER:HB3	1:A:179:LYS:HZ3	1.64	0.60
1:A:125:GLU:N	1:A:126:PRO:HD3	2.16	0.60
1:B:113:ARG:HD2	3:B:463:HOH:O	2.01	0.60
1:A:85:ARG:HH22	1:B:91:GLN:HE22	1.49	0.59
1:A:143:GLY:O	1:A:146:VAL:HG22	2.03	0.58
1:A:84:HIS:HD2	3:A:403:HOH:O	1.86	0.58
1:A:192:THR:OG1	1:A:195:VAL:HG23	2.05	0.57
1:A:176:THR:O	1:A:180:GLN:HG3	2.04	0.56
1:A:71:SER:OG	1:A:73:VAL:HG22	2.05	0.56
1:B:75:ALA:HB3	1:B:76:PRO:HD3	1.88	0.56
1:A:10:PRO:HD2	1:A:16:ALA:HA	1.88	0.54
1:A:56:HIS:HD2	3:A:395:HOH:O	1.91	0.53
1:A:10:PRO:HG3	1:A:15:GLU:HG2	1.90	0.53
1:A:116:LYS:O	1:A:120:GLU:HG3	2.09	0.52
1:A:157:LEU:HB2	1:A:158:PRO:HD2	1.90	0.52
1:B:144:GLY:HA3	3:B:335:HOH:O	2.09	0.51
1:A:23:GLU:HB3	3:A:340:HOH:O	2.10	0.51
1:B:109:PRO:O	1:B:113:ARG:HG3	2.11	0.51
1:B:211:ILE:O	1:B:215:GLU:HG3	2.11	0.50
1:B:56:HIS:HD2	3:B:406:HOH:O	1.94	0.50
1:B:125:GLU:N	1:B:126:PRO:HD3	2.27	0.49
1:A:176:THR:HG23	3:A:441:HOH:O	2.13	0.49
1:A:29:GLU:HA	2:A:300:HEM:HBB2	1.94	0.49
1:A:33:PHE:CD2	1:A:34:MET:HE2	2.48	0.49
1:A:212:GLN:NE2	3:A:484:HOH:O	2.45	0.49
1:A:211:ILE:O	1:A:215:GLU:HG3	2.12	0.48
1:A:33:PHE:HD2	1:A:34:MET:HE2	1.79	0.48
1:B:17:LEU:HD23	1:B:200:ILE:HD13	1.96	0.47
1:B:86:LYS:O	1:B:90:GLU:HG3	2.15	0.47
1:A:212:GLN:HG3	3:A:316:HOH:O	2.13	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:THR:C	1:A:126:PRO:HD3	2.35	0.46
1:B:122:GLY:O	1:B:126:PRO:HG3	2.14	0.46
1:A:134:TYR:CD1	1:A:138:LEU:HD22	2.52	0.44
1:A:142:SER:HB3	1:A:179:LYS:HZ2	1.78	0.43
1:B:56:HIS:HE1	1:B:106:PRO:O	2.01	0.43
1:A:196:ARG:O	1:A:200:ILE:HG12	2.18	0.43
3:A:304:HOH:O	1:B:84:HIS:HE1	2.01	0.42
1:A:75:ALA:HB3	1:A:76:PRO:HD3	2.01	0.42
1:A:44:ARG:NH2	1:A:157:LEU:HA	2.34	0.41
1:B:96:TRP:CH2	1:B:157:LEU:HD21	2.56	0.41
1:A:209:LEU:O	1:A:212:GLN:HG2	2.21	0.40
1:B:134:TYR:OH	2:B:300:HEM:HBD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/233 (91%)	207 (98%)	5 (2%)	0	100	100
1	B	212/233 (91%)	209 (99%)	3 (1%)	0	100	100
All	All	424/466 (91%)	416 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/202 (91%)	182 (99%)	2 (1%)	80	66
1	B	184/202 (91%)	180 (98%)	4 (2%)	60	35
All	All	368/404 (91%)	362 (98%)	6 (2%)	70	51

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	TYR
1	A	157	LEU
1	B	23	GLU
1	B	37	PHE
1	B	114	TYR
1	B	162	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS
1	A	68	ASN
1	A	84	HIS
1	A	91	GLN
1	A	102	GLN
1	A	112	GLN
1	A	152	GLN
1	A	171	ASN
1	B	30	ASN
1	B	36	ASN
1	B	56	HIS
1	B	68	ASN
1	B	84	HIS
1	B	91	GLN
1	B	102	GLN
1	B	112	GLN
1	B	171	ASN
1	B	180	GLN
1	B	197	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	300	1	30,50,50	2.75	10 (33%)	24,82,82	1.94	6 (25%)
2	HEM	B	300	1	30,50,50	2.80	10 (33%)	24,82,82	2.01	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	300	1	-	0/10/54/54	0/0/8/8
2	HEM	B	300	1	-	0/10/54/54	0/0/8/8

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	HEM	C2D-C3D	-6.44	1.35	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	HEM	C3B-CAB	-6.43	1.39	1.51
2	B	300	HEM	C3B-CAB	-6.20	1.39	1.51
2	B	300	HEM	C2D-C3D	-6.16	1.36	1.54
2	B	300	HEM	C3C-CAC	-6.09	1.39	1.51
2	A	300	HEM	C3C-CAC	-5.98	1.40	1.51
2	B	300	HEM	C3D-C4D	-5.42	1.44	1.51
2	B	300	HEM	C3B-C4B	-5.19	1.47	1.51
2	A	300	HEM	C3B-C4B	-5.12	1.47	1.51
2	A	300	HEM	C3D-C4D	-4.51	1.45	1.51
2	B	300	HEM	C2C-C1C	-3.67	1.45	1.52
2	A	300	HEM	C2C-C1C	-3.01	1.46	1.52
2	A	300	HEM	CBC-CAC	2.68	1.44	1.29
2	A	300	HEM	CBB-CAB	2.75	1.45	1.29
2	B	300	HEM	CBB-CAB	2.83	1.45	1.29
2	B	300	HEM	CBC-CAC	2.94	1.46	1.29
2	B	300	HEM	C1C-NC	3.17	1.39	1.36
2	A	300	HEM	C4C-NC	3.30	1.40	1.36
2	B	300	HEM	C4C-NC	3.30	1.40	1.36
2	A	300	HEM	C1C-NC	3.45	1.40	1.36

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	HEM	CMD-C2D-C3D	2.42	125.03	114.35
2	B	300	HEM	CMD-C2D-C3D	2.45	125.17	114.35
2	B	300	HEM	C2D-C3D-C4D	3.18	106.89	101.50
2	A	300	HEM	C2D-C3D-C4D	3.29	107.07	101.50
2	A	300	HEM	CAD-C3D-C2D	3.32	122.77	113.22
2	B	300	HEM	CAD-C3D-C4D	3.46	124.68	112.47
2	A	300	HEM	CMC-C2C-C3C	3.49	125.23	116.53
2	B	300	HEM	CMB-C2B-C3B	3.75	125.88	116.53
2	A	300	HEM	CMB-C2B-C3B	3.95	126.38	116.53
2	B	300	HEM	CMC-C2C-C3C	4.00	126.52	116.53
2	A	300	HEM	CAD-C3D-C4D	5.01	130.15	112.47
2	B	300	HEM	CAD-C3D-C2D	5.29	128.41	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	HEM	1	0
2	B	300	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.