



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:59 AM GMT

PDB ID : 2OZN  
Title : The Cohesin-Dockerin Complex of NagJ and NagH from *Clostridium perfringens*  
Authors : Adams, J.J.; Boraston, A.; Smith, S.P.  
Deposited on : 2007-02-26  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	<div> <div>3%</div> <div>73%</div> <div>7%</div> <div>19%</div> </div>
2	B	140	<div> <div>10%</div> <div>86%</div> <div>8%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	403	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-GlcNAcase nagJ.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	0	0	0
			991	620	157	214			

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	745	MET	-	INITIATING METHIONINE	UNP Q0TR53
A	746	GLY	-	EXPRESSION TAG	UNP Q0TR53
A	747	SER	-	EXPRESSION TAG	UNP Q0TR53
A	748	SER	-	EXPRESSION TAG	UNP Q0TR53
A	749	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	750	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	751	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	752	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	753	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	754	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	755	SER	-	EXPRESSION TAG	UNP Q0TR53
A	756	SER	-	EXPRESSION TAG	UNP Q0TR53
A	757	GLY	-	EXPRESSION TAG	UNP Q0TR53
A	758	LEU	-	EXPRESSION TAG	UNP Q0TR53
A	759	VAL	-	EXPRESSION TAG	UNP Q0TR53
A	760	PRO	-	EXPRESSION TAG	UNP Q0TR53
A	761	ARG	-	EXPRESSION TAG	UNP Q0TR53
A	762	GLY	-	EXPRESSION TAG	UNP Q0TR53
A	763	SER	-	EXPRESSION TAG	UNP Q0TR53
A	764	HIS	-	EXPRESSION TAG	UNP Q0TR53
A	765	MET	-	EXPRESSION TAG	UNP Q0TR53
A	766	ALA	-	EXPRESSION TAG	UNP Q0TR53
A	767	SER	-	EXPRESSION TAG	UNP Q0TR53

- Molecule 2 is a protein called Hyaluronoglucosaminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	131	Total	C	N	O	S	0	0	0
			1004	618	166	219	1			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1497	MET	-	EXPRESSION TAG	UNP P26831
B	1629	LEU	-	EXPRESSION TAG	UNP P26831
B	1630	GLU	-	EXPRESSION TAG	UNP P26831
B	1631	HIS	-	EXPRESSION TAG	UNP P26831
B	1632	HIS	-	EXPRESSION TAG	UNP P26831
B	1633	HIS	-	EXPRESSION TAG	UNP P26831
B	1634	HIS	-	EXPRESSION TAG	UNP P26831
B	1635	HIS	-	EXPRESSION TAG	UNP P26831
B	1636	HIS	-	EXPRESSION TAG	UNP P26831

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ca	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	181	Total	O	0	0
			181	181		
5	B	134	Total	O	0	0
			134	134		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: O-GlcNAcase nagJ



- Molecule 2: Hyalurononglucosaminidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.49Å 74.59Å 94.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	13.88 – 1.60 13.88 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.8 (13.88-1.60) 98.8 (13.88-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.83 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.204 , 0.246 0.202 , 0.242	Depositor DCC
$R_{free}$ test set	1689 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.4	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 61.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 33593 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2313	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/1000	0.69	0/1357
2	B	0.46	0/1012	0.60	0/1360
All	All	0.51	0/2012	0.64	0/2717

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	991	0	970	13	0
2	B	1004	0	967	7	0
3	A	1	0	0	0	0
4	B	2	0	0	0	0
5	A	181	0	0	2	0
5	B	134	0	0	5	0
All	All	2313	0	1937	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1531:ASP:HB2	5:B:1695:HOH:O	1.65	0.95
2:B:1604:THR:C	5:B:1720:HOH:O	2.18	0.82
2:B:1624:HIS:HD2	5:B:1722:HOH:O	1.65	0.79
2:B:1625:ARG:HG2	5:B:1739:HOH:O	1.87	0.73
1:A:828:SER:HB2	1:A:832:VAL:CG2	2.26	0.65
1:A:787:VAL:CG2	1:A:791:GLU:HB3	2.29	0.63
1:A:828:SER:HB2	1:A:832:VAL:HG23	1.84	0.59
1:A:787:VAL:HG22	1:A:791:GLU:HB3	1.84	0.59
1:A:785:GLU:HG2	5:A:951:HOH:O	2.04	0.56
2:B:1517:VAL:HG13	2:B:1524:GLU:HG2	1.91	0.52
1:A:828:SER:HB3	1:A:834:VAL:HG21	1.91	0.52
1:A:787:VAL:HG23	1:A:791:GLU:CD	2.32	0.51
1:A:828:SER:HB3	1:A:834:VAL:CG2	2.46	0.45
2:B:1616:GLU:HB2	5:B:1668:HOH:O	2.18	0.43
1:A:837:LYS:HZ2	2:B:1628:ASN:HD21	1.68	0.42
1:A:774:GLU:HG2	1:A:774:GLU:O	2.20	0.42
1:A:784:LEU:HD12	5:A:1090:HOH:O	2.18	0.41
1:A:787:VAL:HG23	1:A:791:GLU:OE1	2.20	0.41
1:A:828:SER:CB	1:A:832:VAL:HG23	2.49	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/165 (79%)	130 (99%)	1 (1%)	0	100	100
2	B	129/140 (92%)	124 (96%)	5 (4%)	0	100	100
All	All	260/305 (85%)	254 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/134 (81%)	105 (97%)	3 (3%)	51	22
2	B	111/120 (92%)	108 (97%)	3 (3%)	52	23
All	All	219/254 (86%)	213 (97%)	6 (3%)	52	23

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	835	ASN
1	A	872	LYS
1	A	906	GLU
2	B	1596	LYS
2	B	1606	LEU
2	B	1609	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	877	ASN
2	B	1624	HIS
2	B	1628	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/165 (80%)	-0.00	5 (3%) 44 41	12, 18, 27, 37	0
2	B	131/140 (93%)	0.57	14 (10%) 8 7	15, 22, 37, 47	0
All	All	264/305 (86%)	0.28	19 (7%) 18 16	12, 20, 34, 47	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1546	LYS	8.9
2	B	1582	GLY	6.9
2	B	1604	THR	6.3
2	B	1547	GLU	5.5
2	B	1581	ASN	5.0
2	B	1548	ASP	4.5
1	A	906	GLU	4.4
1	A	830	ASP	3.6
2	B	1584	GLY	3.1
1	A	774	GLU	3.0
2	B	1498	ASP	2.8
2	B	1583	ASN	2.7
2	B	1556	LEU	2.7
2	B	1549	ALA	2.7
1	A	872	LYS	2.4
2	B	1550	THR	2.2
1	A	904	ILE	2.1
2	B	1559	GLU	2.1
2	B	1552	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	403	1/1	1.00	0.14	2.77	26,26,26,26	0
4	CA	B	402	1/1	0.97	0.07	-1.20	27,27,27,27	0
4	CA	B	401	1/1	0.99	0.06	-1.68	20,20,20,20	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.