



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:26 PM GMT

PDB ID : 1P06
Title : STRUCTURE ANALYSIS OF SPECIFICITY. ALPHA-LYTIC PROTEASE
COMPLEXES WITH ANALOGUES OF REACTION INTERMEDIATES
Authors : Bone, R.; Agard, D.A.
Deposited on : 1989-04-24
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1509 (2.36-2.32)
Ramachandran outliers	100387	1490 (2.36-2.32)
Sidechain outliers	100360	1491 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	198	
2	P	5	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALPHA-LYTIC PROTEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	1391	846	262	275	8	0	0	0

- Molecule 2 is a protein called METHOXYSUCCINYL-ALA-ALA-PRO-PHENYLALANINE BORONIC ACID INHIBITOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	B	C	N	O			
2	P	4	28	1	19	4	4	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	161	Total 161	O 161	0	0
4	P	2	Total 2	O 2	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ALPHA-LYTIC PROTEASE



- Molecule 2: METHOXYsuccinyl-ALA-ALA-PRO-PHENYLALANINE BORONIC ACID INHIBITOR



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	66.62Å 66.62Å 80.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.34	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.34)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.140 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1587	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: B2F, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.28	0/1409	2.92	83/1909 (4.3%)
2	P	1.12	0/17	2.45	1/23 (4.3%)
All	All	1.28	0/1426	2.92	84/1932 (4.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (84) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	125	ARG	NE-CZ-NH2	-38.44	101.08	120.30
1	A	125	ARG	NE-CZ-NH1	26.70	133.65	120.30
1	A	138	ARG	NE-CZ-NH1	-26.69	106.95	120.30
1	A	125	ARG	CD-NE-CZ	21.84	154.18	123.60
1	A	117	ARG	NE-CZ-NH2	-21.08	109.76	120.30
1	A	48(B)	ARG	NE-CZ-NH1	-20.68	109.96	120.30
1	A	138	ARG	NE-CZ-NH2	18.06	129.33	120.30
1	A	91	ARG	NE-CZ-NH2	16.47	128.53	120.30
1	A	67	ARG	NE-CZ-NH1	16.04	128.32	120.30
1	A	178	ARG	NE-CZ-NH1	15.67	128.14	120.30
1	A	91	ARG	CD-NE-CZ	11.94	140.32	123.60
1	A	91	ARG	NH1-CZ-NH2	-11.88	106.33	119.40
1	A	141	ARG	NE-CZ-NH1	-11.36	114.62	120.30
1	A	230	ARG	NE-CZ-NH1	10.54	125.57	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	ASN	CB-CG-OD1	-10.06	101.48	121.60
1	A	161	THR	CA-CB-CG2	9.96	126.34	112.40
1	A	117	ARG	NE-CZ-NH1	9.73	125.17	120.30
1	A	120(D)	SER	N-CA-CB	-9.58	96.13	110.50
1	A	91	ARG	NE-CZ-NH1	9.31	124.96	120.30
1	A	178	ARG	NH1-CZ-NH2	-9.27	109.20	119.40
1	A	192(B)	ARG	CD-NE-CZ	-8.95	111.07	123.60
1	A	120(D)	SER	CB-CA-C	8.86	126.94	110.10
1	A	129	GLU	OE1-CD-OE2	8.63	133.66	123.30
1	A	141	ARG	NE-CZ-NH2	8.62	124.61	120.30
1	A	143	THR	CA-CB-CG2	8.57	124.40	112.40
1	A	192(B)	ARG	NE-CZ-NH2	-8.46	116.07	120.30
1	A	201	THR	CA-CB-CG2	8.30	124.02	112.40
1	A	64	ASN	CB-CG-ND2	8.24	136.48	116.70
1	A	125	ARG	N-CA-CB	-7.86	96.45	110.60
1	A	67	ARG	NE-CZ-NH2	-7.78	116.41	120.30
1	A	49	THR	CA-CB-OG1	-7.76	92.70	109.00
1	A	48(B)	ARG	CD-NE-CZ	7.50	134.11	123.60
1	A	199	TRP	N-CA-CB	7.25	123.66	110.60
1	A	109	THR	CA-CB-CG2	-7.19	102.34	112.40
1	A	117	ARG	O-C-N	7.15	134.15	122.70
2	P	3	ALA	N-CA-CB	7.01	119.91	110.10
1	A	195	SER	CB-CA-C	-6.97	96.87	110.10
1	A	122	VAL	CA-CB-CG2	6.95	121.32	110.90
1	A	174	GLU	OE1-CD-OE2	6.82	131.48	123.30
1	A	17	VAL	CG1-CB-CG2	-6.59	100.36	110.90
1	A	48(B)	ARG	NE-CZ-NH2	6.57	123.59	120.30
1	A	90	ALA	CB-CA-C	-6.57	100.25	110.10
1	A	67	ARG	CD-NE-CZ	-6.50	114.50	123.60
1	A	103	ARG	NE-CZ-NH2	6.44	123.52	120.30
1	A	41	LEU	O-C-N	6.42	132.96	122.70
1	A	156	GLY	C-N-CA	6.41	137.72	121.70
1	A	190	ALA	CA-C-N	6.36	131.20	117.20
1	A	48(B)	ARG	NH1-CZ-NH2	6.29	126.31	119.40
1	A	85	VAL	CA-CB-CG2	6.28	120.31	110.90
1	A	106	VAL	O-C-N	6.27	132.73	122.70
1	A	117	ARG	CD-NE-CZ	-6.26	114.84	123.60
1	A	209	ALA	N-CA-CB	-6.25	101.34	110.10
1	A	105	TRP	CB-CA-C	6.08	122.56	110.40
1	A	139	SER	N-CA-CB	-6.07	101.39	110.50
1	A	114	LEU	N-CA-CB	-6.00	98.40	110.40
1	A	63	VAL	O-C-N	5.99	132.28	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	ALA	N-CA-CB	-5.87	101.88	110.10
1	A	224	ARG	CB-CG-CD	-5.84	96.41	111.60
1	A	48(B)	ARG	N-CA-CB	-5.83	100.11	110.60
1	A	31	TYR	CB-CA-C	5.76	121.93	110.40
1	A	164	ALA	N-CA-CB	-5.71	102.10	110.10
1	A	48(B)	ARG	CA-CB-CG	-5.71	100.84	113.40
1	A	143	THR	CA-C-N	5.64	127.47	116.20
1	A	102	ASP	CB-CG-OD1	-5.62	113.24	118.30
1	A	216	GLY	O-C-N	5.43	131.38	122.70
1	A	203	ALA	CA-C-O	-5.42	108.71	120.10
1	A	105	TRP	N-CA-CB	5.42	120.35	110.60
1	A	217(E)	GLY	CA-C-O	-5.39	110.90	120.60
1	A	237	TYR	CB-CG-CD1	-5.37	117.78	121.00
1	A	34	ASN	N-CA-CB	-5.35	100.98	110.60
1	A	143	THR	CA-CB-OG1	-5.33	97.81	109.00
1	A	242	VAL	CA-CB-CG1	-5.25	103.03	110.90
1	A	165	LYS	N-CA-CB	-5.25	101.16	110.60
1	A	90	ALA	N-CA-CB	-5.24	102.76	110.10
1	A	184	ASN	CA-C-O	-5.22	109.13	120.10
1	A	117	ARG	NH1-CZ-NH2	5.16	125.07	119.40
1	A	125	ARG	NH1-CZ-NH2	5.15	125.06	119.40
1	A	48(C)	GLY	N-CA-C	-5.10	100.34	113.10
1	A	113	THR	O-C-N	5.07	130.81	122.70
1	A	228	PHE	CA-CB-CG	5.07	126.06	113.90
1	A	244	GLY	CA-C-O	-5.07	111.48	120.60
1	A	237	TYR	CB-CG-CD2	5.05	124.03	121.00
1	A	163	THR	O-C-N	5.02	130.73	122.70
1	A	88	PHE	CB-CG-CD2	-5.01	117.30	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	192(B)	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1391	0	1359	17	0
2	P	28	0	25	2	0
3	A	5	0	0	0	0
4	A	161	0	0	2	0
4	P	2	0	0	0	0
All	All	1587	0	1384	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ILE:HD12	1:A:85:VAL:HG21	1.71	0.73
1:A:221(C):ALA:HA	1:A:224:ARG:HD2	1.90	0.52
1:A:94:PHE:CE1	1:A:174:GLU:HG3	2.44	0.52
1:A:171:TYR:CE1	2:P:2:PRO:HB3	2.48	0.49
1:A:91:ARG:HB3	1:A:104:ALA:HB2	1.95	0.48
1:A:48(B):ARG:HD2	4:A:355:HOH:O	2.15	0.47
1:A:171:TYR:CD1	2:P:2:PRO:HB3	2.51	0.46
1:A:94:PHE:HE1	1:A:174:GLU:HG3	1.81	0.46
1:A:30:GLU:HG2	1:A:31:TYR:N	2.29	0.46
1:A:191:CYS:HA	1:A:219:ASN:OD1	2.16	0.46
1:A:65:ALA:O	1:A:87:THR:HA	2.15	0.46
1:A:232:GLN:N	1:A:233:PRO:HD2	2.32	0.45
1:A:48(B):ARG:HG3	1:A:239:LEU:HD23	1.98	0.45
1:A:137:CYS:HA	1:A:159:CYS:HA	1.99	0.44
1:A:46:PHE:O	1:A:52:PHE:HA	2.21	0.41
1:A:129:GLU:HG3	4:A:364:HOH:O	2.21	0.40
1:A:138:ARG:O	1:A:157:TYR:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/198 (99%)	188 (96%)	8 (4%)	0	100	100
2	P	2/5 (40%)	2 (100%)	0	0	100	100
All	All	198/203 (98%)	190 (96%)	8 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/142 (100%)	136 (96%)	6 (4%)	36	46
2	P	1/1 (100%)	1 (100%)	0	100	100
All	All	143/143 (100%)	137 (96%)	6 (4%)	36	46

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48(B)	ARG
1	A	91	ARG
1	A	110	SER
1	A	141	ARG
1	A	166	ASN
1	A	192	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B2F	P	1	1,2	9,11,12	0.97	1 (11%)	9,13,15	1.10	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B2F	P	1	1,2	-	0/3/6/8	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	1	B2F	CA-N	2.03	1.51	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	1	B2F	CG-CB-CA	-2.26	108.81	113.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	1	-	4,4,4	1.05	0	6,6,6	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	1	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.