



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 05:00 AM GMT

PDB ID : 2P0V
Title : Crystal structure of BT3781 protein from Bacteroides thetaiotaomicron, Northeast Structural Genomics Target BtR58
Authors : Forouhar, F.; Chen, Y.; Seetharaman, J.; Janjua, H.; Xiao, R.; Liu, J.; Baran, M.C.; Acton, T.B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2007-03-01
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	489	
1	B	489	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein BT3781.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	443	3566	2283	599	663	6	15	0	0	0
1	B	443	3566	2283	599	663	6	15	0	0	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	22	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	75	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	106	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	131	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	154	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	162	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	168	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	243	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	344	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	401	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	405	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	408	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	409	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	426	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	428	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	437	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
A	482	LEU	-	CLONING ARTIFACT	UNP Q8A185
A	483	GLU	-	CLONING ARTIFACT	UNP Q8A185
A	484	HIS	-	CLONING ARTIFACT	UNP Q8A185
A	485	HIS	-	CLONING ARTIFACT	UNP Q8A185
A	486	HIS	-	CLONING ARTIFACT	UNP Q8A185
A	487	HIS	-	CLONING ARTIFACT	UNP Q8A185
A	488	HIS	-	CLONING ARTIFACT	UNP Q8A185
A	489	HIS	-	CLONING ARTIFACT	UNP Q8A185

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	22	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	75	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	106	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	131	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	154	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	162	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	168	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	243	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	344	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	401	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	405	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	408	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	409	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	426	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	428	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	437	MSE	MET	MODIFIED RESIDUE	UNP Q8A185
B	482	LEU	-	CLONING ARTIFACT	UNP Q8A185
B	483	GLU	-	CLONING ARTIFACT	UNP Q8A185
B	484	HIS	-	CLONING ARTIFACT	UNP Q8A185
B	485	HIS	-	CLONING ARTIFACT	UNP Q8A185
B	486	HIS	-	CLONING ARTIFACT	UNP Q8A185
B	487	HIS	-	CLONING ARTIFACT	UNP Q8A185
B	488	HIS	-	CLONING ARTIFACT	UNP Q8A185
B	489	HIS	-	CLONING ARTIFACT	UNP Q8A185


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	240	Total O 240 240	0	0
2	B	228	Total O 228 228	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical protein BT3781



- Molecule 1: Hypothetical protein BT3781



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.07Å 64.01Å 74.19Å 80.74° 84.14° 75.05°	Depositor
Resolution (Å)	29.61 – 2.10 29.60 – 2.07	Depositor EDS
% Data completeness (in resolution range)	79.9 (29.61-2.10) 89.6 (29.60-2.07)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.08Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.211 , 0.234 0.228 , 0.251	Depositor DCC
R_{free} test set	4661 reflections (9.50%)	DCC
Wilson B-factor (Å ²)	16.5	Xtrriage
Anisotropy	0.386	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 99728 reflections	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7600	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3644	0.67	6/4913 (0.1%)
1	B	0.35	0/3644	0.70	6/4913 (0.1%)
All	All	0.35	0/7288	0.68	12/9826 (0.1%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	187	ARG	NE-CZ-NH1	15.42	128.01	120.30
1	B	187	ARG	NE-CZ-NH2	-14.96	112.82	120.30
1	B	58	ARG	NE-CZ-NH2	-13.23	113.68	120.30
1	B	58	ARG	NE-CZ-NH1	13.06	126.83	120.30
1	A	58	ARG	NE-CZ-NH2	12.93	126.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3566	0	3474	79	0
1	B	3566	0	3474	85	0
2	A	240	0	0	5	0
2	B	228	0	0	6	0
All	All	7600	0	6948	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:ARG:H	1:A:242:THR:HG23	1.30	0.97
1:B:234:ARG:H	1:B:242:THR:HG23	1.31	0.96
1:A:402:ILE:HD11	1:A:407:ILE:HD11	1.53	0.91
1:B:402:ILE:HD11	1:B:407:ILE:HD11	1.54	0.88
1:B:389:GLY:HA3	1:B:426:MSE:HE3	1.64	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/489 (90%)	417 (95%)	22 (5%)	2 (0%)	34	30
1	B	441/489 (90%)	418 (95%)	20 (4%)	3 (1%)	26	21
All	All	882/978 (90%)	835 (95%)	42 (5%)	5 (1%)	30	24

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PRO
1	B	44	PRO
1	B	451	ARG
1	B	254	PRO
1	A	254	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/404 (95%)	370 (97%)	12 (3%)	47	50
1	B	382/404 (95%)	369 (97%)	13 (3%)	44	45
All	All	764/808 (95%)	739 (97%)	25 (3%)	45	47

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	454	PHE
1	B	110	ASP
1	B	454	PHE
1	B	107	TRP
1	B	246	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	42	ASN
1	B	104	HIS
1	B	442	ASN
1	B	62	HIS
1	B	114	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	428/489 (87%)	0.20	14 (3%) 50 59	10, 21, 35, 48	0
1	B	428/489 (87%)	0.23	14 (3%) 50 59	12, 23, 36, 55	0
All	All	856/978 (87%)	0.21	28 (3%) 50 59	10, 22, 36, 55	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	39	TYR	4.3
1	B	481	GLN	3.6
1	A	297	ARG	3.5
1	A	63	ILE	3.2
1	A	481	GLN	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.