



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:28 PM GMT

PDB ID : 4P0Q
Title : Crystal structure of Human Mus81-Eme1 in complex with 5'-flap DNA
Authors : Gwon, G.H.; Baek, K.; Cho, Y.
Deposited on : 2014-02-22
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2228 (2.90-2.82)
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)
RSRZ outliers	91569	2236 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>16%</div> <div> <div>75%</div> <div>16%</div> <div>•</div> <div>7%</div> </div> </div>
2	B	393	<div> <div>15%</div> <div> <div>48%</div> <div>21%</div> <div>•</div> <div>28%</div> </div> </div>
3	E	17	<div> <div>65%</div> <div> <div>35%</div> <div>65%</div> </div> </div>
4	F	6	<div> <div>83%</div> <div> <div>33%</div> <div>67%</div> </div> </div>
5	G	16	<div> <div>31%</div> <div> <div>38%</div> <div>31%</div> <div>31%</div> </div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Crossover junction endonuclease MUS81.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2255	1414	422	411	8			

- Molecule 2 is a protein called Crossover junction endonuclease EME1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	284	Total	C	N	O	S	0	0	0
			2227	1401	397	415	14			

- Molecule 3 is a DNA chain called DNA GAATGTGTGTCTCAATC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	17	Total	C	N	O	P	0	0	0
			349	167	61	104	17			

- Molecule 4 is a DNA chain called DNA GGATTG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	6	Total	C	N	O	P	0	0	0
			127	60	24	37	6			

- Molecule 5 is a DNA chain called DNA TAACCAGACACACATT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	G	11	Total	C	N	O	P	0	0	0
			224	107	43	63	11			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	18	Total	O	0	0
			18	18		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	11	Total 11	O 11	0	0
6	E	1	Total 1	O 1	0	0
6	F	2	Total 2	O 2	0	0
6	G	1	Total 1	O 1	0	0



● Molecule 4: DNA GGATTG



● Molecule 5: DNA TAACCAGACACACATT



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.41Å 228.38Å 52.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.45 – 2.85 47.80 – 2.85	Depositor EDS
% Data completeness (in resolution range)	98.1 (42.45-2.85) 98.4 (47.80-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 2.86Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.243 , 0.276 0.244 , 0.274	Depositor DCC
R_{free} test set	1269 reflections (5.13%)	DCC
Wilson B-factor (Å ²)	95.6	Xtriage
Anisotropy	0.447	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 67.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 24814 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5215	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2295	0.41	0/3103
2	B	0.23	0/2258	0.42	0/3054
3	E	0.49	0/390	0.93	1/600 (0.2%)
4	F	0.43	0/142	0.79	0/218
5	G	0.40	0/251	0.94	0/384
All	All	0.27	0/5336	0.53	1/7359 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	E	1	DG	C3'-C2'-C1'	-5.72	95.63	102.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2255	0	2293	31	0
2	B	2227	0	2265	50	0
3	E	349	0	194	18	0
4	F	127	0	69	8	0
5	G	224	0	124	5	0
6	A	18	0	0	4	0
6	B	11	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	E	1	0	0	2	0
6	F	2	0	0	0	0
6	G	1	0	0	0	0
All	All	5215	0	4945	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:12:DT:H71	6:E:101:HOH:O	1.56	1.04
4:F:24:DT:H2''	4:F:25:DT:H5''	1.54	0.90
4:F:25:DT:H2''	4:F:26:DG:O5'	1.75	0.87
4:F:23:DA:H4'	4:F:24:DT:OP1	1.80	0.80
2:B:256:LEU:O	6:B:610:HOH:O	2.02	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/306 (91%)	267 (96%)	11 (4%)	1 (0%)	39	71
2	B	276/393 (70%)	267 (97%)	9 (3%)	0	100	100
All	All	555/699 (79%)	534 (96%)	20 (4%)	1 (0%)	52	82

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	259	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246 / 259 (95%)	227 (92%)	19 (8%)	16	39
2	B	242 / 334 (72%)	214 (88%)	28 (12%)	7	18
All	All	488 / 593 (82%)	441 (90%)	47 (10%)	10	28

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	269	LEU
2	B	291	VAL
2	B	531	GLN
2	B	282	ILE
2	B	301	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/306 (93%)	1.18	49 (17%) 2 1	31, 61, 114, 138	0
2	B	284/393 (72%)	1.31	59 (20%) 1 1	33, 65, 127, 148	0
3	E	17/17 (100%)	3.32	11 (64%) 0 0	64, 123, 170, 171	0
4	F	6/6 (100%)	4.24	5 (83%) 0 0	147, 156, 175, 177	0
5	G	11/16 (68%)	2.15	5 (45%) 0 0	109, 120, 144, 151	0
All	All	603/738 (81%)	1.35	129 (21%) 1 1	31, 64, 131, 177	0

The worst 5 of 129 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	542	SER	11.4
1	A	277	GLU	10.9
2	B	369	CYS	9.5
1	A	372	GLY	9.0
4	F	23	DA	8.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.