



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:29 PM GMT

PDB ID : 1P72  
Title : Crystal structure of EHV4-TK complexed with Thy and ADP  
Authors : Gardberg, A.; Shuvalova, L.; Monnerjahn, C.; Konrad, M.; Lavie, A.  
Deposited on : 2003-04-30  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>5%</div> <div>82%</div> <div>13%</div> <div>••</div> </div>
1	B	334	<div> <div>9%</div> <div>86%</div> <div>11%</div> <div>••</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5585 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	2	0
			2502	1583	439	461	19			
1	B	330	Total	C	N	O	S	0	1	0
			2507	1587	432	469	19			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	GLY	-	CLONING ARTIFACT	UNP P24425
A	20	SER	-	CLONING ARTIFACT	UNP P24425
A	21	HIS	-	CLONING ARTIFACT	UNP P24425
A	22	MET	-	CLONING ARTIFACT	UNP P24425
B	19	GLY	-	CLONING ARTIFACT	UNP P24425
B	20	SER	-	CLONING ARTIFACT	UNP P24425
B	21	HIS	-	CLONING ARTIFACT	UNP P24425
B	22	MET	-	CLONING ARTIFACT	UNP P24425

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is THYMIDINE (three-letter code: THM) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			17	10	2	5		
4	B	1	Total	C	N	O	0	0
			17	10	2	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	238	Total 238	O 238	0	0
5	B	205	Total 205	O 205	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Thymidine kinase



#### • Molecule 1: Thymidine kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.30Å 121.01Å 118.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.70 – 2.10 29.70 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.70-2.10) 96.8 (29.70-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.84 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.184 , 0.232 0.190 , 0.188	Depositor DCC
$R_{free}$ test set	4637 reflections (11.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.9	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 55.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 45956 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5585	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: THM, SO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/2564	0.78	6/3484 (0.2%)
1	B	0.57	0/2562	0.78	5/3488 (0.1%)
All	All	0.57	0/5126	0.78	11/6972 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	ASP	CB-CG-OD2	6.89	124.50	118.30
1	B	332	ASP	CB-CG-OD2	6.73	124.36	118.30
1	B	79	ASP	CB-CG-OD2	6.29	123.96	118.30
1	A	233	ASP	CB-CG-OD2	6.15	123.83	118.30
1	B	170	LEU	CA-CB-CG	5.76	128.56	115.30
1	A	79	ASP	CB-CG-OD2	5.71	123.44	118.30
1	A	253	ASP	CB-CG-OD2	5.39	123.16	118.30
1	B	307	ASP	CB-CG-OD2	5.31	123.08	118.30
1	A	114	ASP	CB-CG-OD2	5.25	123.03	118.30
1	B	170	LEU	CB-CG-CD2	5.08	119.64	111.00
1	A	132	ASP	CB-CG-OD2	5.05	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2502	0	2474	25	0
1	B	2507	0	2429	33	0
2	A	25	0	0	0	0
2	B	20	0	0	0	0
3	A	27	0	12	0	0
3	B	27	0	12	0	0
4	A	17	0	14	0	0
4	B	17	0	14	1	0
5	A	238	0	0	11	0
5	B	205	0	0	8	0
All	All	5585	0	4955	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:MET:HE2	5:B:971:HOH:O	1.69	0.92
1:B:91:VAL:O	1:B:93:ASP:N	2.07	0.85
1:B:259:VAL:HG23	1:B:260:SER:H	1.51	0.76
1:B:100:HIS:CE1	1:B:104:ARG:HE	2.03	0.75
1:A:296:LEU:HG	5:A:976:HOH:O	1.86	0.74
1:A:102:GLN:HG2	1:A:158:MET:HE1	1.71	0.72
1:A:252:MET:HG3	5:A:969:HOH:O	1.91	0.70
1:B:241:SER:O	1:B:273:THR:HG23	1.92	0.69
1:B:100:HIS:HE1	1:B:104:ARG:HH21	1.45	0.65
1:A:352:VAL:HA	5:A:1012:HOH:O	1.96	0.64
1:A:117:CYS:HB2	5:A:970:HOH:O	1.99	0.63
1:B:352:VAL:O	5:B:990:HOH:O	2.16	0.63
1:A:126:GLN:HG3	5:A:1028:HOH:O	2.02	0.58
1:B:100:HIS:HE1	1:B:104:ARG:NH2	2.02	0.57
1:A:118:THR:HG21	5:A:1020:HOH:O	2.05	0.56
1:B:100:HIS:CE1	1:B:104:ARG:NE	2.74	0.56
1:A:275:GLU:HG3	5:A:938:HOH:O	2.06	0.55
1:A:103:SER:HA	1:A:166[A]:MET:HE1	1.87	0.55
1:B:259:VAL:HG22	5:B:830:HOH:O	2.06	0.55
1:B:113:HIS:CE1	1:B:117[A]:CYS:SG	3.01	0.54
1:B:103:SER:OG	1:B:166:MET:HE1	2.08	0.53
1:A:285:GLU:OE2	5:A:997:HOH:O	2.19	0.52
1:A:241:SER:O	1:A:273:THR:HG23	2.11	0.51
1:A:191:ARG:O	1:A:195:THR:HG23	2.12	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:302:ASN:ND2	5:B:906:HOH:O	2.45	0.50
1:A:118:THR:CG2	5:A:1020:HOH:O	2.60	0.49
1:B:125:LEU:HD23	1:B:127:ARG:HG2	1.95	0.49
1:B:204:ASP:HA	5:B:980:HOH:O	2.13	0.47
1:B:153:TYR:CZ	1:B:246:LYS:HE3	2.49	0.47
1:A:100:HIS:CE1	1:B:99:ALA:HB1	2.50	0.47
1:B:246:LYS:O	1:B:250:THR:HG23	2.16	0.46
1:B:128:GLY:HA2	1:B:324:PRO:O	2.16	0.46
1:B:241:SER:O	1:B:273:THR:CG2	2.63	0.46
1:A:102:GLN:HG3	1:A:149:PRO:HG3	1.98	0.46
1:B:259:VAL:HG23	1:B:260:SER:N	2.27	0.45
1:A:197:ALA:O	1:A:198:ARG:O	2.34	0.45
1:A:229:ARG:NH2	1:A:261:PRO:O	2.48	0.45
1:A:252:MET:CG	5:A:969:HOH:O	2.57	0.44
1:B:98:THR:HG22	1:B:158:MET:HE2	1.99	0.44
1:B:246:LYS:NZ	5:B:987:HOH:O	2.50	0.43
1:B:344:ARG:HE	1:B:344:ARG:HB2	1.68	0.43
1:B:75:SER:O	1:B:79:ASP:HB2	2.18	0.43
1:B:166:MET:CE	5:B:971:HOH:O	2.42	0.43
1:A:164:MET:HB3	1:A:164:MET:HE3	1.83	0.43
1:B:43:ARG:HD3	5:B:990:HOH:O	2.18	0.42
1:A:36:ILE:HG22	1:A:308:LEU:HD12	2.01	0.42
1:B:59:PRO:HG3	1:B:352:VAL:HA	2.02	0.41
1:B:100:HIS:CE1	1:B:104:ARG:HH21	2.31	0.41
1:A:193:LEU:HD23	1:A:193:LEU:HA	1.94	0.41
1:A:55:THR:HA	1:A:134:THR:O	2.20	0.41
1:B:36:ILE:HG23	1:B:184:LEU:HB2	2.03	0.41
1:A:100:HIS:CE1	1:B:99:ALA:CB	3.04	0.41
1:B:242:CYS:HB2	1:B:243:GLY:H	1.63	0.41
1:A:39:SER:O	1:A:43[A]:ARG:HG3	2.21	0.41
1:A:299:ARG:NH1	5:A:951:HOH:O	2.53	0.41
1:B:259:VAL:CG2	1:B:260:SER:H	2.29	0.40
4:B:702:THM:H6	4:B:702:THM:O5'	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/334 (96%)	310 (97%)	9 (3%)	2 (1%)	30	24
1	B	327/334 (98%)	317 (97%)	7 (2%)	3 (1%)	21	15
All	All	648/668 (97%)	627 (97%)	16 (2%)	5 (1%)	24	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	ARG
1	B	92	ASP
1	B	90	ALA
1	B	91	VAL
1	A	199	ILE

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/279 (96%)	254 (95%)	14 (5%)	29	25
1	B	261/279 (94%)	250 (96%)	11 (4%)	36	35
All	All	529/558 (95%)	504 (95%)	25 (5%)	31	30

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	30	LEU
1	A	62	MET
1	A	75	SER
1	A	102	GLN
1	A	125	LEU
1	A	170	LEU
1	A	194	ARG
1	A	196	ARG
1	A	273	THR
1	A	280	ARG
1	A	299	ARG
1	A	302	ASN
1	A	344	ARG
1	B	50	SER
1	B	166	MET
1	B	170	LEU
1	B	196	ARG
1	B	237	GLU
1	B	242	CYS
1	B	248	ARG
1	B	273	THR
1	B	302	ASN
1	B	340	GLU
1	B	344	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	GLN
1	A	302	ASN
1	B	100	HIS
1	B	302	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	ADP	A	501	-	22,29,29	2.23	3 (13%)	27,45,45	2.80	5 (18%)
4	THM	A	701	-	13,18,18	0.51	0	16,26,26	2.04	2 (12%)
2	SO4	A	801	-	4,4,4	0.18	0	6,6,6	0.14	0
2	SO4	A	803	-	4,4,4	0.51	0	6,6,6	0.56	0
2	SO4	A	804	-	4,4,4	0.15	0	6,6,6	0.19	0
2	SO4	A	805	-	4,4,4	0.21	0	6,6,6	0.17	0
2	SO4	A	806	-	4,4,4	0.31	0	6,6,6	0.25	0
3	ADP	B	502	-	22,29,29	2.29	4 (18%)	27,45,45	2.60	4 (14%)
4	THM	B	702	-	13,18,18	0.44	0	16,26,26	2.28	3 (18%)
2	SO4	B	802	-	4,4,4	0.15	0	6,6,6	0.46	0
2	SO4	B	807	-	4,4,4	0.42	0	6,6,6	0.31	0
2	SO4	B	808	-	4,4,4	0.17	0	6,6,6	0.16	0
2	SO4	B	809	-	4,4,4	0.13	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	501	-	-	0/12/32/32	0/3/3/3
4	THM	A	701	-	-	0/2/18/18	0/2/2/2
2	SO4	A	801	-	-	0/0/0/0	0/0/0/0
2	SO4	A	803	-	-	0/0/0/0	0/0/0/0
2	SO4	A	804	-	-	0/0/0/0	0/0/0/0
2	SO4	A	805	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	806	-	-	0/0/0/0	0/0/0/0
3	ADP	B	502	-	-	0/12/32/32	0/3/3/3
4	THM	B	702	-	-	0/2/18/18	0/2/2/2
2	SO4	B	802	-	-	0/0/0/0	0/0/0/0
2	SO4	B	807	-	-	0/0/0/0	0/0/0/0
2	SO4	B	808	-	-	0/0/0/0	0/0/0/0
2	SO4	B	809	-	-	0/0/0/0	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	ADP	C6-N6	-2.69	1.26	1.34
3	A	501	ADP	C6-N6	-2.57	1.27	1.34
3	B	502	ADP	O4'-C1'	2.06	1.43	1.41
3	A	501	ADP	C2-N1	5.94	1.45	1.33
3	B	502	ADP	C2-N1	6.08	1.45	1.33
3	B	502	ADP	C2-N3	7.61	1.45	1.32
3	A	501	ADP	C2-N3	7.62	1.45	1.32

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	ADP	N3-C2-N1	-13.02	118.92	128.89
3	B	502	ADP	N3-C2-N1	-12.20	119.55	128.89
4	B	702	THM	C5-C4-N3	-4.64	119.97	125.14
4	A	701	THM	C5-C4-N3	-3.69	121.03	125.14
3	A	501	ADP	PA-O3A-PB	-2.49	124.31	132.67
3	B	502	ADP	PA-O3A-PB	-2.47	124.39	132.67
3	B	502	ADP	C4-C5-N7	-2.36	107.31	109.48
3	A	501	ADP	O3A-PA-O5'	-2.31	96.81	102.94
3	A	501	ADP	C2'-C1'-N9	-2.16	110.99	114.29
3	B	502	ADP	O3A-PA-O5'	-2.07	97.45	102.94
3	A	501	ADP	C4'-O4'-C1'	2.02	111.93	109.72
4	B	702	THM	C5M-C5-C4	2.33	123.05	120.05
4	A	701	THM	C4-N3-C2	6.42	120.80	115.25
4	B	702	THM	C4-N3-C2	7.18	121.45	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	702	THM	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	323/334 (96%)	0.08	17 (5%) 30 39	14, 23, 42, 53	0
1	B	330/334 (98%)	0.30	29 (8%) 12 17	15, 27, 49, 57	0
All	All	653/668 (97%)	0.19	46 (7%) 19 26	14, 25, 46, 57	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	89	LEU	6.7
1	B	20	SER	6.2
1	B	259	VAL	6.1
1	A	197	ALA	5.9
1	A	199	ILE	5.5
1	B	21	HIS	4.8
1	B	48	ALA	4.8
1	B	352	VAL	4.6
1	A	200	GLY	4.5
1	B	279	ASP	4.4
1	B	195	THR	4.3
1	B	199	ILE	4.2
1	A	352	VAL	3.9
1	B	90	ALA	3.8
1	B	198	ARG	3.6
1	B	333	PHE	3.3
1	B	85	GLN	3.3
1	A	259	VAL	3.2
1	A	195	THR	3.1
1	A	78	TYR	3.1
1	B	88	ASN	3.1
1	B	351	GLY	3.0
1	A	242	CYS	3.0
1	B	92	ASP	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	49	ALA	2.9
1	A	80	THR	2.9
1	B	22	MET	2.9
1	A	196	ARG	2.7
1	A	280	ARG	2.6
1	B	237	GLU	2.5
1	B	236	GLY	2.5
1	B	91	VAL	2.4
1	B	82	ASN	2.4
1	A	260	SER	2.3
1	B	274	GLN	2.3
1	A	198	ARG	2.3
1	B	30	LEU	2.3
1	A	70	GLU	2.1
1	B	191	ARG	2.1
1	B	146	VAL	2.1
1	B	242	CYS	2.1
1	B	311	THR	2.1
1	B	83	ARG	2.0
1	A	142	VAL	2.0
1	A	243	GLY	2.0
1	A	20	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ADP	B	502	27/27	0.94	0.17	0.54	22,40,42,42	0
2	SO4	B	802	5/5	0.96	0.13	-0.15	43,44,45,45	0
3	ADP	A	501	27/27	0.97	0.09	-0.44	15,25,28,29	0
4	THM	A	701	17/17	0.92	0.11	-0.60	20,27,33,35	0
2	SO4	A	801	5/5	0.98	0.10	-0.60	39,42,42,42	0
4	THM	B	702	17/17	0.95	0.10	-1.19	21,24,30,30	0
2	SO4	B	808	5/5	0.96	0.35	-	54,55,56,56	0
2	SO4	A	803	5/5	0.99	0.13	-	32,33,35,35	0
2	SO4	A	806	5/5	0.98	0.18	-	48,48,48,50	0
2	SO4	B	809	5/5	0.97	0.17	-	60,60,60,61	0
2	SO4	B	807	5/5	0.98	0.15	-	34,34,37,38	0
2	SO4	A	804	5/5	0.92	0.17	-	61,61,61,62	0
2	SO4	A	805	5/5	0.97	0.20	-	52,52,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.