



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:29 PM GMT

PDB ID : 1P7D  
Title : Crystal structure of the Lambda Integrase (residues 75-356) bound to DNA  
Authors : Aihara, H.; Kwon, H.J.; Nunes-Duby, S.E.; Landy, A.; Ellenberger, T.  
Deposited on : 2003-05-01  
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2552 (3.00-2.92)
Ramachandran outliers	100387	2468 (3.00-2.92)
Sidechain outliers	100360	2471 (3.00-2.92)
RSRZ outliers	91569	2201 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	13	
1	E	13	
2	D	26	
2	F	26	
3	A	283	
3	B	283	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*CP\*AP\*AP\*TP\*GP\*CP\*CP\*AP\*AP\*CP\*TP\*TP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	13	Total	C	N	O	P	0	0	0
			259	126	45	76	12			
1	E	13	Total	C	N	O	P	0	0	0
			259	126	45	76	12			

- Molecule 2 is a DNA chain called 26-MER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	26	Total	C	N	O	P	0	0	0
			541	257	103	155	26			
2	F	26	Total	C	N	O	P	0	0	0
			541	257	103	155	26			

- Molecule 3 is a protein called Integrase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	A	280	Total	C	N	O	P	S	0	0	0
			2161	1358	389	402	1	11			
3	B	280	Total	C	N	O	P	S	0	0	0
			2123	1332	383	397	1	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	MET	VAL	SEE REMARK 999	UNP P03700
A	342	PTR	TYR	MODIFIED RESIDUE	UNP P03700
B	74	MET	VAL	SEE REMARK 999	UNP P03700
B	342	PTR	TYR	MODIFIED RESIDUE	UNP P03700

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(\*CP\*AP\*AP\*TP\*GP\*CP\*CP\*AP\*AP\*CP\*TP\*TP\*T)-3'

Chain C: 





- Molecule 1: 5'-D(\*CP\*AP\*AP\*TP\*GP\*CP\*CP\*AP\*AP\*CP\*TP\*TP\*T)-3'

Chain E: 



- Molecule 2: 26-MER

Chain D: 




- Molecule 2: 26-MER


Chain F: 




- Molecule 3: Integrase

Chain A: 





● Molecule 3: Integrase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.02Å 106.16Å 73.17Å 90.00° 92.10° 90.00°	Depositor
Resolution (Å)	50.00 – 2.95 35.08 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.95) 99.3 (35.08-2.81)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.47 (at 2.81Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.231 , 0.259 0.230 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	57.6	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.0	EDS
Estimated twinning fraction	0.037 for h,-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 21931 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5884	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	C	0.49	0/289	0.85	0/443
1	E	0.51	0/289	0.84	0/443
2	D	0.52	0/608	0.87	0/936
2	F	0.51	0/608	0.86	0/936
3	A	0.38	0/2179	0.63	0/2931
3	B	0.38	0/2141	0.63	0/2888
All	All	0.42	0/6114	0.71	0/8577

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	E	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	10	DC	Sidechain
1	E	10	DC	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	259	0	148	17	0
1	E	259	0	148	9	0
2	D	541	0	294	17	0
2	F	541	0	294	24	0
3	A	2161	0	2165	78	0
3	B	2123	0	2075	81	0
All	All	5884	0	5124	210	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 210 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:21:DA:H2''	2:D:22:DG:H5''	1.08	1.06
3:A:328:GLN:HG2	3:A:343:ARG:HH21	1.31	0.95
3:B:328:GLN:HG2	3:B:343:ARG:HH21	1.31	0.94
3:B:345:ASP:OD2	3:B:349:GLU:HB2	1.68	0.93
2:D:21:DA:C2'	2:D:22:DG:H5''	1.97	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	275/283 (97%)	245 (89%)	19 (7%)	11 (4%)	4 18

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	B	275/283 (97%)	242 (88%)	22 (8%)	11 (4%)	4	18
All	All	550/566 (97%)	487 (88%)	41 (8%)	22 (4%)	4	18

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	115	PRO
3	A	340	SER
3	A	354	GLU
3	A	355	ILE
3	B	90	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	219/236 (93%)	214 (98%)	5 (2%)	58	86
3	B	208/236 (88%)	201 (97%)	7 (3%)	44	79
All	All	427/472 (90%)	415 (97%)	12 (3%)	51	83

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	97	LEU
3	B	109	ARG
3	B	324	ASP
3	A	345	ASP
3	B	179	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	328	GLN
3	B	94	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	B	328	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PTR	A	342	1,3	11,15,17	0.63	0	13,19,24	0.77	0
3	PTR	B	342	1,3	11,15,17	0.73	0	13,19,24	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PTR	A	342	1,3	-	0/6/10/13	0/1/1/1
3	PTR	B	342	1,3	-	0/6/10/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	13/13 (100%)	-0.04	0 100 100	23, 37, 72, 89	0
1	E	13/13 (100%)	0.01	0 100 100	25, 41, 81, 98	0
2	D	26/26 (100%)	0.22	0 100 100	17, 48, 98, 108	0
2	F	26/26 (100%)	0.38	1 (3%) 44 26	19, 54, 123, 143	0
3	A	279/283 (98%)	0.06	4 (1%) 78 59	20, 49, 69, 79	0
3	B	279/283 (98%)	0.07	9 (3%) 51 32	20, 49, 72, 82	0
All	All	636/644 (98%)	0.08	14 (2%) 65 44	17, 49, 73, 143	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	335	SER	3.5
3	B	162	THR	3.4
3	B	164	HIS	2.9
3	A	74	MET	2.6
3	A	172	LYS	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	PTR	B	342	15/17	0.97	0.20	-	27,34,45,45	0
3	PTR	A	342	15/17	0.98	0.19	-	30,34,47,47	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.