



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:29 PM GMT

PDB ID : 1P7Z  
Title : Crystal structure of the D181S variant of catalase HP11 from E. coli  
Authors : Chelikani, P.; Carpena, X.; Fita, I.; Loewen, P.C.  
Deposited on : 2003-05-06  
Resolution : 2.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	753	<div> <div>3%</div> <div>80% 15% . .</div> </div>
1	B	753	<div> <div>2%</div> <div>76% 18% . .</div> </div>
1	C	753	<div> <div>2%</div> <div>77% 17% . . .</div> </div>
1	D	753	<div> <div>2%</div> <div>76% 18% . .</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 25893 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase HPIL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	727	Total	C	N	O	S	0	1	0
			5747	3649	1005	1081	12			
1	B	727	Total	C	N	O	S	0	1	0
			5747	3649	1005	1081	12			
1	C	727	Total	C	N	O	S	0	1	0
			5747	3649	1005	1081	12			
1	D	727	Total	C	N	O	S	0	1	0
			5747	3649	1005	1081	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	181	SER	ASP	ENGINEERED	UNP P21179
B	181	SER	ASP	ENGINEERED	UNP P21179
C	181	SER	ASP	ENGINEERED	UNP P21179
D	181	SER	ASP	ENGINEERED	UNP P21179

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	739	Total O 739 739	0	0
3	B	610	Total O 610 610	0	0
3	C	674	Total O 674 674	0	0
3	D	710	Total O 710 710	0	0







• Molecule 1: Catalase HP1I



• Molecule 1: Catalase HP1I



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.34Å 132.88Å 121.45Å 90.00° 109.40° 90.00°	Depositor
Resolution (Å)	29.80 – 2.21 17.93 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.80-2.21) 99.1 (17.93-2.21)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.21 (at 2.21Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.144 , 0.211 0.142 , 0.201	Depositor DCC
$R_{free}$ test set	6953 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.528	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 62.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 138642 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	25893	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/5908	1.34	42/8032 (0.5%)
1	B	0.55	0/5908	1.35	44/8032 (0.5%)
1	C	0.55	0/5908	1.35	55/8032 (0.7%)
1	D	0.56	0/5908	1.42	56/8032 (0.7%)
All	All	0.55	0/23632	1.37	197/32128 (0.6%)

There are no bond length outliers.

The worst 5 of 197 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	D	612	ARG	CD-NE-CZ	30.99	166.98	123.60
1	B	37	ARG	CD-NE-CZ	18.81	149.94	123.60
1	A	542	ARG	NE-CZ-NH2	-16.78	111.91	120.30
1	C	445	ARG	NE-CZ-NH1	15.90	128.25	120.30
1	D	61	ARG	NE-CZ-NH2	13.58	127.09	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5747	0	5586	63	0
1	B	5747	0	5586	101	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	5747	0	5586	75	0
1	D	5747	0	5586	80	0
2	A	43	0	30	1	0
2	B	43	0	30	3	0
2	C	43	0	30	3	0
2	D	43	0	30	3	0
3	A	739	0	0	3	0
3	B	610	0	0	14	0
3	C	674	0	0	10	0
3	D	710	0	0	8	0
All	All	25893	0	22464	291	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:750:LYS:HD3	1:D:751:ILE:H	1.49	0.78
1:B:583:LYS:NZ	1:B:583:LYS:H	1.84	0.75
1:D:267:ARG:HG2	1:D:332:PRO:HB3	1.69	0.74
1:C:267:ARG:HG3	3:C:3862:HOH:O	1.89	0.72
1:B:710:ILE:HD13	1:B:718:ILE:HG13	1.70	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/753 (96%)	699 (96%)	21 (3%)	6 (1%)	24	21
1	B	726/753 (96%)	703 (97%)	18 (2%)	5 (1%)	26	25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	726/753 (96%)	705 (97%)	19 (3%)	2 (0%)	46	50
1	D	726/753 (96%)	698 (96%)	24 (3%)	4 (1%)	30	29
All	All	2904/3012 (96%)	2805 (97%)	82 (3%)	17 (1%)	30	29

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	709	LYS
1	B	28	SER
1	D	28	SER
1	D	750	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	613/636 (96%)	597 (97%)	16 (3%)	54	65
1	B	613/636 (96%)	589 (96%)	24 (4%)	39	47
1	C	613/636 (96%)	577 (94%)	36 (6%)	24	26
1	D	613/636 (96%)	581 (95%)	32 (5%)	29	32
All	All	2452/2544 (96%)	2344 (96%)	108 (4%)	35	41

5 of 108 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	252	ASN
1	C	571	LEU
1	D	616	LEU
1	C	375	LEU
1	C	459	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	459	ASN
1	C	556	GLN
1	D	629	HIS
1	C	507	HIS
1	C	629	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	760	1	30,50,50	2.51	8 (26%)	24,82,82	2.66	12 (50%)
2	HEM	B	760	1	30,50,50	2.52	10 (33%)	24,82,82	2.94	13 (54%)
2	HEM	C	760	1	30,50,50	2.41	6 (20%)	24,82,82	2.59	10 (41%)
2	HEM	D	760	1	30,50,50	2.59	10 (33%)	24,82,82	2.63	10 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	760	1	-	0/10/54/54	0/0/8/8
2	HEM	B	760	1	-	0/10/54/54	0/0/8/8
2	HEM	C	760	1	-	0/10/54/54	0/0/8/8
2	HEM	D	760	1	-	0/10/54/54	0/0/8/8

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	760	HEM	C3B-C4B	-7.40	1.45	1.51
2	A	760	HEM	C2D-C3D	-7.35	1.32	1.54
2	C	760	HEM	C2D-C3D	-7.34	1.32	1.54
2	B	760	HEM	C2D-C3D	-7.26	1.32	1.54
2	B	760	HEM	C3B-C4B	-7.17	1.45	1.51

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	760	HEM	C3C-CAC-CBC	-5.54	115.97	124.46
2	D	760	HEM	CAA-C2A-C1A	-4.21	122.44	127.01
2	D	760	HEM	C3C-CAC-CBC	-4.17	118.07	124.46
2	B	760	HEM	C3B-CAB-CBB	-4.09	118.19	124.46
2	A	760	HEM	C3C-CAC-CBC	-4.05	118.24	124.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	760	HEM	1	0
2	B	760	HEM	3	0
2	C	760	HEM	3	0
2	D	760	HEM	3	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	727/753 (96%)	-0.57	19 (2%) 59 59	10, 19, 41, 73	1 (0%)
1	B	727/753 (96%)	-0.44	18 (2%) 61 60	11, 20, 47, 65	1 (0%)
1	C	727/753 (96%)	-0.50	18 (2%) 61 60	11, 20, 44, 64	1 (0%)
1	D	727/753 (96%)	-0.51	14 (1%) 70 68	10, 19, 42, 65	1 (0%)
All	All	2908/3012 (96%)	-0.50	69 (2%) 62 61	10, 19, 44, 73	4 (0%)

The worst 5 of 69 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	ASP	7.7
1	C	27	ASP	6.5
1	A	28	SER	5.5
1	C	28	SER	5.0
1	A	712	ASP	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	HEM	B	760	43/43	0.97	0.10	0.53	2,12,14,15	0
2	HEM	D	760	43/43	0.98	0.09	0.16	2,12,15,17	0
2	HEM	A	760	43/43	0.98	0.08	-0.09	9,13,15,17	0
2	HEM	C	760	43/43	0.98	0.07	-0.48	11,14,17,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.