wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 05:03 AM GMT

PDB ID	:	2P7M
Title	:	Crystal structure of monoclinic form of genomically encoded fosfomycin resistance protein, FosX, from Listeria monocytogenes at pH 6.5
Authors	:	Fillgrove, K.L.; Pakhomova, S.; Schaab, M.; Newcomer, M.E.; Armstrong, R.N.
Deposited on	:	2007-03-20
Resolution	:	1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	133	66% 29% • 5%
1	B	133	71% 20% • 8%
1	C	133	81% 16% •
1	D	133	76% 18% 6%
1	E	133	75% 20% • •
1	F	133	70% 27% • •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyoxalase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	127	1044	669	174	197	4	0	0	0
1	B	122	981	632	156	188	5	0	2	0
1	C	129	1054	675	173	201	5	0	1	0
1	D	125	1031	659	171	196	5	0	2	0
1	E	128	1049	671	172	201	5	0	1	0
1	F	130	1065	680	177	203	5	0	1	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	46	46	46	0	0
2	B	35	35	35	0	0
2	C	51	51	51	0	0
2	D	45	45	45	0	0
2	E	66	66	66	0	0
2	F	28	28	28	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Glyoxalase family protein

Chain A:



- Molecule 1: Glyoxalase family protein

Chain B:

- Molecule 1: Glyoxalase family protein

Chain C:

- Molecule 1: Glyoxalase family protein

Chain D:

- Molecule 1: Glyoxalase family protein

Chain E:

- Molecule 1: Glyoxalase family protein

Chain F:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	169.43Å 69.69Å 84.87Å 90.00° 113.98° 90.00°	Depositor
Resolution (Å)	18.62 – 1.85	Depositor
% Data completeness (in resolution range)	81.9 (18.62-1.85)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.239 , 0.267	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6495	wwPDB-VP
Average B, all atoms ($Å^2$)	43.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.35	0/1064	0.58	0/1430
1	B	0.34	0/1007	0.56	0/1355
1	C	0.35	0/1079	0.56	0/1451
1	D	0.35	0/1059	0.60	0/1424
1	E	0.35	0/1074	0.57	0/1445
1	F	0.31	0/1090	0.56	0/1466
All	All	0.34	0/6373	0.57	0/8571

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1044	0	1021	35	0
1	B	981	0	941	19	0
1	C	1054	0	1032	17	0
1	D	1031	0	996	17	0
1	E	1049	0	1025	22	0
1	F	1065	0	1040	29	0
2	A	46	0	0	0	0
2	B	35	0	0	0	0
2	C	51	0	0	0	0
2	D	45	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	66	0	0	2	0
2	F	28	0	0	0	0
All	All	6495	0	6055	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:GLN:HE22	1:B:123:THR:HG22	1.24	0.99
1:F:47:PHE:HB2	1:F:54:ILE:HG23	1.64	0.78
1:F:99:ARG:HH12	1:F:105:ARG:HG2	1.49	0.76
1:A:52:LEU:HD11	1:B:5:LEU:HB2	1.71	0.73
1:C:13:LYS:NZ	1:F:63:GLN:HE22	1.88	0.72

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	123/133 (92%)	122 (99%)	1 (1%)	0	100 100
1	B	120/133 (90%)	115 (96%)	3 (2%)	2 (2%)	11 2
1	C	128/133 (96%)	123 (96%)	5 (4%)	0	100 100
1	D	123/133 (92%)	119 (97%)	3 (2%)	1 (1%)	24 9
1	E	127/133 (96%)	121 (95%)	6 (5%)	0	100 100
1	F	129/133 (97%)	124 (96%)	4 (3%)	1 (1%)	24 9
All	All	750/798 (94%)	724 (96%)	22 (3%)	4 (0%)	34 17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	129	LYS
1	D	34	SER
1	F	104	GLY
1	B	2	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/119 (94%)	110 (98%)	2 (2%)	66	52
1	B	105/119 (88%)	104 (99%)	1 (1%)	82	76
1	C	115/119 (97%)	114 (99%)	1 (1%)	84	79
1	D	111/119 (93%)	111 (100%)	0	100	100
1	E	115/119 (97%)	113 (98%)	2 (2%)	68	54
1	F	116/119 (98%)	115 (99%)	1 (1%)	84	79
All	All	674/714 (94%)	667 (99%)	7 (1%)	82	76

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	39	PHE
1	F	80	ASP
1	E	39	PHE
1	A	131	TYR
1	E	48	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	27	ASN
1	D	27	ASN
1	E	73	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	23	GLN
1	E	101	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates (i)

EDS was not executed - this section will therefore be empty.

6.4 Ligands (i)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers (i)

EDS was not executed - this section will therefore be empty.