



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:34 PM GMT

PDB ID : 4P7I
Title : Crystal structure of the Merlin FERM/DCAF1 complex
Authors : Wei, Z.; Li, Y.; Zhang, M.
Deposited on : 2014-03-27
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	<div> <div>2%</div> <div>84%</div> <div>9%</div> <div>7%</div> </div>
1	B	317	<div> <div>14%</div> <div>81%</div> <div>10%</div> <div>8%</div> </div>
2	C	67	<div> <div>7%</div> <div>13%</div> <div>9%</div> <div>78%</div> </div>
2	D	67	<div> <div>7%</div> <div>9%</div> <div>90%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Merlin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	0	0
			2458	1592	411	442	13			
1	B	292	Total	C	N	O	S	0	0	0
			2442	1581	408	440	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P46662
A	-2	PRO	-	expression tag	UNP P46662
A	-1	GLY	-	expression tag	UNP P46662
A	0	SER	-	expression tag	UNP P46662
B	-3	GLY	-	expression tag	UNP P46662
B	-2	PRO	-	expression tag	UNP P46662
B	-1	GLY	-	expression tag	UNP P46662
B	0	SER	-	expression tag	UNP P46662

- Molecule 2 is a protein called Protein VPRBP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	15	Total	C	N	O	0	0	0
			116	75	16	25			
2	D	7	Total	C	N	O	0	0	0
			54	35	8	11			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	1441	GLY	-	expression tag	UNP Q9Y4B6
C	1442	PRO	-	expression tag	UNP Q9Y4B6
C	1443	GLY	-	expression tag	UNP Q9Y4B6
C	1444	SER	-	expression tag	UNP Q9Y4B6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1445	GLU	-	expression tag	UNP Q9Y4B6
C	1446	PHE	-	expression tag	UNP Q9Y4B6
D	1441	GLY	-	expression tag	UNP Q9Y4B6
D	1442	PRO	-	expression tag	UNP Q9Y4B6
D	1443	GLY	-	expression tag	UNP Q9Y4B6
D	1444	SER	-	expression tag	UNP Q9Y4B6
D	1445	GLU	-	expression tag	UNP Q9Y4B6
D	1446	PHE	-	expression tag	UNP Q9Y4B6

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	24	Total	O	0	0
			24	24		
4	B	7	Total	O	0	0
			7	7		

GLY	I1501	GLY	D1500
PRO	I1502	PRO	
GLY	L1503	GLY	
SER	S1504	SER	
GLU	L1505	GLU	
PHE	H1506	PHE	
GLY	GLU	GLY	
GLU		GLU	
ASP		ASP	
GLY		GLY	
ASP		ASP	
ASN		ASN	
ASP		ASP	
PHE		PHE	
SER		SER	
PRO		PRO	
SER		SER	
ASP		ASP	
GLU		GLU	
GLU		GLU	
LEU		LEU	
ALA		ALA	
ASN		ASN	
LEU		LEU	
LEU		LEU	
GLU		GLU	
GLU		GLU	
GLY		GLY	
GLU		GLU	
ASP		ASP	
GLY		GLY	
GLU		GLU	
ASP		ASP	
ALA		ALA	
ASP		ASP	
GLU		GLU	
GLU		GLU	
VAL		VAL	
GLU		GLU	
LEU		LEU	
ILE		ILE	
LEU		LEU	
GLY		GLY	
ASP		ASP	
THR		THR	
ASP		ASP	
SER		SER	
SER		SER	
ASP		ASP	
ASN		ASN	
SER		SER	
ASP		ASP	
LEU		LEU	
GLU		GLU	
ASP		ASP	
D1500		D1500	

I1501	GLU
I1502	
L1503	
S1504	
L1505	
H1506	
GLU	

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	97.10Å 97.10Å 224.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.47 – 2.60 31.47 – 2.60	Depositor EDS
% Data completeness (in resolution range)	97.9 (31.47-2.60) 88.0 (31.47-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.224 , 0.256 0.213 , 0.248	Depositor DCC
R_{free} test set	1680 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	67.9	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 60.3	EDS
Estimated twinning fraction	0.033 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 37760 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5107	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2518	0.41	0/3396
1	B	0.23	0/2501	0.38	0/3374
2	C	0.22	0/114	0.45	0/152
2	D	0.20	0/53	0.41	0/71
All	All	0.25	0/5186	0.40	0/6993

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2458	0	2464	15	0
1	B	2442	0	2443	19	0
2	C	116	0	120	4	0
2	D	54	0	58	1	0
3	A	6	0	8	0	0
4	A	24	0	0	0	0
4	B	7	0	0	2	0
All	All	5107	0	5093	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:PRO:O	1:B:159:LYS:NZ	2.13	0.81
1:A:149:LYS:NZ	1:A:166:GLU:OE2	2.13	0.81
1:A:74:TRP:O	1:A:76:LYS:NZ	2.14	0.80
1:A:233:LEU:HB2	1:A:244:TYR:HB2	1.72	0.71
1:B:85:ASP:OD2	1:B:291:ARG:NH1	2.31	0.64
2:C:1487:GLY:N	2:C:1500:ASP:O	2.32	0.58
1:B:116:HIS:ND1	1:B:204:GLU:OE2	2.38	0.57
1:A:179:MET:HB3	1:A:183:MET:HB2	1.88	0.56
1:A:161:GLY:N	1:A:185:GLU:OE2	2.35	0.55
2:C:1482:VAL:HG13	2:C:1505:LEU:HD23	1.90	0.54
1:B:150:TYR:OH	1:B:166:GLU:OE2	2.19	0.54
1:B:233:LEU:HB2	1:B:244:TYR:HB2	1.91	0.52
1:A:100:PHE:CZ	1:A:302:GLY:HA3	2.44	0.52
2:D:1500:ASP:OD1	2:D:1500:ASP:N	2.43	0.52
1:B:220:ASN:ND2	4:B:406:HOH:O	2.42	0.52
1:A:136:GLU:OE1	1:A:187:ARG:NH2	2.40	0.51
1:B:240:GLY:HA2	1:B:258:TRP:CE2	2.48	0.49
1:A:240:GLY:HA2	1:A:258:TRP:CE2	2.48	0.48
1:A:301:ILE:HG12	2:C:1503:LEU:HD11	1.95	0.48
1:B:270:GLU:OE2	1:B:284:LYS:HD3	2.14	0.48
1:B:98:ALA:O	4:B:401:HOH:O	2.20	0.47
1:A:48:ASP:O	1:A:52:ARG:HG3	2.15	0.47
1:A:225:ARG:NH2	1:A:231:GLU:OE1	2.43	0.46
1:A:74:TRP:CD1	1:A:99:LYS:HE3	2.51	0.45
1:A:269:LYS:HD3	1:A:288:SER:HA	1.98	0.45
1:A:308:MET:HB3	2:C:1480:GLU:HA	1.98	0.45
1:B:160:ARG:HG2	1:B:182:GLU:OE2	2.17	0.45
1:B:219:VAL:HG22	1:B:237:ASP:HB3	2.00	0.44
1:B:100:PHE:CZ	1:B:302:GLY:HA3	2.54	0.43
1:B:306:LEU:O	1:B:310:ARG:HG3	2.18	0.43
1:B:43:GLY:HA3	1:B:76:LYS:O	2.19	0.43
1:B:38:GLU:HB2	1:B:41:TRP:CG	2.54	0.42
1:B:79:LYS:HE3	1:B:79:LYS:HB2	1.85	0.42
1:B:56:LEU:HD21	1:B:62:PHE:CD2	2.54	0.41
1:B:305:ASP:OD2	1:B:309:ARG:NE	2.49	0.41
1:B:154:ASP:HA	1:B:155:PRO:HD3	1.92	0.41
1:A:111:GLN:O	1:A:115:GLN:HG3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/317 (92%)	289 (99%)	3 (1%)	0	100	100
1	B	290/317 (92%)	278 (96%)	10 (3%)	2 (1%)	26	51
2	C	11/67 (16%)	11 (100%)	0	0	100	100
2	D	5/67 (8%)	5 (100%)	0	0	100	100
All	All	598/768 (78%)	583 (98%)	13 (2%)	2 (0%)	46	72

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	160	ARG
1	B	199	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/284 (94%)	266 (99%)	2 (1%)	88	96
1	B	266/284 (94%)	265 (100%)	1 (0%)	93	98
2	C	14/58 (24%)	14 (100%)	0	100	100
2	D	7/58 (12%)	7 (100%)	0	100	100
All	All	555/684 (81%)	552 (100%)	3 (0%)	92	98

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	GLU
1	A	281	ASP
1	B	41	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	401	-	5,5,5	0.29	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	401	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/317 (92%)	0.12	7 (2%) 62 56	38, 64, 117, 149	0
1	B	292/317 (92%)	0.85	44 (15%) 3 1	75, 118, 155, 195	0
2	C	15/67 (22%)	1.37	5 (33%) 0 0	65, 96, 132, 138	0
2	D	7/67 (10%)	3.71	5 (71%) 0 0	122, 139, 152, 166	0
All	All	608/768 (79%)	0.54	61 (10%) 9 6	38, 96, 149, 195	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	178	GLN	10.0
2	D	1506	ASN	7.4
2	D	1502	ILE	7.2
2	C	1506	ASN	5.6
1	B	35	PHE	5.5
1	B	279	LYS	5.4
2	C	1480	GLU	5.3
1	B	278	LYS	4.3
1	B	177	TYR	4.2
1	B	83	ASP	4.0
1	B	82	LEU	3.9
2	D	1501	ILE	3.9
1	B	40	LYS	3.8
1	A	177	TYR	3.8
1	B	87	SER	3.6
1	B	175	ASN	3.6
1	B	24	VAL	3.5
1	B	78	ASP	3.5
1	A	196	ARG	3.4
1	B	92	VAL	3.3
2	C	1486	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	39	MET	3.2
1	B	36	ASN	3.1
1	B	276	LEU	3.1
2	D	1500	ASP	3.1
1	B	179	MET	3.0
1	B	312	LYS	2.9
1	B	156	SER	2.9
1	B	68	ILE	2.9
1	B	42	LYS	2.9
1	B	290	LEU	2.9
1	B	232	LEU	2.8
1	B	90	GLU	2.7
1	B	280	ILE	2.7
1	B	37	CYS	2.7
1	A	69	LYS	2.7
1	B	89	GLU	2.7
1	A	178	GLN	2.7
1	B	110	VAL	2.7
1	B	31	ALA	2.6
1	B	226	ASN	2.5
1	B	157	VAL	2.5
2	D	1504	SER	2.5
1	A	312	LYS	2.5
1	B	41	TRP	2.4
1	B	199	ALA	2.4
1	B	23	THR	2.4
1	B	74	TRP	2.4
1	B	299	LEU	2.4
1	B	196	ARG	2.4
2	C	1500	ASP	2.4
1	B	140	LEU	2.3
1	B	22	PHE	2.3
1	B	277	ASP	2.3
1	B	63	GLY	2.2
1	B	311	ARG	2.2
2	C	1483	GLU	2.1
1	A	171	LYS	2.1
1	B	245	ASP	2.0
1	A	157	VAL	2.0
1	B	291	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	A	401	6/6	0.90	0.23	1.44	71,72,78,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.