



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:29 PM GMT

PDB ID : 1P8H
Title : BACTERIORHODOPSIN M1 INTERMEDIATE PRODUCED AT ROOM TEMPERATURE
Authors : Lanyi, J.K.
Deposited on : 2003-05-07
Resolution : 1.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2887 (1.54-1.50)
Ramachandran outliers	100387	2818 (1.54-1.50)
Sidechain outliers	100360	2816 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	1-A	249	
1	2-A	249	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LI1	1-A	601	X	-	-	-
3	LI1	1-A	611	X	-	-	-
3	LI1	1-A	613	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LI1	2-A	601	X	-	-	-
3	LI1	2-A	611	X	-	-	-
3	LI1	2-A	613	X	-	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4145 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriorhodopsin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1-A	222	Total	C	N	O	S	0	0	0
			1720	1159	262	290	9			
1	2-A	222	Total	C	N	O	S	0	0	0
			1719	1159	261	290	9			

- Molecule 2 is RETINAL (three-letter code: RET) (formula: C₂₀H₂₈O).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1-A	1	Total	C	0	0
			20	20		
2	2-A	1	Total	C	0	0
			20	20		

- Molecule 3 is 1-[2,6,10,14-TETRAMETHYL-HEXADECAN-16-YL]-2-[2,10,14-TRIMETHYL-HEXADECAN-16-YL]GLYCEROL (three-letter code: LI1) (formula: C₄₂H₈₆O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	1-A	1	Total C O 32 29 3	0	0
3	2-A	1	Total C O 32 29 3	0	0
3	1-A	1	Total C O 40 37 3	0	0
3	2-A	1	Total C O 40 37 3	0	0
3	1-A	1	Total C 18 18	0	0
3	2-A	1	Total C 18 18	0	0
3	1-A	1	Total C 16 16	0	0
3	2-A	1	Total C 16 16	0	0
3	1-A	1	Total C 8 8	0	0
3	2-A	1	Total C 8 8	0	0
3	1-A	1	Total C 8 8	0	0
3	2-A	1	Total C 8 8	0	0
3	1-A	1	Total C O 38 35 3	0	0
3	2-A	1	Total C O 38 35 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	1-A	1	Total C 18 18	0	0
3	2-A	1	Total C 18 18	0	0
3	1-A	1	Total C 16 16	0	0
3	2-A	1	Total C 16 16	0	0
3	1-A	1	Total C O 40 37 3	0	0
3	2-A	1	Total C O 40 37 3	0	0
3	1-A	1	Total C 17 17	0	0
3	2-A	1	Total C 17 17	0	0
3	1-A	1	Total C 18 18	0	0
3	2-A	1	Total C 18 18	0	0
3	1-A	1	Total C 13 13	0	0
3	2-A	1	Total C 13 13	0	0

- Molecule 4 is 2,10,23-TRIMETHYL-TETRACOSANE (three-letter code: SQU) (formula: C₂₇H₅₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	1-A	1	Total C 27 27	0	0
4	2-A	1	Total C 27 27	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	1-A	25	Total O 25 25	0	0
5	2-A	23	Total O 23 23	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Bacteriorhodopsin



• Molecule 1: Bacteriorhodopsin



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	60.91Å 60.91Å 108.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 1.52	Depositor
% Data completeness (in resolution range)	92.1 (25.00-1.52)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.175 , 0.207	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4145	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SQU, RET, LI1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1-A	0.42	0/1767	1.19	7/2413 (0.3%)
1	2-A	0.70	0/1765	1.51	17/2408 (0.7%)
All	All	0.58	0/3532	1.36	24/4821 (0.5%)

There are no bond length outliers.

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1-A	216	LYS	CB-CG-CD	12.97	145.31	111.60
1	2-A	175	ARG	NE-CZ-NH1	12.08	126.34	120.30
1	1-A	82	ARG	NE-CZ-NH1	-8.42	116.09	120.30
1	1-A	216	LYS	CA-CB-CG	8.14	131.31	113.40
1	2-A	175	ARG	NE-CZ-NH2	-7.87	116.37	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	1720	0	1776	100	0
1	2-A	1719	0	1774	54	0
2	1-A	20	0	27	2	0
2	2-A	20	0	27	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	1-A	282	0	452	55	0
3	2-A	282	0	452	62	0
4	1-A	27	0	53	7	0
4	2-A	27	0	53	7	0
5	1-A	25	0	0	10	0
5	2-A	23	0	0	1	0
All	All	4145	0	4614	229	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:609:LI1:C26	3:A:609:LI1:C25	1.74	1.60
3:A:609:LI1:C25	3:A:609:LI1:C26	1.74	1.60
1:A:164:ARG:HG3	1:A:166:GLU:HG3	1.46	0.97
1:A:19:LEU:C	1:A:20:MET:CA	2.37	0.92
3:A:609:LI1:C25	3:A:609:LI1:C27	2.52	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1-A	218/249 (88%)	208 (95%)	9 (4%)	1 (0%)	34	10
1	2-A	216/249 (87%)	214 (99%)	2 (1%)	0	100	100
All	All	434/498 (87%)	422 (97%)	11 (2%)	1 (0%)	52	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1-A	198	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1-A	178/195 (91%)	160 (90%)	18 (10%)	9	0
1	2-A	177/195 (91%)	168 (95%)	9 (5%)	29	4
All	All	355/390 (91%)	328 (92%)	27 (8%)	16	1

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1-A	186	PRO
1	1-A	202	ASN
1	2-A	164	ARG
1	1-A	191	ILE
1	1-A	74	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	1-A	76	ASN
1	1-A	202	ASN
1	2-A	105	GLN
1	2-A	202	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	RET	1-A	301	1	19,20,21	0.88	0	27,27,28	5.41	13 (48%)
3	LI1	1-A	601	-	31,31,44	1.08	2 (6%)	33,33,51	1.12	2 (6%)
3	LI1	1-A	602	-	39,39,44	0.97	2 (5%)	39,43,51	1.36	4 (10%)
3	LI1	1-A	603	-	17,17,44	1.03	1 (5%)	18,18,51	1.13	1 (5%)
3	LI1	1-A	604	-	15,15,44	1.11	1 (6%)	14,14,51	1.20	2 (14%)
3	LI1	1-A	605	-	7,7,44	0.46	0	6,6,51	0.66	0
3	LI1	1-A	606	-	7,7,44	0.44	0	6,6,51	0.55	0
3	LI1	1-A	607	-	37,37,44	1.00	2 (5%)	36,38,51	1.01	2 (5%)
3	LI1	1-A	608	-	17,17,44	1.03	1 (5%)	18,18,51	1.29	1 (5%)
3	LI1	1-A	609	-	15,15,44	1.13	1 (6%)	14,14,51	1.17	2 (14%)
3	LI1	1-A	610	-	39,39,44	1.02	2 (5%)	41,41,51	1.12	2 (4%)
3	LI1	1-A	611	-	16,16,44	0.97	1 (6%)	17,18,51	2.22	5 (29%)
3	LI1	1-A	612	-	17,17,44	1.04	1 (5%)	18,18,51	1.15	2 (11%)
3	LI1	1-A	613	-	12,12,44	0.44	0	12,12,51	0.94	0
4	SQU	1-A	701	-	26,26,26	1.35	6 (23%)	28,28,28	1.72	9 (32%)
2	RET	2-A	301	1	19,20,21	1.03	2 (10%)	27,27,28	1.98	11 (40%)
3	LI1	2-A	601	-	31,31,44	1.08	2 (6%)	33,33,51	1.12	2 (6%)
3	LI1	2-A	602	-	39,39,44	0.97	2 (5%)	39,43,51	1.36	4 (10%)
3	LI1	2-A	603	-	17,17,44	1.03	1 (5%)	18,18,51	1.13	1 (5%)
3	LI1	2-A	604	-	15,15,44	1.11	1 (6%)	14,14,51	1.20	2 (14%)
3	LI1	2-A	605	-	7,7,44	0.46	0	6,6,51	0.66	0
3	LI1	2-A	606	-	7,7,44	0.44	0	6,6,51	0.55	0
3	LI1	2-A	607	-	37,37,44	1.00	2 (5%)	36,38,51	1.01	2 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	LI1	2-A	608	-	17,17,44	1.03	1 (5%)	18,18,51	1.29	1 (5%)
3	LI1	2-A	609	-	15,15,44	1.13	1 (6%)	14,14,51	1.17	2 (14%)
3	LI1	2-A	610	-	39,39,44	1.02	2 (5%)	41,41,51	1.12	2 (4%)
3	LI1	2-A	611	-	16,16,44	0.97	1 (6%)	17,18,51	2.22	5 (29%)
3	LI1	2-A	612	-	17,17,44	1.04	1 (5%)	18,18,51	1.15	2 (11%)
3	LI1	2-A	613	-	12,12,44	0.44	0	12,12,51	0.94	0
4	SQU	2-A	701	-	26,26,26	1.35	6 (23%)	28,28,28	1.72	9 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	1-A	301	1	-	0/13/30/31	0/1/1/1
3	LI1	1-A	601	-	2/2/3/8	0/32/32/49	0/0/0/0
3	LI1	1-A	602	-	-	0/42/42/49	0/0/0/0
3	LI1	1-A	603	-	-	0/16/16/49	0/0/0/0
3	LI1	1-A	604	-	-	0/13/13/49	0/0/0/0
3	LI1	1-A	605	-	-	0/5/5/49	0/0/0/0
3	LI1	1-A	606	-	-	0/5/5/49	0/0/0/0
3	LI1	1-A	607	-	-	0/38/38/49	0/0/0/0
3	LI1	1-A	608	-	-	0/16/16/49	0/0/0/0
3	LI1	1-A	609	-	-	0/13/13/49	0/0/0/0
3	LI1	1-A	610	-	-	0/39/39/49	0/0/0/0
3	LI1	1-A	611	-	2/2/3/8	0/17/17/49	0/0/0/0
3	LI1	1-A	612	-	-	0/16/16/49	0/0/0/0
3	LI1	1-A	613	-	1/1/1/8	0/11/11/49	0/0/0/0
4	SQU	1-A	701	-	-	0/25/25/25	0/0/0/0
2	RET	2-A	301	1	-	0/13/30/31	0/1/1/1
3	LI1	2-A	601	-	2/2/3/8	0/32/32/49	0/0/0/0
3	LI1	2-A	602	-	-	0/42/42/49	0/0/0/0
3	LI1	2-A	603	-	-	0/16/16/49	0/0/0/0
3	LI1	2-A	604	-	-	0/13/13/49	0/0/0/0
3	LI1	2-A	605	-	-	0/5/5/49	0/0/0/0
3	LI1	2-A	606	-	-	0/5/5/49	0/0/0/0
3	LI1	2-A	607	-	-	0/38/38/49	0/0/0/0
3	LI1	2-A	608	-	-	0/16/16/49	0/0/0/0
3	LI1	2-A	609	-	-	0/13/13/49	0/0/0/0
3	LI1	2-A	610	-	-	0/39/39/49	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LI1	2-A	611	-	2/2/3/8	0/17/17/49	0/0/0/0
3	LI1	2-A	612	-	-	0/16/16/49	0/0/0/0
3	LI1	2-A	613	-	1/1/1/8	0/11/11/49	0/0/0/0
4	SQU	2-A	701	-	-	0/25/25/25	0/0/0/0

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	2-A	701	SQU	C14-C12	-3.31	1.35	1.52
4	1-A	701	SQU	C14-C12	-3.31	1.35	1.52
4	2-A	701	SQU	C23-C22	-2.98	1.34	1.51
4	1-A	701	SQU	C23-C22	-2.98	1.34	1.51
4	2-A	701	SQU	C18-C17	-2.85	1.35	1.51

The worst 5 of 88 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	1-A	301	RET	C20-C13-C14	-11.87	86.95	123.36
2	1-A	301	RET	C19-C9-C10	-7.72	111.50	122.90
2	1-A	301	RET	C11-C12-C13	-7.52	104.19	126.32
2	1-A	301	RET	C7-C6-C5	-7.14	105.02	121.37
2	2-A	301	RET	C11-C10-C9	-5.46	119.32	127.20

5 of 10 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	1-A	613	LI1	C18
3	1-A	601	LI1	C2
3	1-A	601	LI1	C18
3	2-A	611	LI1	C13
3	2-A	611	LI1	C23

There are no torsion outliers.

There are no ring outliers.

27 monomers are involved in 135 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	1-A	301	RET	2	0
3	1-A	601	LI1	3	0
3	1-A	602	LI1	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	1-A	603	LI1	3	0
3	1-A	604	LI1	1	0
3	1-A	605	LI1	1	0
3	1-A	606	LI1	2	0
3	1-A	607	LI1	5	0
3	1-A	609	LI1	3	0
3	1-A	610	LI1	18	0
3	1-A	611	LI1	2	0
3	1-A	612	LI1	2	0
4	1-A	701	SQU	7	0
2	2-A	301	RET	2	0
3	2-A	601	LI1	3	0
3	2-A	602	LI1	19	0
3	2-A	603	LI1	5	0
3	2-A	604	LI1	1	0
3	2-A	605	LI1	1	0
3	2-A	606	LI1	2	0
3	2-A	607	LI1	3	0
3	2-A	608	LI1	1	0
3	2-A	609	LI1	3	0
3	2-A	610	LI1	17	0
3	2-A	611	LI1	2	0
3	2-A	612	LI1	6	0
4	2-A	701	SQU	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.